Abstracts of the Joint Annual Meeting of the ASHS North Central Region and Western Canadian Society for Horticulture

Winnipeg, Manitoba, Canada

11-15 July 1989

APPLICATION OF SIMULATION MODELLING TO HORTICULTURE

Dennis D. Schulte*, George E. Meyer, Jay B. Fitzgerald and Kevin G. Power, Department of Agricultural Engineering and Horticulture, University of Nebraska, Lincoln, NE 68583

This paper is a forward-oriented review of computer-based simulation models. It focusses on applications to horticultural crops in nurseries and greenhouses. Highlights of systems simulation models for horticultural enterprises are presented. The paper closes with an outline of needs for future simulation models for horticultural crops and trends in computer science and engineering which will facilitate useful applications for the industry.

GRHEAT GREENHOUSE HEATING/COOLING ANALYSIS AND DESIGN MODEL

Kevin C. Power*, George E. Meyer, Dennis Schulte, and Jay B. Fitzgerald, Departments of Agricultural Engineering and Horticulture, University of Nebraska, Lincoln, NE 68583

A microcomputer program has been developed in a spreadsheet to perform heat/mass balance on several greenhouse architectural designs. Databases containing construction material and crop requirement data are automatically accessed for engineering data used for the heat/mass balance calculations. Output from the system includes: design heat gain/loss loads, required ventilation rates, evaporative cooler design and greenhouse orientation comparisons.

PLANT SPACING AND NITROGEN INFLUENCE ON STRAWBERRY GROWTH AND FRUITING

Elden J. Stang* and Gavin G. Weis
Department of Horticulture, University of Wisconsin-Madison, Madison, WI 53706

'Raritan' and 'Guardian' strawberry were grown in the matted row system with controlled plant densities of 1, 2, 3, 4 or 5 plants/0.09m² for comparison to a non-thinned matted row averaging 9 plants/0.09m². Nitrogen treatments were superimposed on plant spacings at 3 week intervals in preharvest and postharvest applications. Total seasonal available N was 0, 36, 54 and 76 kg/ha. Fruit yield per plant decreased as plant population increased. Berry size declined with increased plant population but number of fruit per plant was not influenced. For both cultivars, plant populations of 4 to 5 plants/0.09m² resulted in maximum fruit yield. Number of branch crowns for all treatments was 2.5-3.5/plant in the second growing season. Branch crown numbers were reduced with higher plant populations. N effects were independent of plant population effects and did not compensate for lower yields at low plant populations in more or larger berries. Optimum water management may be more important than N fertilizer in determining strawberry plant growth and yield.

ROOT INFECTION BY MYCORRHIZAL FUNGI REDUCES THE SOIL PHOSPHORUS REQUIREMENTS OF SOME HORTICULTURAL CROPS

Doug R. Waterer* and Robert R. Colman, University of Saskatchewan, Saskatoon, Sask. S7N 0W0

The influence of root colonization by vesicular-arbuscular mycorrhizal fungi (VAMF) on the growth, yields, tissue phosphorus (P) levels and soil P requirements of lettuce, onion, leek, garlic, bell pepper, and Saskatoon berry were studied in field and greenhouse experiments. Colonization by a mixture of VAMF species or by pure isolates of the VAMF *Glomus aggregatum* either had no effect or improved growth and yields of all species tested. Colonization improved the P-uptake efficiency of the host, reducing their requirements for fertilizer P. Onion, leek, garlic and bell pepper were more responsive to VAMF infection than lettuce or Saskatoon berry.

ECONOMIC ANALYSIS OF GREENHOUSE ENERGY CONSERVATION DESIGNS

Kevin C. Power*, George E. Meyer, Dennis Schulte, and Jay B. Fitzgerald, Departments of Agricultural Engineering and Horticulture, University of Nebraska, Lincoln, NE 68583

The Greenhouse Heating/Cooling Analysis and Design model (GRHEAT) utilizes annual heat requirement data to compare various energy-saving alternative greenhouse designs. GRHEAT includes an economic model that conducts a present value method of investment analysis for energy systems to compare various greenhouse designs. A statistical analysis of the heat requirement estimations from GRHEAT is conducted to validate the level of accuracy of the simulation results.

A GENETIC ANALYSIS OF SELF-INCOMPATIBILITY IN *POTENTILLA FRUTICOSA* L.

Bert L. Innes and Louis M. Lenz, Department of Plant Science, University of Manitoba, Winnipeg, Manitoba R3T 2N2

The inheritance of self-incompatibility in *Potentilla fruticosa* was investigated by intercrossing F₁ hybrids between two self-incompatible parents in a complete diallel design. Self-incompatibility was determined to be controlled by a monofactorial system operating gametophytically in the pollen. Pollen of *P. fruticosa* was found to be binucleate at anthesis, which is consistent with a gametophytic self-incompatibility system. Anomalies were observed in the segregation of extra petals in a series of backcrosses designed to study the inheritance of extra petals. These results could be explained by the linkage of the self-incompatibility locus to the modifier locus influencing petal number.

THE EFFECT OF POLYETHYLENE SOIL MULCH & ROW TUNNELS ON THE YIELD OF BELL PEPPER.

Richard St. Pierre, Todd Kabaluk and David Schmitz*, University of Saskatchewan, Saskatoon, Sask. S7N 0W0

The influence of environmental modification through the use of plastic mulches and tunnels on the total and marketable fruit yields and plant weights of 3 cultivars of bell pepper were investigated. Early harvests indicate few yield differences between the mulched treatments and the non-mulched control. In late harvests black plastic mulch with perforated plastic tunnels, and plastic mulch alone increased fruit numbers and fruit weights relative to the non-mulched control. Tunnels with mulch increased yields relative to mulch alone.

THE EFFECTS OF BENZYLAMINOPURINE AND CHLORFLURENOL ON SEEDLING MORPHOLOGY OF FOUR CONIFER SPECIES

Gerald A. Tuskan, Emile J. Poisson and Wayne A. Sargent*, Department of Horticulture and Forestry, North Dakota State University, Fargo, ND 58105

Benzylaminopurine and chlorflurenol were applied to 2-1 nursery stock of Scots pine and ponderosa pine and 2-0 nursery stock Colorado blue spruce and Black Hills spruce to determine if crown morphology was influenced by varying combinations of the two plant growth regulators. Four levels of benzylaminopurine, 0, 250, 750 and 1250 ppm, and two chlorflurenol levels, 0 and 1% (v/v) were tested. Morphological response to treatments was significantly enhanced when treatments were applied to open, elongating buds. Benzylaminopurine significantly increased bud and shoot formation, while chlorflurenol significantly reduced height and increased branch length at species dependent concentrations. The two plant growth regulators lacked positive synergistic effects.

ELECTRONIC VIDEO INSPECTION OF GROWING GREENHOUSE CROPS
W.W. Troyer*, G. E. Meyer and J. B. Fitzgerald, Departments
of Agricultural Engineering and Horticulture, University of
Nebraska, Lincoln, NE 68583

Spectral reflectance and electronic color image analyses were performed on the foliage of growing poinsettia plants in a greenhouse. The plants were subjected to high, moderate, and zero nitrogen treatments. Plants deficient in nitrogen showed increased visible reflectance and decreased near-Infrared reflectance and color image components of higher red and green intensity.

AN EVALUATION OF THE HORTICULTURE PROGRAM, PSYCHIATRIC
WARD, UNIVERSITY HOSPITAL. Sara Williams, University
of Saskatchewan, Saskatoon, Sask. S7N 0W0

A therapeutic horticulture program developed for patients in a short-term psychiatric facility was evaluated to determine the extent to which the goals of the program were being met. Participation fostered both social interaction and cooperative activities. 75% of the patients perceived the group to be both enjoyable and relaxing. 75% took responsibility for the care of their plants. Resource material was requested by less than 30%. The majority of the patients perceived the program as beneficial and felt satisfaction in their accomplishment. Previous gardening experience did not seem to influence the response distribution of perceived benefit or enjoyment of the program as much as it did the patient's decision to join the group initially.

CARROT CULTIVAR TOLERANCE TO SCLEROTINIA INFECTION, M.K. Pritchard* and J.E. Finlayson, Dept. of Plant Science,
University of Manitoba, Winnipeg, Canada R3T 2N2.

Five carrot cultivars were assessed for their tolerance to infection by *Sclerotinia sclerotiorum*. After infection by mycelium in the field, Six Pak II had more diseased roots than Dess Dan after 3 months storage at 5-6°C. When a conductivity assay measuring electrolyte leakage from carrot root discs after infection by the organism was used, there was more electrolyte leakage from discs of Six Pak II than from Dess Dan. A significant polynomial regression was obtained for leakage response to temperature between 2 and 20°C for all five cultivars. Resistance of root tissue to cellular disruption by *Sclerotinia* may play a role in the varying degrees of susceptibility of carrot cultivars to the disease.

PLANT PRODUCTION COST ACCOUNTING/MANAGEMENT (PPCAM) SYSTEM
Kevin C. Power*, George E. Meyer, and Jay Fitzgerald,
Departments of Agricultural Engineering and Horticulture,
University of Nebraska, Lincoln, NE 68583

A microcomputer program has been developed to keep records on energy, labor, product pricing, and revenue. The program manages plant production data enabling the grower to optimize production and profits. The grower can use the program to determine how much it costs to produce individual plants, to ascertain how much labor is costing and where to reallocate employees. Advertising and other indirect costs can be included to determine where the grower's money is spent on a per-plant or per-square foot basis.

GROWING ARBORVITAE CULTIVARS FROM SEED

A. A. Bee, Department of Horticulture and Forestry, North
Dakota State University, Fargo, ND 58105

Seeds from arborvitae, *Thuja occidentalis* L., cultivars were grown to determine phenotypic variations which might be useable as new cultivars. Seeds were germinated and grown under greenhouse and lath house condition. Many seedlings from isolated plants of *T. occidentalis* 'Hetz Midget' and 'Holmstrup' were similar or identical to the parent plant (98% and 58%, respectively). Seedlings grown from seeds of plants in a mixed planting produced seedlings similar to the seed parent as follows: 'Hetz Midget' 100%, 'Minima' 78%, 'Little Giant' 0%, 'Sherwood Moss' 11%, 'Spiralis' 24%, 'Hoseri' 36%; cultivars that produce a high percentage of true to type seedling could be propagated sexually.

A SEED SELECTION SYSTEM FOR VIGOROUS TRUE POTATO SEEDS
Vergel C. Concibido*, Department of Horticulture, North
Dakota State University, Fargo, ND 58105

In developing countries, the use of True Potato Seed (TPS) technology offers some potential economic and biological advantages compared to using seed tubers. One problem with using TPS technology is the lack of seedling vigor, poor

germination and non-uniformity of seedlings derived from TPS. Open-pollinated true seeds were divided into three classes based on differences in embryo morphology. Highly significant differences were observed among these seed classes for 100-seed weights, percent germination, rate of germination, 10-seedling weights and percent seedlings with leaves unfurled, ten days after sowing. Seed classes were quantified in terms of 100-seed weight with Seed class 1 having the lowest weight and Seed class 3 the highest; A significantly high *r* value (0.99) was obtained between seed classes and 10-seedling weight. High *r* values (0.93) and (0.94) were also obtained between seed classes and percent germination and percent seedlings with unfurled leaves, respectively. Results indicate that it may be possible to use these differences on embryo morphology as a basis for selecting vigorous true potato seeds.

EVALUATION OF LONICERA FOR SUSCEPTIBILITY TO HONEYSUCKLE
APHID, *HYADAPHIS TATARICAE* (AIZENBERG), WINTER HARDINESS AND
LANDSCAPE USE.

Dale E. Herman* and Lawrence J. Chaput, Dept. of Horticulture
and Forestry, North Dakota State Univ., Fargo, ND 58105

One-hundred honeysuckle taxa were evaluated and/or reviewed in the literature for relative susceptibility to honeysuckle aphid. Degree of susceptibility included 4 categories: 1) Apparent Resistance, 2) Lightly Susceptible, 3) Susceptible and 4) Highly Susceptible. Although discrepancies appeared in levels of reported susceptibility, a vast majority of taxa resulted in a clear-cut picture of susceptibility or resistance. Approximately 50% of the taxa reviewed are classified as having apparent resistance. The latter group formed the basis from which recommended taxa are derived. Further, this group was evaluated for winter hardiness and compared with hardiness zone recommendations in the literature. Honeysuckle taxa are categorized according to landscape characteristics for recommendation in USDA hardiness Zones 2 through 5. Landscape categories included 1) Highly Recommended, 2) Very Acceptable, 3) Fairly Acceptable and 4) Undesirable. Only 12 taxa are listed as highly recommended or very acceptable. Aphid-resistant taxa for potential replacements in conservation plantings are listed.

SEGREGATION FOR RESISTANCE TO HOLLOW HEART IN POTATOES

S.H. Jansky* and D.M. Thompson, Department of Horticulture
and Forestry, North Dakota State University, Fargo, ND 58105

Hollow heart is a physiological disorder that affects potato tuber quality. Although environmental factors influence the expression of hollow heart, there are genetic differences in expression among clones. Nine families from crosses between resistant and susceptible clones were grown in a replicated trial at Grand Forks, ND, in 1988. Five of these families were also included under irrigated conditions at Oakes, ND. Non-significant differences for hollow heart resistance were present among families grown at Grand Forks, probably because hot, dry growing conditions inhibited hollow heart expression. Significant differences were detected among families at Oakes. The number of clones/family with hollow heart was highest in a susceptible × resistant and lowest in a resistant × resistant cross. Clones in families with high susceptibility tended to express hollow heart more severely than those in less susceptible families. Resistant × resistant crosses produced some susceptible progeny, suggesting that resistance may be partially dominant.

RESISTANCE TO THE COLORADO POTATO BEETLE AND GREEN PEACH
APHID IN INTERSPECIFIC *SOLANUM* HYBRIDS.

Jim Mooney* and Shelley H. Jansky, Department of Horticulture
and Forestry, North Dakota State University, Fargo, ND 58105

Resistance to the Colorado potato beetle (CPB) and green peach aphid (GPA) would be valuable if it could be effectively transferred from wild potato species to the cultivated potato. Eighteen diploid interspecific hybrids have been developed using *Solanum tuberosum* Gp. *Tuberosum* haploids ($2n = 2x = 24$) and the diploid wild species *S. berthaultii* (ber), *S. chacoense* (CHC), *S. jamesii* (jam), and *S. tarijense* (tar). Twenty-five genotypes per family were screened for resistance to CPB and GPA. Feeding trials were carried out on intact leaves. The degree of resistance to CPB was determined by the stage of instar development and weight of larvae after a four day feeding period; resistance to GPA was evaluated by aphid reproduction and survival after a fifteen day feeding period. Highly CPB or GPA resistant clones, compared to 'Norgold Russet', have been identified thus far. Some clones express high levels of resistance to both CPB and GPA. Crosses between resistant clones and *S. tuberosum* will be carried out at the diploid level in an attempt to combine resistance with good tuberization qualities.