Literature Cited

- Boxus, P. H., M. Ouoirin, and J.M. Laine. 1977. Large scale propagation of strawberry plants from tissue culture. p. 130-143. In: J. Reinert and Y.P.S. Bajaj (eds.). Applied and fundamental aspects of plant cell, tissue and organ culture. Springer-Verlag, New York.
- Damiano, C. 1980. Strawberry micropropagation. Proc. Conf. on Nursery Production of Fruit Plants—Through Tissue Culture Applications and Feasibility. USDA-SEA, Agr. Res. Results ARR-NE-11. p. 11-22.
- Guttridge, C.G. and S. Bright. 1978. Accelerating and synchronizing germination of strawberry seeds by osmotic pre-treatments. Euphytica 27:843-848.
- Jones, O. P., B.J. Wailer, and M.G. Beech. 1988. The production of strawberry plants from callus cultures. Plant Cell Tissue Org. Cult. 12:235-241.
- Liu, Z.R. and J.C. Sanford. 1988. Plant regeneration by organogenesis from strawberry leaf and runner tissue. HortScience 23:1056-1059.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassay with tobacco tissue cultures. Physiol. Plant. 15:473-497.
- Scott, D.H. and A.D. Draper. 1967. Light in relation to seed germination of blueberries, strawberries and *Rubus*. HortScience 2:107-108.
- Scott, D.H. and F.J. Lawrence. 1975. Strawberries. p. 71-97. In: J. Janick and J.N. Moore (eds.). Advances in fruit breeding. Purdue Univ. Press, West Lafayette, Ind.
- Shoemaker, N. P., H.J. Swartz, and G.J. Galletta. 1985. Cultivar dependent variation in pathogen resistance due to tissue culture-propagation of strawberries. HortScience 20:253-254.
- Swartz, H. J., G.J. Galletta, and R.H. Zimmerman. 1981. Field performance and phenotypic stability of tissue culture propagated strawberries. J. Amer. Soc. Hort. Sci. 106:667-673.
- Wang, D., W.P. Wergin, and R.H. Zimmerman. 1984. Somatic embryogenesis and plant regeneration from immature embryos of strawberry. HortScience 19:71-72.
- Wilson, D., A. Goodall, and J. Reeves. 1973. An improved technique for the germination of strawberry seeds. Euphytica 12:362-366.
- Yurgalevitch, C. M., H.W. Janes, and C. Chin. 1985. Somaclonal variation in Fragaria. HortScience 20:450. (Abstr.).

HORTSCIENCE 25(5):571-572. 1990.

Callus Induction and Plant Regeneration of American Ginseng

Andrew S. Wang¹

Sungene Technologies Cop, 2050 Concourse Drive, San Jose, CA 95131

Additional index words. *Panax quinquefolium*, tissue culture

Abstract. Friable embryogenic callus of American ginseng (*Panax quinquefolium* L.) was induced from root pith on Murashige and Skoog medium supplemented with 2 mg 2,4-D and 1 mg KIN/liter. Optimal callus growth occurred on medium containing 1.5 mg dicamba/liter. Plants were regenerated on MS medium supplemented with various concentrations of plant growth regulators (PGRs); the best PGR combination was 0.5 mg IBA and 0.1 mg NAA/liter. Chemical names used: (2,4-dichlorophenoxy) acetic acid (2,4-D); 3,6-dichloro-o-anisic acid (dicamba); 6-benzylaminopurine (BA); gibberellic acid (GA); indole-3-butyric acid (IBA); kinetin (KIN); and naphthaleneacetic acid (NAA).

The root of ginseng (*Panax* spp.), cultivated in northern China, Korea, the United States, Canada, and the Soviet Union, has been extensively used as a cure-all drug or tonic in the Orient (Choi, 1988). American ginseng, grown in North America, is *Panax quinquefolium* L. (Carlson, 1986; Proctor and Bailey, 1987). In 1987 and 1988, the United States exported 578 and 448 t, respectively, of American ginseng root (valued at nearly \$100 million) to other countries, mainly in Asia (US Bureau of Census). The majority of exported ginseng is produced in Wisconsin (Wis. Dept. of Agr.). Wisconsin harvested 592 and 472 t of the ginseng root in 1987 and 1988, respectively.

Improvement of ginseng root production through conventional breeding is slow and difficult, since the plant is a perennial and

requires 3 to 4 years to flower and 4 to 5 years to harvest. Tissue-culture technology might be able to accelerate the rate of improvement. For Korean ginseng (*Panax ginseng* C.A. Meyer), callus cultures have been successfully established (Butenko et al., 1968), plantlets have been regenerated from

root-derived embryogenic callus (Chang and Hsing, 1980a; Choi, 1988; Furuya et al., 1986), and in vitro flowering of somatic embryos from root-derived callus has been reported (Chang and Hsing, 1980b). Callus cultures of Korean ginseng have been transformed with *Agrobacterium rhizogenes* to produce many hairy roots that synthesize the same saponins and ginsenosides as did untransformed roots (Yoshikawa and Furuya, 1987).

Although callus cultures have been established using leaves, roots, and stems of American ginseng (Butenko et al., 1968; Jhang et al., 1974), plant regeneration from these cultures has not been reported. This paper reports studies on embryogenic callus induction, callus growth, and plant regeneration of American ginseng.

A 4-year-old fresh field-grown ginseng root was washed in 70% alcohol for 40 sec, then blotted and dried with paper towels. The cleaned ginseng root was then sterilized in 2.5% sodium hypochlorite for 20 min under vacuum with occasional agitation, rinsed with sterile, distilled water three times, and sliced into 2-mm sections. Pith tissue of each section was removed with a cork borer 1 cm in

Fig. 1. Net callus growth of American ginseng cultured in MS medium containing 2 mg dicamba (□) or 2 mg 2,4-D (■)/liter. Each point represents the average callus growth from at least four plates. Each plate initially contained 50 ml of the maintenance medium and 1 g of callus. The vertical bars represent these of the mean.

Received for publication 1 May 1989. I thank Ross Eccleshall and Alexi Miller for reviewing the manuscript, and Faye Wang for technical assistance. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked advertisement solely to indicate this fact.

¹Present address: Northrup King Co. Research Center, Highway 19, Stanton, MN 55081.

Fig. 2. Friable embryogenic callus(a), mature somatic embryos(b), and regenerated plantlets(c) of American ginseng.

Table 1. Effects of the plant growth regulators (PGR) BA, GA, IBA, KIN, and NAA on plating efficiency of American ginseng cultures.^a

PGR concn (mg·liter ⁻¹)	Plating efficiency (%) ^b				
	BA	GA	IBA	KIN	NAA
0.10	0	17	72	5	56
0.50	5	56	56	22	22
1.00	33	89	33	22	0
2.00	22	89	27	17	5

^a0.5 g of embryogenic callus was evenly spread on a 60 × 20-mm petri dish that contained 20 ml of MS medium supplemented with various PGRs and maintained at 27C with a 16-hr photoperiod.

^bAll percentages based on 18 plates.

diameter, transferred onto a callus induction medium, and incubated in the dark at 27C.

The callus induction medium consisted of Murashige and Skoog (MS) salts (Murashige and Skoog, 1962), thiamine HCl (10 mg·liter⁻¹), casein hydrolysate (100 mg·liter⁻¹), sucrose (30 g·liter⁻¹), Difco-Bacto agar (8 g·liter⁻¹), 2,4-D (2 mg·liter⁻¹), and KIN (1 mg·liter⁻¹). Dark-greenish, compact calli were observed on the root-pith sections 4 weeks after initiation.

Four weeks later, the calli, were transferred to a maintenance medium similar to the initiation medium except that KIN was omitted and gelrite (Kelco) (2 g·liter⁻¹) was used instead of agar. Friable yellowish callus formed on the edge of the compact callus 2 to 3 weeks after the transfer. This friable callus grew very slowly. To enhance callus growth, dicamba (2 mg·liter⁻¹) was substituted for 2,4-D in the maintenance medium. Not only did the callus grow faster on the

medium containing dicamba than on the 2,4-D medium (Fig. 1), but the callus became embryogenic (Fig. 2a). The callus has been maintained in the dark for more than 3 years and subculture at 3-week intervals.

The friable embryogenic callus grows well in a liquid maintenance medium supplemented with dicamba. American ginseng seems to develop more readily on a medium supplemented with dicamba, in contrast to Korean ginseng, which grows better on a medium containing 2,4-D (unpublished data).

For plant regeneration, embryogenic callus was transferred onto an embryo-maturation medium that was composed of MS salts, B5 vitamins (Gamborg et al., 1968), NAA (0.4 mg·liter⁻¹), 2,4-D (1 mg·liter⁻¹), sucrose (30 g·liter⁻¹), and agar (7 g·liter⁻¹) and maintained at 27C with a 16-hr photoperiod under cool-white fluorescent light (60 μmol·s⁻¹·m⁻²). The somatic embryos matured in about 3 weeks (Fig. 2b). Mature somatic embryos were then transferred to MS medium supplemented with various concentrations of BA, GA, IBA, KIN, or NAA and also grown at 27C with a photoperiod of 16 hr.

The effect of these plant growth regulators on germination of somatic embryos was studied (Table 1). Overall, GA gave the highest plating efficiency (89%) and BA and KIN the lowest (≈20%). Also, plantlets derived from callus grown on medium containing BA, GA, or KIN were vitreous and abnormal, and roots were not formed. These plantlets were difficult to maintain.

Plantlets obtained from callus grown on medium containing IBA or NAA were nor-

real, although their growth was relatively slow. Therefore, the regeneration medium was modified by using a combination of IBA and NAA in ratios of 0.1:0.1, 0.1:0.5, 0.5:0.1, and 0.5:0.5 mg·liter⁻¹. Plantlets regenerated on medium supplemented with 0.5 mg IBA and 0.1 mg NAA/liter were the healthiest (Fig. 2c), with a regeneration frequency of 30%.

This study reports the initiation of friable embryogenic callus from American ginseng root and plant regeneration from its somatic embryos. The method should be useful for genetic improvement of ginseng.

Literature Cited

- Butenko, R. G., LV. Brushwitzky, and L.I. Slepian. 1968. Organogenesis and somatic embryogenesis in the tissue culture of *Panax ginseng* C.A. Meyer. *Bet. Zh.* 7:906-913.
- Carlson, A.W. 1986. Ginseng: America's botanical drug connection to the Orient. *Econ. Bot.* 40:233-249.
- Chang, W.C. and Y.I. Hsing. 1980a. Plant regeneration through somatic embryogenesis in root-derived callus of ginseng (*Panax ginseng* C.A. Meyer). *Theor. Applied Genet.* 57:133-135.
- Chang, W.C. and Y.I. Hsing. 1980b. *In vitro* flowering of embryoids derived from mature root callus of ginseng (*Panax ginseng*). *Nature* 284:341-342.
- Choi, K.T. 1988. *Panax ginseng* C.A. Meyer: Micropropagation and the *in vitro* production of saponins, p. 484-500. In: Y.P.S. Bajaj (ed.). *Biotechnology in agriculture and forestry*, vol. 4. Medicinal and aromatic plants I. Springer-Verlag, Berlin.
- Furuya, T., T. Yoshikawa, K. Ushiyama, and H. Oda. 1986. Formation of plantlets from callus culture of ginseng (*Panax ginseng*). *Experientia* 42:193-194.
- Gamborg, O. L., R.A. Miller and K. Ojima. 1968. Nutrient requirements of suspension culture of soybean root callus. *Expt. Cell Res.* 50:151-158.
- Jhang, J. J., E.J. Staba, and J.Y. Kim. 1974. American and Korean ginseng tissue cultures: growth, chemical analysis, and plantlet production. *In Vitro* 9:253-259.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15:473-479.
- Proctor, J.T.A. and W.G. Bailey. 1987. Ginseng: industry, botany, and culture. *Hort. Rev.* 9: 187-236.
- Yoshikawa, T. and T. Furuya. 1987. Saponin production by cultures of *Panax ginseng* transformed with *Agrobacterium rhizogenes*. *Plant Cell Rpt.* 6:449-453.