Curd Formation and Flowering of Plantlets Regenerated from Cauliflower Curd Explants

Ghazanfar Ali Shah¹ and Guo Deping²

Department of Horticulture, Zhejiang Agricultural University, Hangzhou, Zhejiang 310029, China

Zeng Guangwen³

Section of Plant Physiology, Zhejiang Agricultural University, Hangzhou, Zhejiang 310029, China

Additional index words. micropropagation, in vitro, curd, flowering, *Brassica oleracea* var. *botrytis*

In vitro flowering has been reported for several plant species (Scorza, 1982). However, to our knowledge, there have been no reports on in vitro flowering in *Brassica* spp., with the exception of a nonheading Chinese cabbage [*Brassica campestris* ssp. *chinensis* (L.) Makino] commonly known as To-Pe-tsai, from which flowers were obtained from seedling shoot-tip cultures (Shinora et al., 1981). We have observed curd formation and flowering on plantlets regenerated from cauliflower (*B. oleracea* var. *botrytis* L. '60 Day') curd explants. The objectives of this work were to confirm the occurrence of in vitro flowering in cauliflower and to investigate the effect of temperature on flowering.

Curd portions collected from plants grown in the vegetable garden of Zhejiang Agricultural Univ. were cut into small pieces and soaked in tap water plus Tween-20 for 30 min. Curd pieces were soaked in 70% ethanol for 15 to 30 sec, rinsed two times in sterile distilled water, soaked in 0.170 mercuric chloride for 3 to 5 min, and rinsed three more times in sterile distilled water.

Curd tissues were dissected aseptically into 2- to 3-mm-diameter pieces and planted on the surface of agar medium. The medium consisted of MS (Murashige and Skoog, 1962) salts and vitamins, 3% sucrose, and (mg-liter⁻¹) 0.1 N⁶-benzyladenine and 1.0 α -naphthaleneacetic acid (Li and Qiu, 1981). The pH was adjusted to 5.8 with 0.1N HCl or NaOH before adding agar (0.7%). The medium was dispensed into 50-ml Erlenmeyer flasks (20 ml/

flask) and autoclave at 121C for 15 min. Three explants were placed in each flask. Cultures were maintained at 25C and a 12-h photoperiod (46 μ mol-m⁻²-s⁻¹) supplied by cool-white fluorescent tubes.

The explants began to form calli from the cut edges after $\approx$ 1 week and redifferentiated shoots and roots on the same medium. After 40 days, regenerated shoots could be harvested from the cultures and planted on agar-based MS medium without growth regulators for rooting. The effect of temperature on curd formation was determined by placing the cultures in incubators at 25 or 10C and a 12-h photoperiod (34 μ mol-m⁻²-s⁻¹). This experiment was repeated three times with 20 replicates per treatment.

Plantlets treated at 10 and 25C were transplanted to pots containing 1 perlite : 1 vermiculite (v/v) and kept at 25C to determine their vernalization response. The plantlets grown in vitro at 10C developed complete inflorescences (Table 1), flowered, and formed pods, but most plantlets kept at 25C developed only curds and a few developed rudimentary inflorescences.

Plantlets at 10C had visible curds after 30 days (Fig. 1); at 25C there were no curds, even after 2 months, when the plantlets wilted and died. Plantlets at 10C developed inflorescences in vitro and on transfer to pots.

We have shown that temperature is an important factor in cauliflower curd formation and flowering in vitro and that in vitro-regenerated shoots are in the receptive stage for cold

Fig. 1. Curd formation on in vitro *Brassica oleracea* var. *botrytis* plantlet grown at 10C and then transferred to 25C.

induction. Further studies concerning the mechanism of in vitro curd and inflorescence formation are needed. Since we used a growth-regulator-free medium, our system can be used for studying hormonal regulation of curd formation and flowering in cauliflower.

Literature Cited

- Li, S.H. and W.D. Qiu. 1981. The production of plants from tissue cultures of cauliflower (*Brassica oleracea* var. *botrytis* L.) (in Chinese). *Acta Hort. Sinica* 8(1):33-36.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15:473-497.
- Scorza, R. 1982. In vitro flowering. *Hort. Rev.* 4:106-127.
- Shinohara, Y., S. Yoshio, and Y.L. Yunn. 1981. Vernalization of non-heading Chinese cabbage in vitro. *Proc. First Intl. Symp. Chinese Cabbage*. Asian Veg. Res. and Dev. Ctr. Publ. 81-138. p. 245-252.

Received for publication 29 June 1992. Accepted for publication 14 Dec. 1992. Research conducted in the Horticultural Dept. of Zhejiang Agricultural Univ., China: This publication is part of the PhD thesis of G.A.S. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Graduate Student.

²Lecturer of horticulture.

³Professor of plant physiology.

Table 1. Effect of temperature on growth and development of plantlets from cauliflower curd explants.

Temp (°C)	Plantlets ¹			Transplants ²		
	Curd formation (%)	Leaves (no.)	Plantlet ht (cm)	Leaves (no.)	Plant ht (cm)	Inflorescence stalk length (cm)
10+ 25 ³	100	7.0 $\pm$ 0.2	4.0 $\pm$ 0.4	10.2 $\pm$ 1.9	17.2 $\pm$ 1.2	11.0 $\pm$ 1.0
25 + 25 ⁴	0	11.4 $\pm$ 0.6	4.1 $\pm$ 0.4	19.3 $\pm$ 1.9	10.8 $\pm$ 0.9	4.1 $\pm$ 1.0

¹Data collected after 30 days in vitro.

²Data collected 90 days after transplanting to pots.

³10C in vitro; 25C after transplanting.

⁴25C in vitro; 25C after transplanting.