Elevated Carbon Dioxide Storage of 'Anjou' Pears Using Purge-controlled Atmosphere

Stephen R. Drake¹

U.S. Department of Agriculture, Agricultural Research Service, Tree Fruit Research Laboratory, 1104 North Western Avenue, Wenatchee, WA 98801

Additional index words. *Pyrus communis*, controlled-atmosphere storage

Abstract. 'Anjou' pears (*Pyrus communis* L.) were placed in controlled-atmosphere (CA) storage immediately after harvest (<24 hours) or after a 10-day delay in refrigerated storage, and held there for 9 months at 1°C. Oxygen in all atmospheres was 1.5% and CO₂ was at either 1% or 3%. Atmospheres in the flow-through system were computer-controlled at ±0.1%. After removal from CA storage, pears were evaluated immediately and after ripening at 21°C for 8 days. Pears stored in 3% CO₂ were firmer, greener, and displayed less scald, internal breakdown, and stem-end decay than pears stored in 1% CO₂. In addition, no internal discoloration of 'Anjou' pears was evident when held with 3% CO₂. 'Anjou' pears held in 3% CO₂ retained the ability to ripen after long-term storage. A 10-day delay in atmosphere establishment had little or no influence on the long-term keeping quality or ripening ability of 'Anjou' pears.

Atmospheres of 1% to 2% O₂ and < 1.0% CO₂ are recommended for controlled-atmosphere (CA) storage of 'Anjou' pears (Hansen and Mellenthin, 1979; Hardenburg et al., 1986). Using 2% or less O₂ for long-term pear storage reduced loss of firmness, acidity, and greenness, and scald severity (Chen et al., 1981; Mellenthin et al., 1980; Richardson and Meheriuk, 1989). Elevated CO₂ in long-term pear storage can cause brown core (Allen and Claypool, 1948; Hansen, 1957). Although higher CO₂ enhances storage life of pears (Claypool, 1973; Hansen and Mellenthin, 1962), the results are conflicting because they involve cultivar, maturity, tree vigor, and storage delay. Exposing 'Anjou' pears to prestorage high-CO₂ treatments for short periods of time prolongs storage life (Wang and Mellenthin, 1975). Removing CO₂ to a level of ≤1% is a costly and time-consuming operation (Blanpied, 1988; Wealti and Cavalieri, 1990).

Previous research in CA pear storage has been conducted with static systems where fruit is held in the same atmosphere over the entire storage period and only O₂ and CO₂ levels are regulated. Many new CA facilities employ a flow-through system where fruit is exposed to a flowing atmosphere of desired concentrations. This research was conducted to determine the quality of 'Anjou' pears stored under two CO₂ levels where the atmospheres

were established at different times after harvest in a flow-through-type facility.

Materials and Methods

This study was conducted over 3 years using 'Anjou' pears grown in the Wenatchee, Wash., district. During the first year, eight boxes (four individual lots) were obtained from a commercial warehouse 1 day after harvest. Pears from three orchards with a known history of differences in storage quality were used in two of the years. Pears were divided into four groups. CA of 1.5% O₂ and 1% CO₂ or 1.5% O₂ and 3% CO₂ at 1°C were established within 24 h after harvest on two groups. Identical CA regimes were established on the remaining two groups after 10 days in cold storage (1°C). Pears were stored for 9 months before evaluation. All atmospheres were established and maintained throughout the storage period at ±0.1% using a computer control system (Technical Consulting Services,

Chelan, Wash.). Nitrogen for this purge-type CA system was supplied by a Generon Membrane Separation System (Generon, Houston). Bottled CO₂ was used as the CO₂ source.

Storage evaluation consisted of 20 pears for all combinations of atmosphere, time delay, and replication. Ten pears were evaluated immediately after removal from storage, and the other 10 were allowed to ripen for 8 days at ambient temperature (20°C) before evaluation. Quality factors evaluated were external and internal color, firmness, soluble solids concentration (SSC), titratable acidity (TA), carbohydrates, and visible disorders.

External and internal color were determined with a CR300 Minolta chromameter (Osaka, Japan) using the Hunter L, a, b mode and calibrated with a white plate (no. 11133144). Three values for external color were determined around the circumference of each fruit, and the average value for 10 fruit was reported. Internal color was measured by cutting each fruit in half equatorially and immediately reading the exposed flesh surface of the calyx end; the average value for 10 fruit was reported. A model EP1 pressure tester (Lake City, Kelowna, B.C., Canada) equipped with a 7.8-mm head was used to determine firmness. Juice prepared from pear slices was titrated to pH 8.2 with 0.1 N NaOH and values were expressed as percentage of malic acid. SSC was determined with an Abbe-type refractometer calibrated at 20°C. Carbohydrates were determined by the high-performance liquid chromatography method described by Bio-Rad (Bio-Rad, Richmond, Calif.). Disorders (CO₂ injury, scald, internal breakdown, cork spot, and stem decay) were evaluated by visual assessment and expressed as the percentage of fruit affected. Analysis of variance was determined by MSTAT (1988) as a factorial design. Based on significant F test, means were separated by Duncan's multiple range test.

Results and Discussion

'Anjou' pears stored in 3% CO₂ were 41% firmer (15.6 N) than those stored in 1% CO₂ after 9 months of storage (Table 1). When

Table 1. Firmness, soluble solids concentration (SSC), titratable acidity (TA), and SSC : TA ratio for 'Anjou' pears as influenced by controlled-atmosphere storage conditions.

Treatment	Firmness (N)	SSC (°Brix)	TA (% malic)	SSC : TA ratio
Atmosphere × ripening (days) ^a				
1.5% O ₂ + 1% CO ₂				
0	37.8 b ^b	13.0 ab	0.24 a	55 b
8	19.1 c	12.5 c	0.21 c	61 a
1.5% O ₂ + 3% CO ₂				
0	53.4 a	13.1 a	0.24 a	54 b
8	19.6 c	12.7 b	0.22 b	59 c
Atmosphere × delay (days) ^a				
1.5% O ₂ + 1% CO ₂				
0	26.2 b	12.8 b	0.22 b	58 a
10	30.2 b	12.8 b	0.22 b	58 a
1.5% O ₂ + 3% CO ₂				
0	36.0 a	12.8 b	0.24 a	54 b
10	37.4 a	13.1 a	0.22 b	59 a

^aPears were held at 21°C for 8 days for ripening.

^bMean separation within columns and treatment groups by Duncan's multiple range test ($P \leq 0.05$).

^cAtmosphere establishment delayed 10 days after harvest; pears were held at 1°C during this period.

Received for publication 25 June 1993. Accepted for publication 2 Nov. 1993. I thank the Washington State Tree Fruit Research Commission for its financial support of this project. Use of a company or product name by the department does not imply approval or recommendation of the product to the exclusion of others that may also be suitable. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked advertisement solely to indicate this fact.
¹Research Horticulturist.

allowed to ripen for 8 days at ambient temperature, pears from either 1% or 3% CO₂ storage ripened to a similar firmness level (19 N). When CA was established immediately after harvest (<24 h), pears stored in 3% CO₂ were 37% firmer (9.8 N) than pears stored in 1% CO₂ (Table 1). When pears were in cold storage (1C) for 10 days before atmosphere establishment, the difference in firmness was 24% (7.2 N) in favor of pears that were stored in 3% rather than in 1% CO₂.

There was no difference in the SSC of 'Anjou' pears due to CO₂ concentration. There was a loss in SSC when pears were ripened for 8 days. Ripened pears from 3% CO₂ lost less SSC during ripening (3%) than pears from 1% CO₂ (4%) (Table 1). There was some difference (2%) in SSC of pears when CA establishment was delayed 10 days but only when stored in the 3% CO₂ atmosphere (Table 1). Differences for individual carbohydrate concentrations (sucrose, fructose, glucose, and sorbitol) between pears from the two CO₂ atmospheres were minimal (data not shown). Neither CO₂ content of the atmosphere nor a 10-day delay in atmosphere establishment resulted in a change in the individual carbohydrate concentrations in 'Anjou' pears.

TA values were similar for pears stored in 1% or 3% CO₂. When ripened for 8 days, 'Anjou' pears from the 3% CO₂ storage did not lose as much acid as pears from the 1% CO₂ storage environment (Table 1). A 10-day delay in CA establishment resulted in pears with similar acid values regardless of CO₂ level. Pears from the 3% CO₂ storage were higher initially in TA, or when CA was established in ≤24 h, but when CA establishment was delayed, TA content was identical (Table 1).

Pears from both CO₂ storage environments had similar SSC : TA ratios after ripening (Table 1). Those in 1% CO₂ had a higher SSC : TA ratio when atmosphere was established immediately, but the SSC : TA ratio was the same for the two CO₂ levels when atmosphere establishment was delayed 10 days (Table 1).

'Anjou' pears stored in 3% CO₂ were greener (higher hue value) than pears stored in 1% CO₂ at removal and after ripening (Table 2). Pears lost green pigmentation during ripening regardless of the CO₂ level in storage, but this loss in color was slightly greater with pears stored in 1% CO₂ than those stored in 3% CO₂. This difference in color was also evident when Hunter "L" values were considered. Fruit from 3% CO₂ had a lower Hunter "L" value (indicative of a darker surface color than pears stored in 1% CO₂), but no difference was present after 8 days of ripening.

The difference in Hunter color values between pears from 1% or 3% CO₂ at removal and after ripening was visually quite apparent. Hue values (arctan b/a values) also indicated a much greener (less yellow) pear surface when pears were stored in 3% than in 1% CO₂. Hue values, after ripening, also indicated that pears stored in 3% CO₂ changed color from green to yellow more slowly than those stored in 1% CO₂. A 10-day delay in atmosphere establishment had no influence on Hunter color "a" and "L" values or hue.

Table 2. External and internal color of 'Anjou' pears as influenced by controlled-atmosphere storage conditions.

Treatments	Hunter color					
	External			Internal		
	L	a	Hue	L	Hue	
Atmosphere × ripening (days) ^z						
1.5% O ₂ + 1% CO ₂						
0	63.7 a ^y	−3.6 c	97.6 b	69.3 a	18.7 a	84.7 b
8	65.6 a	0.4 a	88.8 d	69.4 a	18.3 a	84.9 ab
1.5% O ₂ + 3% CO ₂						
0	57.6 b	−5.9 d	102.8 a	72.1 a	18.4 a	86.3 a
8	66.3 a	−2.1 b	94.2 c	70.5 a	18.0 a	84.3 ab
Atmosphere × delay (days) ^x						
1.5% O ₂ + 1% CO ₂						
0	62.8 a	−1.0 a	91.9 b	67.8 a	18.4 a	84.5 a
8	66.5 a	2.1 a	94.4 b	70.8 a	18.6 a	85.2 a
1.5% O ₂ + 3% CO ₂						
0	58.5 a	−4.1 b	98.6 a	69.8 a	17.9 a	85.0 a
8	65.0 a	−3.9 b	98.4 a	72.8 a	18.5 a	85.6 a

^zPears were held at 21C for 8 days for ripening.

^yMean separation within columns within groups by Duncan's multiple range test ($P \leq 0.05$).

^xAtmosphere establishment delayed 10 days after harvest; pears were held at 1C during this period.

Hansen and Mellenthin (1962) and Claypool (1973) reported that high CO₂ in the storage atmosphere will result in a darker internal color of 'Anjou' pears. Internal color was not affected for pears stored in 3% CO₂ regardless of ripening time or delay in atmosphere establishment (Table 2). Internal Hunter "L" values were identical for pears from both CO₂ atmospheres before and after ripening, regardless of delay in CA establishment. Hunter "b" values or yellowness were also similar for all treatments. Internal hue values were not influenced by either ripening time or a delay for atmosphere establishment, but there was a change in pears from the different CO₂ atmospheres before ripening. This difference in hue values was slight (1.6 units) and was considered of no commercial consequence even though a change of 1.0 unit is visually detectable (Hunter and Harold, 1987).

Scald, internal breakdown, and stem decay (Table 3) were significantly reduced in 'Anjou' pears that were stored in 3% CO₂ when compared to 1% CO₂. Internal breakdown (core browning) was of particular concern, but in this study pears had a lower incidence of the disorder in 3% CO₂ than in 1% CO₂. Pears from one of the orchards used in this study have a history of internal breakdown and are poor candidates for CA storage, but reduced internal breakdown was also apparent when

pears from this orchard were stored in 3% CO₂. Dilley (1993), working with apples (*Malus domestica* Borkh.), suggested that purge CA may flush volatile compounds that contribute to disorders. Gast (1993) noted a reduction in volatile production when apples were stored in a purge CA system. A higher incidence of cork spot was found in fruit stored in 3% CO₂ (11%) than in 1% CO₂ (5%). No external surface injury from 3% CO₂ was evident after long-term storage.

After 8 days of ripening, scald and internal breakdown were more apparent, but cork spot was less evident than immediately after storage (Table 3). The increase (34%) in scald was of major concern, particularly if no use of scald control chemicals is required by law. The increase in internal breakdown, although significant, was much less (19%). After ripening, stem decay was less apparent, presumably because the stem dried. A 10-day delay in atmosphere establishment resulted in reduced cork spot but increased internal breakdown with no change in amount of scald or stem decay present. A delay in atmosphere establishment might allow the fruit to become more acclimated to one environment before subjecting fruit to a second environment.

Maintaining a higher CO₂ in the storage atmosphere reduces operational cost (Wealti and Cavalieri, 1990). Three percent CO₂ in the

Table 3. Incidence of disorders and stem decay of 'Anjou' pears as influenced by controlled-atmosphere storage conditions.

Treatments	Scald (%)	Internal breakdown (%)	Cork spot (%)	Stem decay (%)
Atmosphere				
1.5% O ₂ + 1% CO ₂	22.3 a ^z	40.3 a	5.1 b	14.4 a
1.5% O ₂ + 3% CO ₂	12.3 b	21.0 b	10.9 a	2.1 b
Ripening (days) ^y				
0	0.0 b	28.1 b	9.8 a	9.3 a
8	34.6 a	33.4 a	6.3 b	7.3 b
Delay (days) ^x				
0	16.9 a	28.9 b	12.2 a	8.1 a
10	17.6 a	32.5 a	3.8 b	8.4 a

^zMean separation within pairs by analysis of variance ($P \leq 0.05$).

^yPears were held at 21C for 8 days for ripening.

^xAtmosphere establishment delayed 10 days after harvest; pears were held at 1C during this period.

storage atmosphere is well above the recommended CO₂ atmosphere of 1% to 1.5% for 'Anjou' pears. In this study there were definite quality advantages for 'Anjou' pears stored at 3% CO₂ for 9 months relative to 1% CO₂ storage. These quality benefits include reductions in loss of firmness and greenness and disorders. A 10-day delay between harvest and atmosphere establishment had little or no influence on the long-term keeping quality or ripening ability of 'Anjou' pears.

Literature Cited

- Allen, F.W. 1939. Influence of carbon dioxide in lengthening the life of Bartlett pears. *Proc. Amer. Soc. Hort. Sci.* 37:473-478.
- Allen, F.W. and L.L. Claypool. 1948. Modified atmosphere in relation to the storage life of Bartlett pears. *Proc. Amer. Soc. Hort. Sci.* 59:192-204.
- Blanpied, G.D. 1988. Items of interest for apple operators in New York, New England and beyond. *Cornell Fruit Handling and Storage Nwsl.*, August.
- Chen, P.M., R.A. Spotts, and W.M. Mellenthin. 1981. Stem-end decay and quality of low oxygen stored 'd'Anjou' pears. *J. Amer. Soc. Hort. Sci.* 106:522-527.
- Claypool, L.L. 1973. Further studies on controlled atmosphere storage of 'Bartlett' pears. *J. Amer. Soc. Hort. Sci.* 98:289-293.
- Dilley, D.R. 1993. Effect on apple quality of purge vs. static systems for atmosphere control for CA storage. *Proc. Wash. Tree Fruit Postharvest Conf.* 9:83-85.
- Gast, W.A., Jr. 1993. Purge vs. static: Is there a quality difference? *Proc. Wash. Tree Fruit Postharvest Conf.* 9:82.
- Hansen, E. 1957. Reactions of Anjou pears to carbon dioxide and oxygen content of the storage atmosphere. *Proc. Amer. Soc. Hort. Sci.* 69:110-115.
- Hansen, E. and W.M. Mellenthin. 1962. Factors influencing susceptibility of pears to carbon dioxide injury. *Proc. Amer. Soc. Hort. Sci.* 80:146-153.
- Hansen, E. and W.M. Mellenthin. 1979. Commercial handling and storage practices for winter pears. *Oregon State Univ. Agr. Expt. Sta. Rpt.* 550.
- Harlenburg, R.E., A.E. Watada, and C.Y. Wang. 1986. The commercial storage of fruits, vegetables and florist and nursery stocks. *U.S. Dept. of Agr., Agr. Hdbk.* 66.
- Hunter, R.S. and R.W. Harold. 1987. The measurement of appearance, 2nd ed. Wiley, New York.
- Mellenthin, W.M., P.M. Chen, and S.B. Kelly. 1980. Low oxygen effects on desert quality, scald prevention and nitrogen metabolism of 'd' Anjou' pear fruit during long term storage. *J. Amer. Soc. Hort. Sci.* 105:522-527.
- MSTAT, 1988. Version 1.0, Michigan State Univ., East Lansing.
- Richardson, D.G. and M. Meheriuk. 1989. CA recommendations for pears (including Asian pears), p. 285-302, vol 2. In: J.K. Fellman (ed.). *Proc. 5th Intl. Controlled Atmosphere Res. Conf.* June 1989, Wenatchee, Wash.
- Wang, C.Y. and W.M. Mellenthin. 1975. Effect of short-term high CO₂ treatment on storage of 'd' Anjou' pears. *J. Amer. Soc. Hort. Sci.* 100:492-495.
- Wealti, H. and R.P. Cavalieri. 1990. Matching nitrogen equipment to your needs. *Tree Fruit Postharvest J.* 1:3-13.