Modeling Flower Bud Elongation in Easter Lily (*Lilium longiflorum* Thunb.) in Response to Temperature

P.R. Fisher and J.H. Lieth

Department of Environmental Horticulture, University of California, Davis, CA 95616-8587

R.D. Heins

Department of Horticulture, Michigan State University, East Lansing, MI 48824-1325

Additional index words. decision support, bud length, bud development meter, simulation

Abstract. A model was developed to quantify the response of Easter lily ('Nellie White') flower bud elongation to average air temperature. Plants were grown in greenhouses set at 15, 18, 21, 24, or 27C after they had reached the visible bud stage. An exponential model fit the data with an R^2 of 0.996. The number of days until open flowering could be predicted using the model because buds consistently opened when they were 16 cm long. The model was validated against data sets of plants grown under constant and varying greenhouse temperatures at three locations, and it was more accurate and mathematically simpler than a previous bud elongation model. Bud length can be used by lily growers to predict the average temperature required to achieve a target flowering date, or the flowering date at a given average temperature. The model can be implemented in a computer decision-support system or in a tool termed a bud development meter.

The Easter lily has a precise market window that varies from year to year depending on the date of Easter. Many Easter lily growers use a flower bud elongation model, the most widely used version of which was developed by Healy and Wilkins (1984), termed the "HW model," to help control flowering date by manipulating temperature. These authors modeled bud elongation in two linear phases, with a junction at the point where bud length reached 6 cm. In a greenhouse validation trial (Fisher, 1995), we found a major bias in the predictions of the HW model, and also in predictions of an exponential function we fitted to their data. Near visible bud, the stage when flower buds first become externally visible, both models underpredicted the average temperature required to achieve a target flowering date. Required temperatures were overpredicted as the crop approached flowering.

Our objective was to develop a bud elongation model to dynamically simulate bud elongation in response to average greenhouse air temperature (T) and other variables (Table 1). We aimed to validate the model under both constant and varying greenhouse temperatures. The model was designed to predict the re-

quired T to achieve a target flowering date or, alternatively, the number of days to first open flower (D) for a given T for use in a decision-support system (Fisher, 1995). A further goal was to develop a new bud development meter (Healy and Wilkins, 1984) to summarize model results as a tool for growers.

Materials and Methods

Model theory. An exponential function was used to describe flower bud length (B in mm) at time t (days) under constant temperature conditions, given an initial bud length measurement of B_0 (mm) at time t_0 :

$$B = B_0 e^{k(t-t_0)} \quad [1]$$

Healy and Wilkins (1984) found that bud elongation rate increased as temperature in-

creased between 15 and 27C. To incorporate the temperature response, the rate parameter k (day^{-1}) in Eq. [1] can be replaced by some function of temperature, $f(T)$, so that Eq. [1] becomes

$$B = B_0 e^{f(T)(t-t_0)} \quad [2]$$

Easter lily flowers consistently open at some maximum bud length (B_f , in mm) of ≈ 160 mm (Healy and Wilkins, 1984). The number of days (D) to the date when the first flower opens (t_f , in days) for any value of B can therefore be calculated as

$$B_f = B e^{f(T)D} \quad [3]$$

and D can be estimated by

$$D = \ln(B_f/B)/f(T) \quad [4]$$

Similarly, T can be calculated by algebraic manipulation of Eq. [4] if D is given and the inverse of $f(T)$ is known.

Experiment design: Calibration data set. Easter lily plants ('Nellie White' in all experiments) with clearly visible flower buds were selected from a commercial greenhouse on 15 Mar. 1994. Five plants per treatment were placed in Michigan State Univ. (MSU) greenhouses set at 15, 18, 21, 24, or 27C [actual daily temperatures were 15.5 ± 0.2 , 18.3 ± 0.1 , 21.2 ± 0.1 , 24.3 ± 0.1 , and 27.0 ± 0.1 C (mean $\pm$ standard error), respectively]. Flower bud length of a single randomly selected bud was measured to the nearest millimeter on each plant three times per week. In all experiments, plants were inspected daily and the date on which the first flower opened on each plant ("open flower" stage) was recorded.

Fixed temperature settings: Validation data set 1. Plants were obtained from a commercial greenhouse on 8 Mar. 1995 immediately after the visible bud stage (which occurs when immature flower buds are externally visible without leaves being moved), and five plants per treatment were placed in MSU greenhouses set at 18, 21, 24, and 27C (actual daily temperatures 19.0 ± 0.2 , 20.9 ± 0.2 , 21.7 ± 0.7 , and 27.7 ± 1.3 C, respectively). An additional five plants were placed first at 18C until they were 60 mm long, then at 24C. A further group

Table 1. List of abbreviations and parameters.

Symbol	Description	Units
B	Flower bud length	mm
B_0	Initial bud length	mm
B_f	Final bud length	mm
c_1	Parameter in $f(T)$	day^{-1}
c_3	Parameter in $f(T)$	$\text{C}^{-2} \text{day}^{-1}$
D	Days to first open flower	days
k	Rate parameter in exponential bud length model	day^{-1}
$f(T)$	Function quantifying k as a function of T	day^{-1}
KVL	Royal Veterinary and Agricultural Univ., Copenhagen, Denmark	
MSU	Michigan State Univ.	
t	Time	days
t_f	Time of first open flower	days
t_0	Time zero	days
T	Average air temperature	$^{\circ}\text{C}$
UCD	Univ. of California at Davis	

Received for publication 25 Sept. 1995. Accepted for publication 31 Jan. 1996. We thank Univ. Outreach at Michigan State Univ. and the California Association of Nurserymen for providing funding for this project. Thanks also to Niels Ehler, Poul Karlsen, and Michael Brogaard for their experimentation at the Royal Veterinary and Agriculture College in Copenhagen, Denmark. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

Fig. 1. Estimates of k , the rate parameter in Eq. [1], obtained using nonlinear regression to fit Eq. [1] separately to data from five temperature treatments in the calibration experiment (+ symbol, $\pm$ asymptotic standard error) and the quadratic $f(T)$ function (solid line).

Fig. 3. Observed flowering dates (+, mean $\pm$ standard error for five plants per treatment) at various temperatures in the calibration experiment, and the flowering dates predicted by the ($\diamond$) exponential model and the ($\bullet$) Healy-Wilkins model.

of five plants were moved from 24 to 18C when the flower buds were 60 mm long.

Variable temperature settings: Validation data set 2. Fifteen plants were grown in each of three greenhouses under varying temperatures between 10 and 27C during Mar. 1994 at MSU, the Univ. of California at Davis (UCD), and the Royal Veterinary and Agriculture Univ. at Copenhagen (KVL). Temperatures and the design were described in detail by Fisher (1995). Temperatures were set twice per week based on a development model (Fisher, 1995) that aimed to have plants in flower by target dates of 1 Apr. at UCD and 7 Apr. at MSU and KVL. Temperatures in these experiments were therefore variable (averaging 18.4 ± 0.5 C, 19.9 ± 0.8 C, and 15.3 ± 0.4 C each day for the UCD, MSU, and KVL experiments, respec-

tively) and similar to those in a commercial greenhouse. Differences in average temperatures between locations existed because of differences in the timing of initial plant development stages and target flowering dates.

Variable temperature settings: Validation data set 3. Bud lengths were recorded on five plants grown at MSU and 15 plants grown at UCD during 1995, using the same design as validation data set 2, with a target flowering date of 1 Apr. Daily temperatures in these experiments averaged 20.0 ± 0.6 C each day at MSU and 21.2 ± 0.4 C at UCD.

Analysis. To calibrate the model, k in Eq. [1] was estimated independently for each temperature treatment by fitting Eq. [1] to the calibration data set with the nonlinear regression procedure (PROC NLIN) in SAS (SAS

Fig. 2. Predicted (solid line) and observed ($\circ$, mean $\pm$ standard error) bud length for each treatment in the calibration experiment.

Institute, 1988). For each plant, B_0 and t_0 were assigned the values of the first recorded bud length and the first measurement date, respectively. A function $f(T)$ was formulated based on the pattern of k values across the range of temperatures and was fit to a data set of k values estimated for each individual plant. $f(T)$ was incorporated into Eq. [2], and it was fit simultaneously to the entire calibration data set using nonlinear regression to estimate the parameters in $f(T)$. The final bud length at flower (B_f) was then estimated by fitting Eq. [3] with nonlinear regression to the observed D and B in the calibration data set and setting the previously estimated $f(T)$ parameters to constants.

To validate the model, the model error between predicted and observed D was exam-

This procedure resulted in $f(T) = c_1 + c_3 T^2$ so that Eq. [2] became:

$$B = B_0 e^{(c_1 + c_3 T^2)(t - t_0)} \quad [5]$$

where c_1 (day^{-1}) and c_3 ($\text{C}^{-2} \text{ day}^{-1}$) are constants. When Eq. [5] was fit to the entire data set simultaneously, c_1 and c_3 were estimated at $0.0313 \pm 0.0006 \text{ day}^{-1}$ and $8.38 \times 10^{-5} \pm 1.6 \times 10^{-6} \text{ C}^{-2} \text{ day}^{-1}$ (estimate $\pm$ asymptotic standard error), respectively. The resulting R^2 was 0.996, and the exponential model closely fit the observed bud length data (Fig. 2).

The maximum bud length at flower (B_f) from fitting Eq. [5] with nonlinear regression was estimated to be $159.8 \pm 1.0 \text{ mm}$ (asymptotic standard error), which agreed with the HW model estimate of final bud length, and the R^2 was 0.988. The predicted flowering date (t_f), which occurred when predicted B equaled B_f , was within one standard error of observed t_f in all treatments except at 24C (Fig. 3). The maximum deviation between observed and predicted t_f was 1 day for the 24C treatment. In comparison, the t_f predicted by the HW model deviated from the observed t_f by nearly 10 days at 15C.

Model validation. The deviation between predicted and observed flowering date was

generally within ± 3 days for all three validation data sets (Figs. 4 and 5) using Eq. [4]. The only exception was an initial error of nearly 4 days at 21C in validation data set 1 (Fig. 4b). In comparison, the HW model showed a consistent tendency (Figs. 4 and 5) to underestimate D until 10 days before flowering, with a deviation of up to 9 days.

Discussion

The model used here for relating Easter lily bud length to time and temperature (Eq. [5]) was more accurate than the HW model, particularly more than 10 days before flowering. The exponential model also was simpler, requiring only three parameters (c_1 , c_3 , and B_f) compared with seven in the HW model. Prediction by the exponential model of the number of days to flowering, D , was generally within ± 3 days, even as early as 4 weeks before flowering. This model performed well when validated against several greenhouse experiments.

The comparison between the exponential and HW model highlights potential problems that may arise from differences in plant temperature in greenhouse vs. growth chamber conditions. Data used by Healy and Wilkins (1984) were obtained from plants grown in

growth chambers, where plant temperature is likely to differ from air temperature due to thermal heating from the lamps.

The exponential model provides horticulturally useful information. The model could be used to predict B , T , or D by simple algebraic manipulation of Eq. [4]. A bud development meter was constructed (Fig. 6) for grower use based directly on the Healy and Wilkins (1984) tool, but incorporating the exponential model. The flower bud is placed onto the meter, and, based on observed bud length, D can be predicted given the T at the right-hand side of the meter. Mathematical equations were chosen that allow the model to be used for dynamic simulation, where changing temperature conditions can be incorporated into the model. This feature makes the exponential model suitable for incorporating into a computer decision-support system (Fisher, 1995).

Literature Cited

- Fisher, P.R. 1995. Prediction and control of stem elongation and flowering in poinsettia and Easter lily. PhD Diss., Michigan State Univ., East Lansing. (Diss. Abstr. 96-05862.)
- Healy, W.E. and H.F. Wilkins. 1984. Temperature effects on 'Nellie White' flower bud development. HortScience 19:843-844.
- SAS Institute. 1988. SAS/STAT users guide, release 6.03 ed. SAS Inst., Cary, N.C.