

Enhancement of Seed Germination in Common Carpetgrass and Centipedegrass Seed

Edward W. Bush¹, Paul Wilson², Dennis P. Shepard³, and Gloria McClure⁴

Department of Horticulture, Louisiana State University Agricultural Center, 137 J.C. Miller Hall, Baton Rouge, LA 70803

Additional index words. *Axonopus affinis*, *Eremochloa ophiuroides*, seed germination, priming, potassium nitrate

Abstract. Priming or presoaking seed of common carpetgrass (*Axonopus affinis* Chase) and centipedegrass [*Eremochloa ophiuroides* Munro. (Kunz)] increased germination percentage and decreased mean time of germination (MTG) at 20, 25, and 30 °C. The effect of presoaking and priming was dependent on grass species and temperature. The optimum seed germination temperature for both of these warm-season species was 30 °C. Maximum effect on common carpetgrass or centipedegrass seeds was achieved by priming in 2% KNO₃; higher concentrations did not improve germination percentage or MTG, and 4% was in some cases detrimental. Germination was higher and MTG lower at 20 and 30 °C than at 15 °C. Presoaking common carpetgrass and centipedegrass seeds was the most efficient seed enhancement treatment for germination at 30 °C.

Viable seed and a suitable environment are necessary for rapid establishment of turfgrass sod. Centipedegrass and common carpetgrass, both warm-season turfgrasses, are frequently established by sowing monostands or using seed mixtures. Common carpetgrass is frequently used as a companion grass with centipedegrass to reduce establishment time. Establishing warm-season grasses from seed during the early spring months when soil temperatures are cold can be challenging. Seed priming prior to planting has been used extensively with other grasses to accelerate germination time, break dormancy, and improve germination percentage and uniformity at suboptimum temperatures. Seed priming is a controlled-hydration treatment in which seeds are exposed to an external water potential sufficiently low to prevent radical protrusion, and yet stimulate physiological and biochemical

activities. Potassium nitrate (KNO₃) has been used successfully in many horticultural crops to control seed hydration (Bradford, 1986; Heydecker and Coolbear, 1977). Germination rate can be increased at temperatures considered suboptimal for germination (Brede and Brede, 1989; Hardegree, 1994; Murray, 1990). Improved seed germination may be the result of physiological changes occurring during imbibition at low-germination temperatures (McClure, 1995).

Priming with KNO₃ significantly advanced the rate of germination of perennial ryegrass (*Lolium perenne* L.), browntop [*Agrostis capillaris* L. (Pers.)], and Kentucky bluegrass (*Poa pratensis* L.) (Lush and Birkenhead, 1987). Maguire and Steen (1971) reported that KNO₃ accelerated the rate of seed germination in Kentucky bluegrass. Additionally, priming perennial ryegrass seed with polyethylene glycol (PEG) at –1.0 MPa increased germination by 35% (Danneberger et al., 1992). Adegbuyi et al. (1981) determined that priming seeds of cool-season grass species sheep fescue (*Festuca ovina* L.), ryegrass, and rough bluegrass (*Poa trivialis* L.) with priming salts (–1.0 to –1.4 MPa) significantly increased the speed of germination at suboptimum temperatures. Hardegree (1994) demonstrated that priming Great Plains native perennial grass seeds (–1.0 to –2.5 MPa) reduced mean time of germination (MTG) by 4 to 8 d. Priming wheat (*Triticum durum* L.) and wild oat (*Avena fatua* L.) seeds at –1.0 MPa accelerated germination, thereby improving seedling competitiveness against weeds (Akalehiyot and Bewley, 1977).

Although seed priming has been used extensively to improve germination of cool-season turfgrass, priming of warm-season turfgrass has not been thoroughly investigated. Presoak-

ing buffalograss [*Buchloë dactyloides* (Nutt.) Engelm.] burrs in water significantly increased MTG by 1 week (Fry et al., 1993), and soaking centipedegrass seeds in water increased germination by 7% (Walker, 1976). Toole and Toole (1939) increased common carpetgrass seed germination rate by moistening germination paper with a 0.2% KNO₃ solution. Recommendations for germinating common carpetgrass include moistening germination filter paper with 0.2% KNO₃, and exposing seeds to at least 8 h of cool-white fluorescent light (11 to 19 μmol·m⁻²·s⁻¹), and alternating temperatures 15/25 °C (Association of Official Seed Analysts, 1994). These recommendations for common carpetgrass seed are not referenced, and no seed testing recommendations are given for centipedegrass by the Association of Official Seed Analysts (1994).

We tested the hypothesis that priming and presoaking would improve common carpetgrass and centipedegrass seed germination at suboptimum (<30 °C) temperatures.

Materials and Methods

We used two grass species (common carpetgrass and centipedegrass), four isothermic germination temperatures (15, 20, 25, and 30 °C), and six seed treatments [0% (distilled water); 1%, 2%, 3%, and 4% KNO₃ solutions; and an untreated control]. These concentrations of KNO₃ provided osmotic pressure of –0.5, –0.9, –1.3, and –1.7 Pa, respectively. The two grass species and four germination temperatures were evaluated separately. A thermogradient table (Scientific Systems Corp., Baton Rouge, La.), with four isothermic temperature lanes calibrated at 15, 20, 25, or 30 °C, was sectioned into two blocks. A block consisted of six seed treatments in petri dishes randomized within each temperature lane. Cool-white fluorescent lights (19 μmol·m⁻²·s⁻¹) provided 14 h of daily illumination. The experiment was repeated twice over time for each grass species and germination temperature to establish four blocks.

Centipedegrass (Lot # 95/05; Patten Seed, Lakeland, Ga.) and common carpetgrass (Lot # U4/410; Pennington Seed, Hammond, La.) seeds were soaked for 48 h in aerated distilled water (–0.06 MPa) or primed for 48 h with KNO₃ solutions maintained at 25 °C. The osmolality of priming solutions (vapor pressure osmometer; Wescor, Logan, Utah) was measured prior to and following the 48-h treatment period. Each solution was aerated using a standard aquarium pump. Following treatments, seeds were rinsed with copious amounts of water, transferred to paper towels, and air-dried for 1 h. Fifty seeds of each seed treatment were placed into separately labeled 6.0-cm-diameter petri dishes lined with Whatman #42 filter papers moistened with 2 mL of distilled water. A control treatment (untreated seed) was also included. Seeds were misted with distilled water daily as needed to prevent desiccation.

Germination counts (radicle protrusion) were taken daily for 21 d and germinated seed discarded. MTG and germination percentage

Received for publication 27 July 1998. Accepted for publication 16 Feb. 1999. Approved for publication by the director of the Louisiana Agricultural Experiment Station as manuscript number 98-28-0211. This research was supported by funds from State and matching funds allocated to the Louisiana State Univ. Agricultural Center. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Assistant Professor. To whom reprint requests should be addressed. E-mail address: ebush@agctr.lsu.edu

²Professor. E-mail address: pwilson@agctr.lsu.edu

³Former Assistant Professor. Present address: Novartis Crop Protection Inc., 11561 Hemlock Drive, Overland Park, KS 66210.

⁴Research Assistant. Present address: Department of Plant Pathology and Crop Physiology, Louisiana State Univ. Agricultural Center, 302 Life Sciences Bldg., Baton Rouge, LA 70803.

Table 1. Effects of soaking and/or priming seeds of common carpetgrass and centipedegrass on germination during 21 d incubation at 15, 20, 25, or 30 °C.

Treatment		Germination (%)				Mean time of germination (days)			
KNO ₃ priming solution (%)	Ψ (MPa)	15 °C	20 °C	25 °C	30 °C	15 °C	20 °C	25 °C	30 °C
<i>Carpetgrass</i>									
0 (Presoaking)	-0.06	4.1 a	64.6 ab ^z	96.0 a	97.6 a	13.6 a	8.3 b	3.2 c	2.1 c
1	-0.5	5.5 a	65.6 ab	98.5 a	98.1 a	15.0 a	7.5 b	3.3 c	2.0 c
2	-0.9	6.1 a	83.7 a	99.0 a	97.0 a	14.6 a	7.2 b	3.6 bc	2.1 c
3	-1.3	2.1 a	85.6 a	96.9 a	99.5 a	12.5 a	7.1 b	3.4 c	2.2 c
4	-1.7	4.7 a	82.7 a	96.6 a	95.9 a	15.3 a	8.0 b	4.0 b	2.9 b
Control	(untreated)	6.0 a	49.3 b	87.9 b	95.1 a	12.3 a	10.2 a	5.6 a	4.4 a
Linear		NS	*	NS	NS	NS	NS	NS	NS
Nonlinear		NS	NS	NS	NS	NS	NS	NS	NS
<i>Centipedegrass</i>									
0 (Presoaking)	-0.06	11.0 a	36.6 a	33.5 b	87.5 a	14.3 a	9.3 bc	5.7 ab	2.8 b
1	-0.5	12.1 a	32.8 a	35.0 b	72.0 b	13.4 a	8.0 c	4.6 b	3.2 b
2	-0.9	6.0 a	29.0 a	58.3 a	73.6 b	16.7 a	9.4 bc	5.7 ab	3.6 b
3	-1.3	4.0 a	26.7 a	36.4 b	62.0 bc	17.0 a	10.3 ab	6.4 ab	3.1 b
4	-1.7	5.0 a	21.0 a	30.0 b	57.0 c	14.6 a	11.0 a	5.4 b	3.1 b
Control	(untreated)	5.0 a	22.2 a	34.5 b	33.0 d	15.0 a	11.4 a	8.6 a	5.5 a
Linear		NS	NS	NS	NS	NS	NS	NS	NS
Nonlinear		NS	NS	NS	NS	NS	NS	NS	NS

^zMean separation within columns and species by Duncan's new multiple range test ($P \leq 0.05$).

ns, *, **Nonsignificant or significant at $P \leq 0.05$ and 0.01, respectively, within columns and species.

of primed seeds were calculated and compared with those of nonprimed seeds (control). MTG was calculated as follows: $MTG = (\sum Ti Ni) / G$, where Ti is the day of germination, Ni is the number of seeds germinating on Ti , and G is the total number of germinated seeds (Hartmann et al., 1990). A tetrazolium test was used to determine the viability of nongerminated seeds (Association of Official Seed Analysts, 1994). Seed germination was calculated based as the sum of percentages of viable seeds and germinated seeds.

The experimental design used for both grass species and temperature was a randomized complete-block with four replications. A general linear model was performed separately on each grass species and germination temperature, using the SAS statistical package (SAS Inst., 1991). Means were separated using Duncan's new multiple range test at $P \leq 0.05$.

Results and Discussion

Data for carpetgrass and centipedegrass data were evaluated separately. There were no significant differences over time for either species at $P \leq 0.05$. Therefore, the experiment was analyzed as a randomized complete-block design.

Common carpetgrass. Presoaking in distilled water did not significantly improve germination at 15 °C, but affected germination at higher temperatures (Table 1). Presoaking hastened germination significantly at 20 °C, but did not increase germination percentage. Presoaking improved germination percentage at 25 °C but not at 30 °C (Table 1), and reduced MTG at both temperatures by ≈ 2 d.

Priming in KNO₃ had a slight effect on germination, but did not reduce MTG at 15 °C (Table 1). The relationship between KNO₃ concentration and germination at 20 °C was linear. Mean time of germination was significantly decreased by ≈ 2 d in comparison to control seeds (Table 1). Priming with KNO₃

(1%, 2%, 3%, and 4%) and exposing to 25 °C significantly increased germination percentage and decreased MTG. Although germination percentage at 30 °C was not improved by priming, MTG was reduced by ≈ 2 d relative to the control.

Centipedegrass. There were no benefits to presoaking seed germinated at 15 or 25 °C (Table 1). Presoaking did not increase germination percentage at 20 °C, but accelerated MTG by ≈ 2 d. The highest germination percentage (87.5%) resulted from presoaking seed and germinating at 30 °C; MTG was also reduced ≥ 2.5 d.

Germination percentage at 15 °C decreased linearly as KNO₃ concentration increased (Table 1), but MTG was not significantly reduced by priming. Priming did not increase germination percentage at 20 °C, but MTG increased linearly with increasing KNO₃ concentration. Priming in 2% KNO₃ significantly increased germination by 23% at 25 °C. Mean time of germination was reduced by 3 d for seeds primed in either 1% or 4% KNO₃. Priming increased germination percentage at 30 °C and reduced MTG ≈ 2 d.

Conclusions

Presoaking and priming may be of little value at 15 °C, but priming common carpetgrass and centipedegrass seeds has potential for improving seed germination or reducing MTG at germination temperatures between 20 and 30 °C. Currently, there is no commercial source of primed or presoaked seed of carpetgrass or centipedegrass. However, the results of this study indicate that MTG of both species could be reduced by presoaking and/or priming.

Literature Cited

- Adegbuyi, E., S.R. Cooper, and R. Don. 1981. Osmotic priming of some herbage grass seed using polyethylene glycol (PEG). *Seed Sci. Technol.* 9:867-878.
- Akalehiyot, T. and J.D. Bewley. 1977. Promotion and synchronization of cereal grain germination

by osmotic pretreatment with polyethylene glycol. *J. Agr. Sci.* 89:503-506.

Association of Official Seed Analysts. 1994. Rules for testing seeds. *J. Seed Technol.* 16:1-113.

Bradford, K.J. 1986. Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *HortScience* 21:1105-1112.

Brede, J. and A.D. Brede. 1989. Seed priming. *Grounds Maintenance* 4:42, 46, 48.

Danneberger, T.K., M.B. McDonald, Jr., C.A. Geron, and P. Kumari. 1992. Rate of germination and seedling growth of perennial ryegrass seed following osmoconditioning. *HortScience* 27:28-30.

Delouche, J.C. 1961. Effect of gibberellin and light on germination of centipedegrass seed (*Eremochloa ophiuroides*). *Proc. Offic. Seed Analysts* 51:147-150.

Fry, J.D., W. Upham, and L. Leuthold. 1993. Seeding month and seed soaking affect buffalograss establishment. *HortScience* 28:902-903.

Hardegree, S.P. 1994. Matric priming increases germination rate of Great Basin native perennial grasses. *Agron. J.* 86:289-293.

Hartmann, H.T., D.E. Kester, and F.D. Davies, Jr. 1990. *Plant propagation: Principles and practices*, 5th ed. Prentice Hall, Englewood Cliffs, N.J.

Heydecker, W. and P. Coolbear. 1977. Seed treatments for improved performance: Survey and attempted prognosis. *Seed Sci. Technol.* 5:353-425.

Lush, W.M. and J.A. Birkenhead. 1987. Establishment of turf using advanced 'pregerminated' seeds. *Austral. J. Expt. Agr.* 27:323-337.

Maguire, J.D. and K.M. Steen. 1971. Effects of potassium nitrate on germination and respiration of dormant and nondormant Kentucky bluegrass (*Poa pratensis* L.) seed. *Crop Sci.* 11:48-50.

McClure, G. 1995. Sweet corn seed reserve changes during imbibition in relation to germination at low temperature. MS Thesis, Louisiana State Univ., Baton Rouge.

Murray, G. 1990. Priming sweet corn seed to improve emergence under cool conditions. *HortScience* 25:231.

SAS Institute. 1991. SAS users guide. SAS Inst., Cary, N.C.

Toole, E.H. and V.K. Toole. 1939. Notes: Germination of carpetgrass seed. *Amer. Soc. Agron.* 31:566-567.

Walker, J.T. 1976. Centipedegrass seed treatments and light-temperature effects on germination. *Plant Dis. Rptr.* 60:393-397.