Large-scale Micropropagation of *Aloe vera*

Enio Tiago de Oliveira¹ and Otto Jesu Crocomó

ESALQ/USP, Biological Science, Av. Pádua Dias, 11, Piracicaba, São Paulo 13418-900 PB09, Brazil

Tatiana Bistaco Farinha

Depto. de Ciências Biológicas, Laboratório de Controle Hormonal do Desenvolvimento Vegetal, Escola Superior de Agricultura Luis de Queiroz, ESALQ/USP, C.P. 9, 13418–900, Piracicaba, SP, Brazil

Luiz Antônio Gallo

Depto. de Ciências Biológicas, Laboratório de Biotecnologia Agrícola (CEBTEC), Escola Superior de Agricultura Luis de Queiroz, ESALQ/USP, C.P. 9, 13418–900, Piracicaba, SP, Brazil

Additional index words. *Aloe*, plant tissue culture, disinfection, organogenesis, plant micropropagation

Abstract. A protocol for large-scale *Aloe vera* (L.) Burm. f. production was established using micropropagation of apical buds. The effects of two chlorine-based disinfectants were evaluated on the survival of the explants in different treatments in a semisolidified Murashige and Skoog (MS) medium in the presence of 6-benzylaminopurine (6-BAP; 2 mg·L⁻¹). During 120 days, 136 green apical shoots bearing axillary buds were multiplied four times at 30-day intervals in the same MS medium, reinoculating seven to nine explants per flask each time. The elongation and rooting processes were carried out in the same MS medium without 6-BAP. A total of 40,495 *Aloe vera* microplants were obtained, a yield of 300 microplants per apical bud at a rate of 1:5.3 in every multiplication period of 30 days. From that total, 38,480 *Aloe vera* microplants were successfully acclimatized transferring to 36- and 64-cell polyethylene trays containing proper substrate in two different ex vitro greenhouse conditions. After a 3-month period, fresh and dry matter weights of the *Aloe vera* plants were determined. All the data from each experimental phase were statistically analyzed. The use of 64-cell (40 cm³/cell) trays represented an economy of 47.37% in greenhouse space and 50% in the amount of substrate per *Aloe vera* plant.

Aloe vera (L.) Burm. f., belonging to the family Asphodelaceae (Souza and Lorenzi, 2005), is one of a few *Aloe* species that has been explored by pharmaceutical and cosmetics industries (Mapp and McCarthy, 1970; Morton, 1961). Biological activities are ascribed to the pulp gel of *Aloe vera* such as antiviral, antibacterial, antifungal, anticancer, wound healing, and many others (Reynolds and Dweck, 1999), the medicinal principles being phenolics and polysaccharide compounds (Okamura et al., 1996; Park et al., 1998). Those qualities have prompted industrial and commercial increase in the production of *Aloe vera* throughout the world. The main constraint in this production chain is to obtain healthy primary culture for a sustainable supply of the plant materials. The common practice for the reproduction of

Aloe is conventional vegetative propagation. However, this practice should be avoided because it can also allow the propagation of diseases frequently present in the mother plants, because it happens during the conventional vegetative propagation of other plant species such as banana, potatoes, strawberry, and sugarcane. This problem can be minimized using in vitro propagation of stock materials (Crocomó and Oliveira, 1995; Debergh and Read, 1991) with the objective of the production of not only healthy, but also standardized sizes of plantlets to be used in the subsequent controlled stages of the process suitable for the clonal production of *Aloe vera*. In this context, this article describes a complete micropropagation system involving disinfection, in vitro multiplication, rooting, and hardening followed by ex vitro acclimatization procedures used to attain that objective, thus producing thousands of *Aloe vera* plants.

Materials and Methods

Plant material sources. *Aloe vera* plants were obtained from the field collection of the Department of Plant Production of the Escola Superior de Agricultura “Luiz de Queiroz,”

University of São Paulo, Piracicaba, SP, Brazil.

Surface disinfection treatments. Four hundred eighty apical buds explants, each ≈1 cm³, were isolated from young lateral shoots bearing six to nine leaves and disinfected using three different disinfection treatments (T₁, T₂, and T₃) as shown in Table 1. In each treatment, the final step (Step 7) was conducted in an air sterile chamber followed by removal of the external tissue layers of the explants up to ≈1 cm³ and inoculated into the shoot proliferation medium (SPM) based on semisolidified Murashige and Skoog medium (Murashige and Skoog, 1962) with 2 mg·L⁻¹ glycine, 100 mg·L⁻¹ myo-inositol, 30 g·L⁻¹ sucrose, and 2 mg·L⁻¹ 6-benzylaminopurine (6-BAP); the pH was adjusted to 5.8 before adding 2.3 g·L⁻¹ Phytagel® (SIGMA-ALDRICH Inc., St Louis, MO). In each treatment, 160 buds were individually inoculated in 12.5 cm diameter × 8 cm long test tubes containing 12.5 mL of the SPM medium and incubated in growth room at 25 °C ± 2 under light (50 μmol·m⁻²·s⁻¹) with a photoperiod of 16/8 h. The effect of the disinfection treatments was evaluated after 30 d in the culture medium taking into account the rates of explant contamination, necrosis, and green apical shoot (GAS) development.

Shoot proliferation. After 30 d in the SPM medium, the best 136 green apical shoot explants bearing axillary buds (GAS) were selected, longitudinally divided, and transferred from the test tubes to 350-mL glass flasks containing 50 mL of the same medium and cultivated for an additional 90 d under the same experimental conditions with replication at 30-d intervals. Seven to nine explants were transferred in each replication.

Elongation and root induction. Developed *Aloe* plantlets were transferred to culture medium without 6-BAP to induce elongation and rooting for a 30-d period.

Table 1. Sequential procedures for *Aloe vera* explant disinfection.

Treatments	Steps							
	0	1	2	3	4	5	6	7
T ₁	—	H	H	H	H	H	H	H
T ₂	H	S	S	S	H	H	H	H
T ₃	S	S	H	S	S	H	H	H

H = sodium hypochlorite (0.5% v/v active chlorine) solution; S = 0.06% (v/v) solution prepared from the sodium dichloroisocyanurate (Sumaveg® Unilever Brasil Ltda, São Paulo, SP) with 3% (v/v) active chlorine solution; Step 0 = only for T₂ and T₃ treatments: previous shoot washing with the respective solution; Step 1 = shoot defoliation (three to four leaves) followed by immersion into the respective solution, ≈30 minutes; Step 2 = shoot defoliation of the remaining leaves followed by immersion into the respective solution, ≈30 minutes; Step 3 = rinsing with the respective solution; Step 4 = resting with the respective solution, ≈2 h; Step 5 = rinsing with the respective solution; Step 6 = rinsing with the respective solution prepared with sterile water; Step 7 = final disinfection by immersion into the respective solution prepared with sterile water for 30 minutes. In each treatment 160 replicates were used.

Received for publication 18 June 2009. Accepted for publication 6 Aug. 2009.

We thank Prof. Keigo Minami, Department of Plant Production/E.S.A. “Luiz de Queiroz”/University of São Paulo, for supplying selected *Aloe vera* plants.

¹To whom reprint requests should be addressed; e-mail etolivei@esalq.usp.br.

Fig. 1. Effect of the disinfection treatments on the survival of apical shoots of *Aloe vera*. Analysis was made using 480 explants.

Fig. 2. Classification of *Aloe vera* microplants by size: (A) Small size (M_s). (B) Medium size (M_m). (C) large size (M_l).

Hardening and acclimatization. The average rate of *Aloe* microplant production was evaluated following the in vitro proliferation; 38,480 microplants were used in the hardening and acclimatization processes. The microplants were classified as: small (M_s), 3 to 4 cm long bearing two to three leaves; medium (M_m), 6 to 8 cm long bearing four to five leaves; and large (M_l), 13 to 16 cm long bearing five to seven leaves (Fig. 1). The microplants were transferred to two different types of black polyethylene trays labeled B₁ (80 cm³/cell) and B₂ (40 cm³/cell) with 36 and 64 cells each, respectively. The trays were maintained in two different types of greenhouses with automatically controlled relative humidity and proper irrigation namely: G₁, pad-house system using microaspiration irrigation and air exhaustion; and G₂, conventional greenhouse using nebulization and forced air ventilation. A mixture of bark pinus, vermiculite, sand, and fertilizers (Terra do Paraíso™; Substratos Agrícolas Ltda, Holambra, SP), pH 6.04 and electric conductivity

Fig. 3. Microplant survival as a function of their sizes. M_s = small size; M_m = medium size; M_l = large size. Columns followed by at least one same letter did not differ among them by Tukey test ($P < 0.05$). Each value is the average of 40 replicates.

0.62 $\mu\text{S}\cdot\text{cm}^{-1}$, was used as substrate. After a period of 3 months, the rate of microplant survival and their fresh and dry matter contents were determined.

Statistical analysis. The data were submitted to the analysis of variance (5% probability) using the Statistical Analysis Systems (Version 8.0; SAS, Cary, NC). The average data for the rates of contamination, necrosis, and green apical shoot production were compared by least significant differ-

ence. The Tukey test was used for hardening and acclimatization, comparing microplant sizes $\times$ tray types $\times$ greenhouse conditions.

Results and Discussion

Disinfection, shoot differentiation, and rooting. The three disinfection treatments were effective in different degrees for disinfecting the *Aloe vera* apical buds in vitro. The statistical analysis of the effect of the disinfection processes on the rates of contamination, mainly by bacteria, and on the green apical shoots with axillary buds (GAS) production (Fig. 2) showed no significant differences ($P > 0.05$) between T₂ and T₃ treatments, both being different from T₁ ($P < 0.05$). In relation to the contamination and the GAS rates, the best results were obtained by T₃ (13.12% for contamination and 32.50% GAS) and T₂ (15% for contamination and 35% for GAS) and T₁ (32.81% for contamination and 17.50% for GAS). This was attributable mainly to washing the start material in the disinfectant solutions at the beginning of the process (Step S0 or H0). In regard to necrosis of the explants (Fig. 2), no significant differences were observed among the three treatments: T₁, 49.69%; T₂, 50.00%; and T₃, 54.38%.

The degrees of injury on the explants depend on the type of disinfection processes and the nature of the plant material. The *Aloe vera* tissue has low lignin content and is very sensitive to disinfectants. This is not likely to be the result of chlorine toxicity, either of the hypochlorite or the sodium dichlorisocyanurate, which is largely used to disinfect “fresh-cut” products (Lund et al., 2005). One can assume that this could be attributed to the alkalinity of the disinfectant solutions. This aggressiveness was observed mainly in the rates of necrosis of the explants.

From 480 inoculated explants, 136 green apical shoots with axillary buds (GAS) were multiplied three times at 30-d intervals in the same SPM medium. The shoots were longitudinally divided and seven to nine explants were reinoculated per flask in the same culture medium each time.

After the stage of multiplication, the plantlets were separated and transferred to the previous culture medium without 6-BAP to induce elongation and rooting. At the end of the multiplication and rooting periods,

Table 2. Effect of acclimatization conditions on the increase of fresh matter weight (g) of *Aloe vera* microplants.

Microplants	Greenhouse 1 (G ₁)		Greenhouse 2 (G ₂)	
	Tray 1 (B ₁)	Tray 2 (B ₂)	Tray 1 (B ₁)	Tray 2 (B ₂)
M_s	10.49 aA*	8.60 aA*	8.36 aA*	7.03 aA*
M_m	15.58 aA*	12.01 aA*	15.39 aA†	9.66 aA*
M_l	32.50 aA†	20.55 bA†	21.69 aB†	18.86 aA†

Averages followed by at least the same letter or symbol did not differ by the Tukey test ($P < 0.05$). Lower case letters: comparison for trays (constant microplants and greenhouse). Upper case letters: comparison for greenhouses (constant microplants and trays). Symbols (* †): comparison in the same column for microplant size (constant tray and greenhouse).

M_s = Small size microplants; M_m = medium size microplants; M_l = large size microplants; G₁ = pad-house type greenhouse with microaspiration irrigation; G₂ = greenhouse with nebulization irrigation and forced air; B₁ = 36-cell trays (80 cm³/cell); B₂ = 64-cell trays (40 cm³/cell). Each value is the average of 10 replicates.

Table 3. Effect of acclimatization conditions on the increase of dry matter weight (g) of *Aloe vera* microplants.

Microplants	Greenhouse 1 (G ₁)		Greenhouse 2 (G ₂)	
	Tray 1 (B ₁)	Tray 2 (B ₂)	Tray 1 (B ₁)	Tray 2 (B ₂)
M _s	0.20 aA*	0.16 aA*	0.18 aA*	0.15 aA*
M _m	0.29 aA*	0.24 aA*†	0.35 aA†	0.21 aA*
M _l	0.60 aA†	0.38 bA†	0.48 aA†	0.39 aA†

Averages followed by at least the same letter or symbol did not differ among them by the Tukey test ($P < 0.05$). Lower case letters: comparison for trays (constant microplants and greenhouse). Upper case letters: comparison for greenhouses (constant microplants and trays). Symbols (* †): comparison in the same column for microplant size (constant tray and greenhouse).

M_s = Small size microplants; M_m = medium size microplants; M_l = large size microplants; G₁ = greenhouse with pad-house and microaspiration irrigation; G₂ = greenhouse with nebulization irrigation and forced air; B₁ = 36-cell trays (80 cm³/cell); B₂ = 64-cell trays (40 cm³/cell). Each value is the average of 10 replicates.

Fig. 4. General view of *Aloe vera* microplants acclimatization in pad-house system greenhouse with microaspiration irrigation (G₁). A total of 800 m² was suitable for 3330 trays.

Fig. 5. *Aloe vera* microplants: 3 months acclimatization in a pad-house system greenhouse with microaspiration irrigation (G₁).

40,495 *Aloe vera* plantlets were obtained, representing a yield of 300 microplants per apical shoot, or a rate of 1:5.3 in every multiplying period of 30 d.

Similar results were observed by Marfori and Malasa (2005) testing the effects of 6-BAP and kinetin on *Aloe vera* development

in vitro; in the presence of 2.3 mg·L⁻¹ 6-BAP, the ratio was 1:11.3 and in the presence of 1 mg·L⁻¹ 6-BAP, it was 1:9. Liao et al. (2004) obtained a ratio of 1:15 for the micropropagation of chinese *Aloe* in the presence of 2 mg·L⁻¹ 6-BAP plus 0.3 mg·L⁻¹ naphthalene-acetic acid (NAA). Araújo et al. (2002)

obtained ratios varying from 1:2 up to 1:8, depending on the genotype, using 1.5 mg·L⁻¹ 6-BAP and 0.75 mg·L⁻¹ NAA in the in vitro multiplication of 16 different *Aloe vera* genotypes.

Hardening and acclimatization. After a period of 3 months of acclimatization, the rate of microplant survival and their fresh and dry matter contents were determined. In terms of rate of microplant survival, there was a statistically significant difference ($P < 0.05$) observed only in size difference. No other factor such as greenhouse type and tray type, nor their interactions, showed significant differences ($P > 0.05$). As far as size is concerned, medium (M_m) and large (M_l) were found to be significantly different from small (M_s) microplants ($P < 0.05$) (Fig. 3).

The statistical analysis showed a positive interaction among the rate of survival, the content of fresh matter weight (FW), and dry matter weight (DW) ($P < 0.05$) during acclimatization. Tables 2 and 3 show the data for FW and DW for the three microplant sizes, respectively. Fresh matter weight content was higher in large microplants (M_l) maintained in 36-cell trays under microaspiration (G₁) than in the other sizes and also higher than in the M_l microplants maintained under nebulization in 36- or 64-cell trays. Figure 4 shows a general view of the *Aloe vera* microplant acclimatization under microaspiration irrigation in a 800-m² greenhouse with a pad-house system suitable for 3300 trays. Figure 5 shows large *Aloe vera* microplants (M_l) after 3 months of acclimatization. Considering that number of cells in the trays showed no influence on the rate of microplants survival and FW and DW content, the choice of the number of cells is a practical one: using 64-cell trays (40 cm³/cell), there is an economy of 47.37% in greenhouse space and 50% in the amount of substrate used per microplant.

Conclusions

At least 300 microplants were produced from a single apical bud of *Aloe vera* in a period of 4 months. The best results of the apical shoot disinfection were obtained when the field-harvested *Aloe vera* stocks were previously rinsed with a solution either of sodium hypochlorite or sodium dichloroisocyanurate.

The *Aloe vera* plantlets were successfully ex vitro acclimatized in a greenhouse bearing a pad-house system with microaspiration and air exhaustion using 36- (80 cm³/cell) or 64- (40 cm³/cell) cell polyethylene trays containing the proper substrate. Because the cell number in the trays showed no influence either on the survival of the plants or on the FW and DW, the use of 64-cell (40 cm³/cell) trays represents an economy of 47.37% in greenhouse space and 50% in the amount of substrate per *Aloe vera* plant.

Literature Cited

Araújo, P.S., J.M.O.D. Silva, C.A. Nechel, C. Ianssem, A.C. Oltramari, R. Passos, E. Tiepo,

- D.B. Bach, and M. Maraschin. 2002. Micropropagação de babosa (*Aloe vera*—*Liliaceae*). Biotecnologia Ciência e Desenvolvimento, Brasília, No. 25, p. 54–57.
- Crocomo, O.J. and E.T. Oliveira. 1995. Biotecnologia de plantas como um programa de produção vegetal. In: Gerald, L.T.S. (ed.). Biofábrica Produção Industrial de Plantas 'In Vitro'. Araras: UFSCar. p. 18–21.
- Debergh, P.C. and P.E. Read. 1991. Micropropagation, p. 1–13. In: Debergh, P.C. and R.H. Zimmerman (eds.). Micropropagation—Technology and application. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Liao, Z., M. Chen, F. Tan, X. Sun, and K. Tang. 2004. Micropropagation of endangered Chinese aloe. Plant Cell, Tissue and Organ Culture 76:83–86.
- Lund, D.G., L.A. Petrini, J.A.G. Aleixo, and C.V. Rombaldi. 2005. Uso de sanitizantes na redução da carga microbiana de mandioca minimamente processada. Ciência Rural, Santa Maria, 35(6):1431–1435.
- Mapp, R.K. and T.J. McCarthy. 1970. The assessment of purgative principles in Aloes. Planta Medica, New York 18:361–365.
- Marfori, E.C. and A.B. Malasa. 2005. Tissue culture for rapid clonal propagation of *Aloe barbadensis* Miller. Philippine Agricultural Scientist, Laguna 88:167–170.
- Morton, J.F. 1961. Folk Uses and commercial exploitation of Aloe leaf pulp. Economic botany Bronx 15:311–319.
- Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiologia Plantarum, Copenhagen 15:473–497.
- Okamura, N., M. Asai, N. Hine, and A. Yagi. 1996. High-performance liquid chromatographic determination of phenolic compounds in *Aloe* species. J. Chromatography 746: 225–231.
- Park, M.K., J.H. Park, N.Y. Kim, Y.G. Shin, Y.S. Choi, J.G. Lee, K.H. Kim, and S.K. Lee. 1998. Analysis of 13 phenolic compounds in *Aloe* species by high performance liquid chromatography. Phytochemical Analysis Sussex 9:186–191.
- Reynolds, T. and A.C. Dweck. 1999. *Aloe vera* leaf gel: A review update. Journal of Ethnopharmacology, Lausanne 68:3–37.
- Souza, V. and H. Lorenzi. 2005. Botânica Sistemática: Guia ilustrado para identificação das famílias de Angiospermas da flora brasileira, baseado em APG II. Instituto Plantarum, Nova Odessa, 640 p.