Development of Simple Sequence Repeat Markers for *Chionanthus retusus* (Oleaceae) and Effective Discrimination of Closely Related Taxa

Renée S. Arias

USDA-ARS Genomics and Bioinformatics Research Unit, Stoneville, MS 38776

Natascha Techen

University of Mississippi, National Center for Natural Products Research, School of Pharmacy, University, MS 38677

Timothy A. Rinehart

USDA-ARS Southern Horticultural Laboratory, Poplarville, MS 39470

Richard T. Olsen

U.S. National Arboretum, USDA-ARS, Washington, DC 20002

Joseph H. Kirkbride, Jr.

U.S. National Arboretum, USDA-ARS, Washington, DC 20002

Brian E. Scheffler¹

USDA-ARS Genomics and Bioinformatics Research Unit, Stoneville, MS 38776

Additional index words. DNA fingerprinting, STR markers, *Chionanthus pygmaeus*, *Chionanthus virginicus*, *Osmanthus americanus*, fringetree

Abstract. The genus *Chionanthus* (Oleaceae Hoffmans. & Link) includes deciduous or evergreen trees and shrubs distributed widely in tropical and sub-tropical areas, including a few temperate species. Although *Chionanthus* species are planted as ornamental garden plants and commercialized for natural products, genetic information for *Chionanthus* spp. is lacking. We created microsatellite-enriched libraries of *Chionanthus retusus* Lindl. & Paxton, assembled 1072 contigs, and detected 1010 repeats. The frequency of the repeats decreased with the increase in repeat length, and the most abundant motifs were: AG, AC, AAG, ACC, AT, and ACTC. We screened 384 markers on 12 accessions of four related taxa that included *C. retusus*, *Chionanthus virginicus* L., *Chionanthus pygmaeus* Small, and *Osmanthus americanus* (L.) Benth. & Hook. A total of 195 simple sequence repeat (SSR) markers amplified and discriminated six accessions of *C. retusus* and 57 SSR markers amplified and discriminated across the four Oleaceae species screened. To identify the best markers to use in future experiments, the “Unique Pattern Informative Combination” (UPIC) values were calculated for all the markers and the 100 markers that were most effective are reported here. The percentage of heterozygous loci across the 384 markers was lowest for *C. retusus* (29.3%) and highest for *O. americanus* (68.9%). The SSR markers developed here could assist in taxonomy and hybridization investigations for breeding programs and authentication of varieties used as medicinal plants.

Received for publication 15 Oct. 2010. Accepted for publication 11 Nov. 2010.

This work was supported by USDA-ARS project no. 6401-21000-001-00 and 6402-21310-003-00. We thank Xiaofen (Fanny) Liu for library sequencing. Mention of trade names or commercial products in this manuscript is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

¹To whom reprint requests should be addressed; e-mail brian.scheffler@ars.usda.gov.

Chionanthus L. is a member of the Oleaceae, a morphologically diverse family that includes economically important genera cultivated for food (*Olea europaea* L.), timber (*Fraxinus* L.), medicinal (*Forsythia* Vahl.), and ornamental (*Jasminum* L., *Ligustrum* L., *Osmanthus* Lour., and *Syringa* L.) purposes. The genus *Chionanthus* has ≈100 species (Chang et al., 1996; Wallander and Albert, 2000) distributed throughout tropical and subtropical areas worldwide but includes three temperate species (Green, 2004). Only

the three temperate species are economically important as ornamentals and natural products. Temperate *Chionanthus* spp. (*C. retusus* from eastern Asia and *C. pygmaeus* and *C. virginicus* from eastern North America) are cultivated for their white flowers in feathery panicles and decorative blue fruits. With the exception of the pink-flowered *C. pubescens* Kunth from Ecuador, the evergreen tropical and subtropical *Chionanthus* spp. (syn *Lino-ciera* Sw.) are usually not cultivated.

Very little is known about the genetics of *Chionanthus*. The base chromosome number in tribe Oleae is $x = 23$ with published counts for *C. virginicus* and *C. retusus* of $2n = 46$ (Chang et al., 1996; Taylor, 1945; Wallander and Albert, 2000). A hypothetical phylogeny for the Oleaceae, including *C. retusus* and *C. virginicus*, was developed using sequences of two noncoding chloroplast regions, the *trnL-F* (intron-L/spacer-F) and *rps16* intron (Wallander and Albert, 2000); however, no other information at the molecular level is available that could assist in *Chionanthus* breeding programs or identification of species. In addition, understanding the level of genetic diversity as well as inter- and intrageneric relationships within the cultivated germplasm is important for planning a systematic breeding program for *Chionanthus*. Furthermore, the pollination and breeding systems of *Chionanthus* appear complex with individual species exhibiting varying levels of polygamodioecy (Dirr, 1998; Nicholson, 1990; Ueda, 1996) combined with occasional selfing that has implications for crossing and inheritance studies.

The development of molecular markers, specifically SSRs, will aid in assessing relationships, diversity, and parentage within the genus *Chionanthus*. In addition, because SSR marker transferability is usually feasible within plant families, i.e., Casuarinaceae (Yasodha et al., 2005), Meliaceae (White and Powell, 1997), Fagaceae (Barreneche et al., 2004), the markers we report here for *Chionanthus* could potentially transfer to species of *Olea*, *Fraxinus*, and *Syringa*.

Materials and Methods

Plant material. Source and accession data for all samples used in this study are shown in Table 1. *Chionanthus retusus*-derived SSR markers were tested on six accessions of *C. retusus*, three accessions of *C. virginicus*, two accessions of *C. pygmaeus*, and one accession of *Osmanthus americanus*, a species related to *Chionanthus* according to Wallander and Albert (2000). Total genomic DNA was extracted from leaf tissue using a Qiagen Plant Mini Kit (Qiagen, Valencia, CA).

Isolation of simple sequence repeats from *Chionanthus retusus*. For the construction of SSR-enriched libraries, we used the method of Techen et al. (2010), briefly described here. DNA from *C. retusus* was cut with restriction enzymes *AluI*, *HaeIII*, *DraI*, *RsaI* (New England Biolabs, Ipswich, MA) and combinations of pairs of these enzymes. The blunt-end DNA fragments were A-tailed with Taq-DNA polymerase and then ligated to an optimized

Table 1. *Chionanthus* and *Osmanthus* samples tested with simple sequence repeats developed from *Chionanthus retusus*.²

Species	Accession no.	Source and provenance	Voucher no.	Sample
<i>Chionanthus retusus</i>	37681-HL	Kunming Inst. Bot.	6013	CR1
<i>Chionanthus retusus</i>	—	Cultivated, Greenbelt, MD	6021	CR2
<i>Chionanthus retusus</i>	37681-HH	Kunming Inst. Bot.	6012	CR3
<i>Chionanthus retusus</i>	45247-J	Coastal Japan 1978	6020	CR4
<i>Chionanthus retusus</i>	14089-L	Arnold Arboretum	6006	CR5
<i>Chionanthus retusus</i>	40121-H	Kyoto Univ. Forest Stat.	6016	CR6
<i>Chionanthus virginicus</i>	—	BARC-E, Beltsville, MD	6002	CV8
<i>Chionanthus virginicus</i>	—	BARC-E, Beltsville, MD	6003	CV9
<i>Chionanthus virginicus</i>	77196-H	Shemin Nurs.	6007	CV10
<i>Chionanthus pygmaeus</i>	76993-3	U.S. Botanic Garden	—	CP11
<i>Chionanthus pygmaeus</i>	76993-2	U.S. Botanic Garden	—	CP12
<i>Osmanthus americanus</i>	56806-H	Tom Dodd Nurs.	6008	OA7

²All voucher specimens are located in the herbarium of the U.S. National Arboretum, Washington, DC, and are collections of J. Kirkbride and R. Olsen. Living plants of *C. pygmaeus* are only listed by accession number because no voucher specimens have been collected.

linker (SSRLIBF3: 5'-CGGGAGAGCAAGG AAGGAGT-3', SSRLIBR3: 5'-/5Phos/CTCC TTCCTTGCTCTCCCGAAAA'-3'). After 20 cycles of polymerase chain reaction (PCR), the amplified products were hybridized to four groups of biotinylated oligo repeats. Sequences containing repeats were captured using streptavidin-coated magnetic beads M-270 (Invitrogen, Carlsbad, CA) and the DNA was eluted. The eluate was PCR-amplified for 20 cycles; the PCR products were cloned in vector TOPO4 (Invitrogen) and sequenced using an ABI 3730XL DNA Analyzer (Applied Biosystems, Foster City, CA). Sequences were assembled in contigs using DNASTar Lasergene7 (DNASTAR, Inc., Madison, WI) and visually checked. Repeats were searched using SSRFinder (Sharopova et al., 2002) and Sputnik (Abajian, 1994). Primers were designed using Primer3 (Rozen and Skaletsky, 2000) with stringent parameter conditions: Tm 63 optimum (60/65) minimum/maximum, length 24 optimum (20/28) minimum/maximum, and maximum overlap of repeat within the primer was 5 bp. Contig sequences containing microsatellites were screened against the NCBI Protein Database (BLASTx) (Altschul et al., 1990).

Fingerprinting. Forward primers were 5' tailed with the sequence 5'-CAGTTTTCCC AGTCACGAC-3' (Waldbieser et al., 2003) to permit product labeling, and reverse primers were tailed at the 5' end with the sequence 5'-GTTT-3' to promote non-template adenylation (Brownstein et al., 1996). Primer 5'-CA GTTTCCCAGTCACGAC-3' labeled with 6-carboxy-fluorescein (IDT-Technologies, Coralville, IA) was used for amplification of 10 ng DNA using Titanium Taq DNA Polymerase (Clontech, Mountain View, CA) in 5-μL reactions on an M&J thermal cycler (BioRad, Hercules, CA) at 95 °C for 1 min, 60 °C for 1 min (two cycles), 95 °C for 30 s, 60 °C for 30 s, 68 °C for 30 s (27 cycles), and a final extension at 68 °C for 4 min. Fluorescently labeled PCR fragments were analyzed on an ABI 3730XL DNA Analyzer and data-processed using GeneMapper Version 3.7 (both from Applied Biosystems). Presence of alleles was converted to a binary matrix. The accessions were clustered using the unweighted paired group method and arithmetic averages (UPGMA) algorithm implemented in

the SAHN program of NTSYSpc Version 2.2 (Exeter Software, Setauket, NY). The confidence level for the dendrograms was assessed by bootstrap resampling (5000 replicates) (Efron et al., 1996; Felsenstein, 1985) using WINBOOT (Yap and Nelson, 1996).

Unique Pattern Informative Combination and heterozygous loci calculations. Markers that effectively discriminated the samples tested were identified using the UPIC software (Arias et al., 2009), and the number of unique patterns (UPIC values) identified by each marker was reported for the best 100 markers. Percentage of heterozygous loci was also calculated using the UPIC software (Arias et al., 2009) for each DNA sample across all 384 SSR markers tested.

Results

Repeats found. SSR-enriched libraries of *Chionanthus retusus* were made using four groups of biotinylated oligo repeats. A total of 2208 clones from those libraries was sequenced. Sequences were assembled to 1072 contigs in which 1010 repeats were detected by SSRFinder and Sputnik combined. We considered repeats only those that were non-monomonucleotides, had a minimum repeat length of 8 bp, and a minimum 20-bp length of flanking region upstream and downstream of the repeat. Sequences of contigs containing repeats were submitted to GenBank with accession numbers (GQ117288 to GQ118148). We designed 394 primers on the flanking regions of the repeats and tested 384 of those primers on 12 DNA samples. Primer sequences and repeat motifs as appeared in the original sequences are provided in Table 2. DNA sequences corresponding to 28 of the markers had significant hits on BLASTx indicated in Table 2 in bold. To simplify the recording of the repeat motifs, those that were circular permutations and reverse complements of each other were grouped together as one type, i.e., AAC, ACA, CAA, GTT, TGT, and TTG were recorded as AAC. This resulted in 55 non-redundant repeat motifs isolated from the *C. retusus* SSR-enriched libraries. The first 11 most abundant motifs detected had frequencies ranging from 560 to six as shown in Figure 1. The remaining 44

non-redundant motifs, including one CG repeat, were found in a total of 57 repeats (Table 2) with frequencies lower than four and were not included in the plot (data not shown). Frequencies of the isolated repeats decreased as their length increased. The number of di-, tri-, tetra-, penta-, and hexanucleotides are shown in Figure 1. Only 10 of 384 markers tested did not amplify any of the 12 DNA samples used in this study.

Markers that amplified four Oleaceae species (genera *Chionanthus* and *Osmanthus*). Based on the electropherograms in GeneMapper and presence across samples, we selected 57 markers that amplified across the four species tested (*C. pygmaeus*, *C. retusus*, *C. virginicus*, and *O. americanus*). Selected primers are listed in Table 3. From those, 43 amplified all 12 DNA samples and 14 amplified nine or more DNA samples. Only two of the 57 markers were monomorphic for the four species tested; the other 55 corresponded to polymorphic loci and amplified between two and 13 alleles. A total of 350 alleles was detected by these 57 markers. From these markers, 18 were polymorphic (detected up to five alleles) within the species *C. pygmaeus* allowing distinction of the two accessions tested, whereas 35 were polymorphic (detected up to five alleles) within the species *C. virginicus* allowing distinction of the three accessions tested.

Because the SSRs in this study were isolated from *C. retusus*, most markers amplified predominantly this species. Of 384 markers tested, based on their ease to score, we selected 195 that amplified the six lines of *C. retusus*; 33 of them were monomorphic, whereas the other 162 polymorphic markers detected between two and 15 alleles. A total of 837 alleles was detected in these 195 loci within the species *C. retusus*.

Unique pattern informative combinations of markers, Unique Pattern Informative Combination values. UPIC indicates the number of DNA samples that can be discriminated by each particular marker (Arias et al., 2009). The UPIC value is more informative for selecting subsets of SSRs than the use of its polymorphism information content. We calculated UPIC values for all the markers and report the UPIC values of the best 100 in Table 3. Twelve DNA samples (Table 1) representing four species of Oleaceae (*Chionanthus retusus*, *C. virginicus*, *C. pygmaeus*, and *Osmanthus americanus*) were used for the analysis. Combinations of these markers run on the 12 DNA samples described can detect as many unique patterns or alleles as to the sum of their UPIC values. In general, the markers allowed a clear distinction among *Chionanthus* taxa as well as within *C. retusus* accessions, which was shown by the high bootstrap resampling coefficients obtained (Fig. 2).

Heterozygosity (%). The percentage of heterozygous loci, based on 384 markers, for each of the four Oleaceae species tested was 29.9% to 30.7% (*C. pygmaeus*), 30.1% to 39.8% (*C. retusus*), 29.3% to 43.4% (*C. virginicus*), and 68.8% for *O. americanus*.

Cluster analysis. Genetic similarity coefficients based on UPGMA were calculated

Table 2. Markers that amplified all the Oleaceae accessions tested detected by Unique Pattern Informative Combination software (Arias et al., 2009).^z

Marker ID	Forward primer (5' → 3')	Reverse primer (5' → 3')	Size range	No. all	Max all/S	Motif
StvChR_15_a	GGAAAAAGAAAGGGAGAAGGAGAA	CTCTGTGACCATGACTGTCTGTGA	114–212	7	3	AG
StvChR_94_a	CGGAGACAATTTAAAGCACGATTC	CGACAATAGTTCAAGCATTGCGTA	151–188	10	5	TTC
StvChR_76_a	TCAGTCTTCACTACCACTACCACGTA	TGGGCTTTTAGACGAGTATTTGGA	105–165	10	2	ATT
StvChR_114_a	CCTTCCCCATTTAATCAATCACAA	TATGTTTGTTTACTTGTGCCCGTG	160–165	4	2	AC
StvChR_124_a	AATTTCTAGCCAATCACCTCATTC	ATGACGTGACCTTTTGTAGAGGAG	110–132	6	3	TC
StvChR_140_a	TTCTTCGCTCTGTAAAATTGGTCC	CCCTCCTTCTGTGAAGTATGACT	151–161	5	2	TG
StvChR_194_a	CAAGATCTGTTTGGGTCATTTCG	CTCCTTTGAAGCTTTCAACCCTTA	167–172	5	2	TC
StvChR_221_a	GGCTGAGGTTGATTACCTCTGATT	TTTCAGCCTGTAAAACCCTACTCTC	82–110	10	2	AG
StvChR_251_b	TTCTGTTTATTCATCTTCCACCGC	TAGACCAGGGTATTGGTCTTGCAT	132–191	9	4	TG
StvChR_284_a	AAAAATGTTGTGGTAGAAGTGCCG	GTTCTCAACTCCCGGATCTTAAT	69–225	9	4	GGC
StvChR_285_a	TTGACAGTGAAGGTGCAGAAGAAG	AGCCAGTCAGTGTATTCTCCAGG	143–144	2	1	AAG
StvChR_291_a	TGCATTTTCGTTACATTTTGTACGG	AGGATAATCACAGGAGGAAGGGAG	110–389	9	3	TC
StvChR_316_a	TCTCTTTCGGCTTCTTTTCTTCT	GCTACCAACATCATTCGTTCTCT	172–193	9	3	TTC
StvChR_337_a	AGAAGGCTACGGAATTGTAAAGG	GCTAAGTAAGGGGGTTCGGACTAAG	104–174	2	1	TC
StvChR_350_a	GCTTCAACAATCTCCAAAATGTCT	AGGCAACGTGATCTCAGTCTTTT	121–196	9	2	TC
StvChR_353_a	CAAATCCATTTGAAAGCAAATCAAC	CCATTGATATGTAGTGCCGTAGCA	140–378	6	2	TC
StvChR_356_a	CGAAGTATGCATCTAGCCTTCTC	TCGAGAATTTAAATCAGTCGCTCC	145–216	4	2	TTC
StvChR_357_a	ATAAGGGAACAAATGTGCCAAGAA	GTATAGCAGGTCAATTGATTTCCCG	175–373	8	3	AG
StvChR_360_a	AAAGTGAAGCAAGAGATGGTGC	CTCCTCTGTTTCTCTCCCATCTTTT	105–318	6	2	AG
StvChR_381_a	GCAGGAAGGATATGAGTCATGTTG	AAGACTAATACAGCCCTCAAATTGC	116–139	6	1	AG
StvChR_429_a	TTTTCAATCCATTTTCTTCTCCCC	AAGAGAGATGTCGAGAGAGGAGGG	104–153	6	3	TC
StvChR_430_b	TAGAAGAATCCATACATCACCGCC	TCACACTTATTTCACTGAGCCAACA	159–327	8	3	AAG
StvChR_438_a	AAAATTAAGAACTCAACCATGCC	TACATCCCACTCACTCAGGGAA	103–138	2	2	AG
StvChR_454_a	AAATTGACACCCACAACCAATACC	GTGTTTTGCAATTGTTGATAGCGA	86–263	4	2	ACC
StvChR_480_a	AGAAGTGAAGATCTGAAGATGGCG	CGGAGATGTGTCGAAAGAAGAGA	103–338	11	6	AG
StvChR_488_a	GTCATGTATAGCGCTAAGCGAGTTT	TCCCATGTGACAATACCGATAAGA	99–120	7	5	TG
StvChR_493_a	ATCTCCCTTCTCTAGCGACGTTTT	GATCTAACAAAATCCCACAGCGAC	98–309	7	4	TC
StvChR_534_a	GGAAATGAGTGATTGGAGAGCAAC	TCCTTTCCTGCTAGTGAGATTGTC	110–116	5	2	AT
StvChR_540_b	ACACCCAATCACAATCACAATCAC	AACATAAAGAAGGCGGAATTTGGT	141–171	9	4	AAG
StvChR_544_b	TCTTTCTTTTGGCTCAGGAAGACT	AATTTGTCATGCACAGGTCTCATC	100–172	8	4	AT
StvChR_563_b	TTTTGGGGGTTGCTCTTGATAATA	CTTTTCATTGAATTTGGTTCAAGCC	161–162	7	4	AAG
StvChR_592_a	CCAAAATTGAGCAAAAATCTGGAG	ATGGATGAAGATAGGGATGGTTGA	126–136	7	3	AAG
StvChR_603_a	TGAAGATTATTCACGTTGGTTGGA	CCCCTATTAGCACAGCAAGAAGAA	132–183	12	4	TC
StvChR_632_a	CATGACCGGAAGAAAACAACATGAG	CAGTTTCCAAAATCCTCCATTCATC	147–170	2	1	AG
StvChR_632_b	CCACGAAAAATGCACATATGTTTGA	TTGTTGATAATTGACCAGTGCAGG	106–114	4	2	AT
StvChR_656_a	AAAATGAGGAAAAATCAGGGGAAG	CATCCTTTTCTTCAATCTCCTCAA	126–151	7	2	AGG
StvChR_680_a	CAAATAATTTCGAGTTTCCACGACC	ATGTTTTCTATTGGAGATGCGACTT	108–240	8	2	AG
StvChR_709_a	CATTGTGTGTTTTGCCTTCATA	CATTTCGCCCATTTGAATGTTTAT	174–183	4	2	CT
StvChR_711_a	GCAAAAACAGCCATGATTATTTCA	TAATGCCACCCATTTCTTACCTCTC	156–168	5	2	AG
StvChR_732_a	CGTCTGCATACATCAAGCCAATTA	ATTCTCGATCTCTCCGATCATTT	154–170	3	2	AG
StvChR_777_a	ACTCCAAGAAAAGTCGGGAAGTCT	ATCAGATCGCTGCCTACTAGTGCT	170	1	1	TA
StvChR_793_a	TGGAACCAACATAAATAGATGCACG	GAGCCTGAAACCAAGCCTTAGAC	109–132	8	3	AG
StvChR_807_a	AAATGTACGGGTGTAAATATTGTCA	ATGAAGTTGGATGTTGACGTTCTT	169–173	3	2	AG
StvChR_813_b	TACAAAATCATCCCGAGGAAGAAA	TCTGAATCTTCTAGGCTCTCTCGTTT	113–114	2	1	AG
StvChR_852_a	TCAGTGTGGTGTGCGTATGTATTG	CTCTCTCAAATCCACCTTGACTCG	107–322	5	2	AC
StvChR_879_a	GTCTGAGACCCGATCTGTCTGTTT	TAATTTTCTTCCCTCAATCGCTG	165–170	4	2	TC
StvChR_881_b	AGGTGCTAAAAATGCTGAAAAACAA	CCCAGTAATATTGTGGAGGAGGG	112–198	13	4	AAG
StvChR_917_c	ATTTGGTGTGTCCAAAACGAATC	AAATAGTGCATCCAATGGCTAAA	159–160	2	1	TG
StvChR_925_a	AGTCTTGCAACTGCTCCTTTGTCT	TGTCCATTCTCCAGACTTGAATGA	98–121	9	3	AC
StvChR_926_a	GTTTGACGGTTACGATCAATCCAT	AATCCTGACATCTCATCATCGTCC	182	1	1	ATG
StvChR_945_a	CCAATCTTACGCCAACTTCAAT	TTCCGCTTTTATTCTCTCGCTCT	108–174	9	2	AG
StvChR_961_b	AACCAGATCATCCAACCACCATAG	TGCAAAATTGTGGTGGTGTTTTAG	92–124	8	3	ACC
StvChR_967_a	TGTCATCCTATTTTCATCATTCAGCA	AGGGAACGTGTTATGAAAAGCAGA	139–151	7	2	AG
StvChR_974_a	CGGTGGTTTTTGGTGGTAGTAG	CGGCACATCTCCTCTATCTTCATT	155–391	6	2	TGG
StvChR_989_a	AATCGGACAAGAACGAGATTGAG	AACCAAGCAAAATAGTAAAAGACAGAT	182–405	6	2	AG
StvChR_1003_c	TGACTAGTGGGTGAATGTGGATGT	TGACCAAACTCTCCAAGGAATC	112–297	2	2	AAG
StvChR_1032_a	CCTTATGACTCCATGGAAGATTGC	TCAGCAATCAAGTCCAAGTCTGAA	129–134	2	1	ACT
StvChR_7_a	CGATAGGTCAAGTCGACTCTCTTGC	TCATCTCCTTCTCCTTTCACCTCA	130–152	6	3	GAA
StvChR_9_a	AAATCATAGAAATGGCGGTTGTTG	GATATTCTATCTCCCCTCTTCCCC	125–279	12	4	GA
StvChR_11_b	TCAATCCATATACAGAACGAAAATTGAA	CCTTCCCCAAATCTCTTCTTCATC	169–194	5	2	CT
StvChR_13_b	TTATTAGGATGCATCTCACATCAATTT	ATTCTCGATGATTGTAGCGCAT	167–232	5	2	AG
StvChR_19_a	TCATCATAGAAAACAACACTTGCAACA	ACTAAGACTGGCAAGGTTTGGACC	182–209	3	2	AG
StvChR_21_a	AAAATCACACCCATTTACTGCTCTTC	GGAATCAAATCTTCGACAATAAACACA	103–105	2	1	TA
StvChR_32_a	AACCGCTCGGTATCAGACTAAAT	AAGTTGTTTGGAAACATCAGCGTTT	149–162	4	1	GAT
StvChR_33_a	AGGCGATGCATAAACACACTTTTT	CAAGGAACGAAATCCCGAATACTA	182–183	2	1	AC
StvChR_35_a	GCCGCTACTAATGAGAGAGGAGAA	ACAGTGTGGTGCAAGTGATTGTT	135–157	7	4	AG
StvChR_42_a	CTCTGTTGAGAGTGTGTTGATTCCG	AAACGGTTTATCATTTCAAGCAGGT	102–109	3	2	TC
StvChR_44_a	CCAAAATCTGCTCATCTCGCTTAT	TGTTGACTCGAAGGATGTCATTT	144–176	7	3	CA
StvChR_52_a	GCTTCTTCTGCAGGCGATTACT	ATCATCTCCCTTCTCTTCTTTG	128–172	5	3	AG
StvChR_54_a	GGTGAAGAAATATCCCCGATTAC	TTGATTACCAAAACAAAGAGGCAA	154	1	1	TC
StvChR_57_a	TTTGCTCTCTGTCAACTCTTCCCC	ATCACCAGAATGCCACTTCTTTTC	116–162	6	2	AC
StvChR_62_a	GAGGAATTCGAGTGCAAAGAATA	TGTCAACAGACGCACTCTTTTCT	161–167	4	3	AG
StvChR_66_a	TTGGCATTGCACCCTTAGTTATTT	TATATCTCTCCCCAACACCCCTT	118–340	7	5	GA

(Continued on next page)

Table 2. (Continued) Markers that amplified all the Oleaceae accessions tested detected by Unique Pattern Informative Combination software (Arias et al., 2009).²

Marker ID	Forward primer (5' → 3')	Reverse primer (5' → 3')	Size range	No. all	Max all/S	Motif
StvChR_73_a	CTGGAGTGAGAGATGCAGTTCAAT	TCCCCGAAACAACAGATTATTAAGGA	181–207	7	3	TC
StvChR_74_a	AGGTTTTCTCTCTTTCTAGGGTT	CGGGAATTGAAAGGAGAAAAACATT	134–144	3	1	GAA
StvChR_77_a	AAACAAAGTCACCACGACCACTACT	GCGAAAAGGCTTATGTAGAAGCA	115–194	8	2	ACC
StvChR_80_a	TAAAACTGGGGCTTAAGGGGTAA	GGCTCTGACCGTCAATTGGAT	154–164	3	1	TC
StvChR_89_d	TTGAACTACAGATCCAATTTCCAAGA	TTTTGATCCATCCCTGTATAGCGT	123–131	4	2	AC
StvChR_99_a	ACTGTGTAGCGGTGCTGTAATCA	CTCGCCATTCTCTATCTCTTCTCG	112–154	8	3	AG
StvChR_112_a	AATTCCTGTGTTTTATTTGACGGCG	TGCAGTTGATTAAAGCATGTAGCAAT	192–260	7	2	AAG
StvChR_116_a	AGGAGCGCATAGAGAGGTAGGAAT	ATCAATTCAAAATTCAGGGGCAAA	162–166	3	1	AG
StvChR_120_a	CCGAACCTATTTTCTTCCATCTT	AGAGAAAGGATGCGTTCACTGTTG	107–136	8	5	TC
StvChR_121_a	AGTGGACCATCGTCTAACTCTTCG	TTGCAGACTTTTCTTTGTTTCAACTG	109–110	2	1	ATG
StvChR_131_a	AACGAAGCTTAACCCCTTCGAGG	TCCACGAAATGTTGATGAAATGAG	148–168	7	3	TC
StvChR_141_a	CTTGAGTGGTTGCCCTACTTGACT	CACAGATTCAAGACACGCAGATTTC	173–184	4	2	TC
StvChR_142_a	TGACCATTCCAACACCAAAATATG	GATCATTGCAAAAATCAGGGTTTC	132–146	4	2	AG
StvChR_146_a	TCCACAGATTAATGGCATTGAAAA	GAAGAAGGAGAGGAAAAGGAGAGA	128–131	3	1	TC
StvChR_150_a	TAATTAGTATTGACGGGATTGG	AAAGTTGAGGCAATTAATGATGCTG	269–324	7	2	TTC
StvChR_159_a	TCATTTACTTTTGGTCTTCCATCAC	CCTTGTTCTTGCTGCTTGATAAAT	165–184	6	3	AG
StvChR_169_a	TTTGATGTATATGGATGTGAGCG	AAGAGAGAAACAGTCCAAATCCCC	103–283	4	2	AG
StvChR_184_b	AAGAAGGAAAAATGTAGCGTCGTT	GTATGGGATATGTGTCGGATCCCTC	173–187	5	2	AAG
StvChR_185_b	TCTGAAACCTTAACCCCTCTATCC	CGCCATCAATGTTCCGTTATATT	122–145	5	2	TC
StvChR_195_a	CCCTTCTATTTAGTGGGTTTGGGT	GATGAAAATGACTTCGACGATTCC	155–169	5	3	AG
StvChR_196_a	GCCAAACAGGGTTTTACTTTCCTT	TAAAAATGGTGAGGCAATAACGG	162–164	2	2	TC
StvChR_198_a	TGCAATAGAACAGAAACAAGGCAA	TGTAGGACAGCAGTGAGTAGTCCG	134–161	10	4	TC
StvChR_209_b	AGAGAGAAACAAACACAGGGCAC	GAGAAATCACTGAATAGACAGCGGA	189–212	8	3	TC
StvChR_227_b	CAGCCGTGAACCTTAAAGCTTCTC	CGATCCTCGCGTTTTCTATCTCTA	155–159	2	1	TC
StvChR_246_a	CTGTCTGAATGCTGAATGTATGC	CCAAACCACAACAACCAACAATAC	168–172	4	2	AT
StvChR_253_a	GGAAAGCAAGAAATATTTGGGTGA	AATTATGGAGCCCAAAAAGGAAAA	165–174	3	2	TG
StvChR_254_a	CCCTAAATAACCAATAATGGATTGAAA	TCACAACAAAACCTAGTCGTCAGA	165–219	5	2	AAG
StvChR_267_a	GGCAATCTAATTAATGAAAGGGTT	AACCTAAATATAACTAACGTGCACACAC	149–468	4	2	ATAC
StvChR_301_a	TGGCAATCTGAGACTTTGAGTGAG	CTACTAACCCATCCATGCGTAAA	125–133	7	3	AG
StvChR_304_a	AAGATTGGAGGTTCAATTTTGTG	GAACATCGGTGACGTGGTAACTC	104–289	6	2	AG
StvChR_312_a	TCATTAGAGGTTGTCCAAGCCATT	TTCGAGAGGTCGTTATTTTAGACA	172–221	7	2	AG
StvChR_314_a	GTAGCAAGTGAATCCAAGGAACAA	TGAAATTTTGGAACTAACTAAATCGTCT	135–160	7	3	AC
StvChR_318_a	GTTGTCGACGAACGTCGCTATTA	CATTTTGTCCAGCTCTACCTCCAT	119–458	4	2	AG
StvChR_321_b	TGATGGGGTACGAATTTTGGTAT	ATTTTGGAGATTTTGGGGTGAG	248–485	7	4	TG
StvChR_322_a	TTTTCATCGATTCAATACCCAAATACA	TAAGATATGTGTTGGTGTGGTG	111–214	7	4	ACC
StvChR_326_b	TGACTCGTATGGAAGCAGGAAAAAT	ACCGTATTCAACCACCTCAATCAT	168–188	4	1	TG
StvChR_327_a	GTTAGATGGGGTGAGTGAGCAAAT	TTTCATCTCCATTGGTTCCTTCT	159–345	11	3	TC
StvChR_341_a	GACTTGGACTCTTGTTGGAAGA	GACCTCACCTTTTGCCTCTTTTT	123–181	7	2	AT
StvChR_343_a	TTCTCTCTAATCTCGCCCTTCT	AGAGATGGAGGTCTGCTTCTTCAA	157–576	5	3	TC
StvChR_352_a	TTCTGAGATATTTGAACTTCTCTTGA	AATGCATCATCTCAAAGACATGC	87–187	6	2	TC
StvChR_371_a	GTTGACAGTGAGCGGTGTTCTCT	TTTTCTCTATTTTCTTACCCCG	132–171	8	3	AG
StvChR_379_a	AACATGCACGTCTTACCTGTCT	ATTTTCTCTCCCTCTCGGGTTTC	111–141	5	2	TC
StvChR_411_a	GAAACTTTTCTTGTGAGGGGTGA	TGGAATAATTGAATGGTAGGTGTT	156–191	4	1	TC
StvChR_423_a	TCTGCATAAAGCAACAATGAATCC	TGTGAGGACATGAAAATTTGCTT	175–184	5	2	AG
StvChR_427_a	AGTCTTCTCTTGTGCTCCATTCTG	CTTCCACAGCAGCAAAATCAGTTA	149–153	2	1	TC
StvChR_433_b	CCACCATCTCAATCCCAAAATACA	TATCAACATCTTCAATTGCATCGG	175	1	1	TC
StvChR_437_a	TATAATCGCCATTGAGGGATAGG	TCACTTTGCATAGATCAGGCGTAA	172–176	5	2	AG
StvChR_440_a	CACCATAATTGCGTTTGACCA	GGGTCAACCTCTTTCATCTTTT	147–187	7	2	AC
StvChR_447_a	ATGGTCGAGATCACACAGTTGAGA	CAATTAGTCAGTAGCAGCCCCATC	178–206	6	2	ATG
StvChR_461_a	CCTCCATCCCTCTTTTGTCTTT	TAGAGCTGTGATTGGTGAAATGGA	151–165	6	2	TC
StvChR_468_a	TCATAATTGACTCCATTTTACCCTG	CGAGAAATGATTGAAAGTGCTC	102–118	6	2	TC
StvChR_481_a	AAGATGAATTCACTCATGGGTTGC	AAGAGATCTTTTAAACCCCAACC	159–175	5	2	AT
StvChR_482_a	TGAGTCATTGACAGAAGTTGAGTTG	AGATTGAACCAAAATCAAGCGATT	135–149	4	2	ATC
StvChR_487_a	TGACCATTGATTAGGTGAGTGAA	TTTTGAAACCTCTACCCGATTTTT	166–182	4	1	ATTT
StvChR_490_a	CTTGTTAGGCACAAAGCATTTCAT	TGTTATTACATCTTCAAACCACATTTT	218–224	5	3	AG
StvChR_494_b	CTTGATTGAATTTGGTAGGATCGC	CCATAATTTCTGCACACTTGCTCA	102–127	9	3	ATGT
StvChR_496_a	TTCTATGCCTCTCTCCCTCTCTT	AGGGCGAGTTGGTGTAGACTGTAG	152–362	3	2	TCC
StvChR_504_b	ATCAATCCCAACAATTTCAAACC	AAATGAGTAGGTCGATAGGGGGAG	96–112	8	4	TC
StvChR_507_a	ATACTCCACACCTGGAATGATG	TCTCTATAGGAGGAGGAGATGGGG	149–184	9	4	ATG
StvChR_508_a	TGATGAACAATCTCGTTATGTCCA	CTCACCTCATCACTCAACTCTC	165	1	1	AAG
StvChR_510_a	GCAAGCGTATATTTTCCAACAAAA	CATGTTGAAGAATTATTTTACCCTCTC	99–131	3	1	AG
StvChR_515_a	TGGCTATGGTTATGGTGTGTGTC	ACCACGCAATCTCTCCCTACATC	143–185	3	2	TTG
StvChR_525_a	GGAGAGTCTGGGGTTTATGACAGA	TTTCAATCCGTTTTTAACTCGTCG	120	1	1	TG
StvChR_543_a	GCGGAAGATGAAGCTATGAATGTT	TGTTGGCTCTCATTTTCACTATT	157–165	2	2	AT
StvChR_546_a	TGGCTGGGTTTTCTTGTACCTTA	CACACACCCATAAAATTTTTCGTCA	149–207	7	2	AC
StvChR_566_a	TTTGAGATCTTGGTGTGGACTGTT	GATCCAAACAGTCTTGGAGAAA	180–193	5	2	ATG
StvChR_577_a	CATAATGCATGCACAAAACAGTCA	TTCTGTCTCCACTACTAACATCC	171–177	3	2	AG
StvChR_580_a	CCCCACTAGCATATTACATTCCA	TCCGACCTCAACATTACTCTTCAA	120–142	7	2	AC
StvChR_583_a	AGCAATGAGTGCTTGTAGTGTGGT	TCCAACGTTAATTATTGTCAATTATCTC	182	1	1	AG
StvChR_587_a	CTGCAGCAGAACGGTATACACACT	AACCGTCCCATCCTAAACTTTGT	129	1	1	AG
StvChR_598_a	TGATTTTGCATTTCTTGACATCTG	AAAGAAAGCGAAAGTTCAAGGACC	123–180	3	2	TTC
StvChR_604_a	CTTCTGCTGGGACATTTTCTTC	TGCTCTTGATTCTATCCCAAAAT	148–165	4	3	TTC

(Continued on next page)

Table 2. (Continued) Markers that amplified all the Oleaceae accessions tested detected by Unique Pattern Informative Combination software (Arias et al., 2009).^a

Marker ID	Forward primer (5' → 3')	Reverse primer (5' → 3')	Size range	No. all	Max all/S	Motif
StvChR_605_a	AATTTGAGTGACACCACACAATCG	TTCATCCTTCTTCTTCTCTCCCT	127–541	13	5	TGG
StvChR_613_a	AATGATCTCAGTGATTCGTCCC	ATTTGGTCCGGAACATGATGAGAAT	143	1	1	AT
StvChR_617_a	TACCACCGCTGGTAAATTTCTC	TAAAAGTGTGTTCTGATTCGCCTT	139–143	3	1	ACC
StvChR_622_a	TCCAAAGCTAGAACTCTCACAGGA	GTGGACTCAACACCACCCTTATTC	175–194	4	2	AG
StvChR_627_a	GGGGTGAGGGTGTTATTAATCTGTTG	GCCACTTCCAAATCCCACTATACA	167–182	4	1	AG
StvChR_645_c	TCAACTCAACAGGACCAAACTCTG	CACAGGTGACATTTTGCCCTCTTA	149	1	1	AT
StvChR_647_a	AAGCCGAAAAAGAGAGGAAAAAGA	TTAAGAATGAAGAAATGGAGCCTCA	117–125	3	2	AAGG
StvChR_655_b	ACGTAGCAGGATGAGATTCAGACA	GATTGGGTTCTGGAGTTTGAATG	152	1	1	AC
StvChR_659_a	TTTTATACTTGGAGGAGTGCTAATGC	GACACTATCAATCAGGGTCAAGAAAA	149–283	4	2	AT
StvChR_662_a	ACTTCAGATGTGTCATCTCTCCGA	CTACCTCTCCCACTCCACCTCAT	178–181	4	1	AG
StvChR_671_a	GAATGAAAACCTAACAGGTCAAGT	ACCTGAGCAAAAAGCATGAGAAAA	163	1	1	TC
StvChR_681_a	AATAATTTGTGATCACCACCCAC	TGATGATCATGGAAGTGGTTATGG	152	1	1	ACC
StvChR_686_b	GAAAGTCCGAAAGCCAGGTAATTT	GCGTATCACTGTCAACACTTCAAC	116–126	5	2	TC
StvChR_689_a	TGATGTGACTTTTTGATGGGGCT	TTTTCATTTTGCACTCTCCTTTCC	175–201	6	2	AAG
StvChR_697_a	CACCACCATCTGCAATGAATA	ACCTCGTGGCAATCTATGAATTT	102–103	2	1	ACC
StvChR_704_c	ATTTTCGAAATGTACCGGAACTGA	GTGGAGGAAGACCTCCTAGGGATA	140–240	2	2	TC
StvChR_730_a	TGCCTGTGATTTTGTGTATATCGTG	CCAGTGACTAGTTGAAACTTGAATTGC	131–158	6	2	TC
StvChR_733_a	GATCGGCTCCATTGAAAGAGAGTA	ATGCCGAGGATGATGCTGTAAT	100–104	3	2	AG
StvChR_735_a	CACATGAAAACCTAACCCCTTTGCT	GTCTGAGCAAGGAACCTTACCAAAA	121–139	7	3	TC
StvChR_740_a	GTTCTGTGATGTTTCGGTTTTCTT	AGGGTGACATAATCTCCACAAGGA	103–111	4	3	TG
StvChR_752_a	GACTTTTAGAAGATTATGAGTGCCCC	TTTATTCCTTCTGTTACTTTTCACAAA	172–199	7	3	TC
StvChR_753_a	TCTCGCTTCATCTTATCCTGATCC	TCAATGTGTTATACAGGGAGGGG	124–125	2	1	TC
StvChR_759_b	TAACACGGACACATCTTAAGCAAC	TGTGTCTGTGAAATGGTTGGTTGT	126–143	5	2	ACC
StvChR_786_a	GATTAACAAAATCATCTCTCCCC	CATGCAAAACATGTTCTATGTGCAG	108–118	6	2	TC
StvChR_795_a	TCATCTACCTCTGAACATAATTCCCA	GGCTGTATTGGCTTAGTGAAATGG	140–176	7	3	AACC
StvChR_808_a	GAGCAAAGGTAACAAAGAGCGAA	GTGCTATTTCTCTCGATGTCTGT	163	1	1	AG
StvChR_825_a	TTGGATTAAATTTTGGTTTGTATTGTT	TGGAAGAGAGTTTGTGCAAGGTAG	172–192	5	2	TC
StvChR_832_b	TGTTTTATTGTGCAAAAATCCGGT	GGTGGAGAATGAGGAGAGAAGTGA	158–185	7	3	TC
StvChR_836_c	CACCAGATCAACAAATCCCATACC	GTGTGATAGATGTGGTGGTGCAG	100–218	14	5	ACCATC
StvChR_858_a	TTTACCGTGAGAGATAATGGGAGC	TTAGTTGAGTGACTTGACCACCGA	165–203	6	2	AG
StvChR_862_a	GCTTTTAAGAACTACGTAAGATGGGG	CGATTTGATTCTTAAAGTCACTCG	159–179	3	2	AG
StvChR_865_a	GTCTCTTCAGGTTTACTCGACCCC	TTGACTGTGAAGTCGTGGGCT	136	1	1	ATC
StvChR_894_a	TGAACATTCACTAATCAGCCTCCTC	AGAGAAGAAGACGGACCGGTAGAC	178–358	5	3	TC
StvChR_896_a	GTCCGGATTCACTTCTTCTCTTTC	ATGATAATGGCGATGGTCTCGTAA	98–119	4	2	TTC
StvChR_918_c	AAAATCTCCACATTCACACAGAA	CACGTTCACTGTTTCGTTCACT	173–184	3	2	AC
StvChR_927_a	TAAAATCCGACTTTTCCCTCACAA	CACCAACTTTGGCTAAGAAAAATGA	172	1	1	TC
StvChR_935_a	TAAGCCTTGAAGTAGTAGCTGCC	ATCGAACTTAAAGTGTGCTGCAATC	83–189	5	2	TC
StvChR_943_a	TTACTCCAATCGCTTTCCTAAACG	CAGCCTACATCCAACTTTGTTC	123–146	2	2	AG
StvChR_950_a	TATTACCCTATTCCCTGACGAGCA	GGGAAGGAGAAAAACAAAGAGGAA	155–181	5	2	TG
StvChR_976_a	TACGTAGGGAGGAATGTTCAAACG	GCAAGGAACATCAAAAGTCCATCT	179	1	1	AAG
StvChR_996_a	ATTTGAGGGGAATAGGGAATCGT	AAGTCGAAGACTCGAAGTCCAC	99–134	5	3	TG
StvChR_1006_a	TGCTGACCACGTTGTATCTGTTT	ACTGACGTGGAACCAAGAAAGAAG	97–113	3	3	TC
StvChR_1016_a	GCTTTCTCTTTTGTTCCTCTTTC	TTTATCCAAAGATTCGAGTGTAATA	155–183	5	2	TTC
StvChR_1017_a	GACCAACCAAAACAACTCAACCTC	CTCCGGTGGTAAATTTGAAAGAGAG	92–127	15	5	TC
StvChR_1025_a	AGCCCATTTTGCATTTGAGTAT	ACAATGATTTTGTGTTTGTATTGTCT	157–182	4	1	AC
StvChR_1031_b	TTTGAATCCATCCATTTGAGAACA	TGGGAGAAAAATGAGACTTGGTTT	139–146	8	4	AC
StvChR_1061_a	CACAATTGCACCAATTCCTACAT	CCATCTCTTAAACCTCATTTC	150	1	1	AT
StvChR_1072_a	TACCAGATGTGGGAACAACACAGA	TTTATGAGGGGTGAAATTTGATGA	146–161	5	4	AAG
StvChR_217_sk/a	ACTTGAAGATTTATCGGCACATCC	GATGGACCTTTCCATGTTCTTGAC	147–169	5	3	TC
StvChR_351_sk/a	AGTTTCTCTCTTACTACGCCACT	AATGAAGGCGACAATCCACTAAA	154–184	6	3	ACTC
StvChR_432_sk/a	GTCTCCGACGATGAACCAAACTAT	TTTCAACCTTTGTTCTCGTCTCC	165–176	2	2	AG
StvChR_568_sk/a	TCCCAAAATACCCAAAGAAAACAA	TTGGAATCCTTTTCTCTATTCCC	149–150	2	1	AAGAT
StvChR_710_sk/a	GTAGAAGAAGGGGAAATGGGAAGA	CTATCCTGGGTGGGTATGAAATTG	174–191	5	2	AAGAG
StvChR_745_sk/a	CACACTCCATCGCAACAATAGAAG	TGTGTGGTTGATTAGTTCGTGGTT	99–157	8	3	AACC
StvChR_955_sk/a	ATTTCGCGCCCATACACATACAC	CGACCGAGTCTCCATAGAGATTT	172–217	7	2	AC

^aMarkers that amplified all the Oleaceae (StvChR_15_a to StvChR_1032_a) and *Chionanthus retusus* only (StvChR_7_a to StvChR_955_sk/a). DNA sequences shown in bold had significant hits on BLASTx. In the columns, size range is: observed on the samples, includes the 23 bp of primer tailing. “No. all” = total number of alleles detected. “Max all/S” = maximum number of alleles detected on an individual sample. Sequences corresponding to the contigs were submitted to GenBank with accession numbers GQ117288 to GQ118.

for four species of Oleaceae (*C. pygmaeus*, *C. retusus*, *C. virginicus*, and *O. americanus*) using 57 markers as shown in Figure 2A. We also calculated genetic similarity coefficients for *C. retusus* using 195 markers as shown in Figure 2B. We observed a high level of polymorphism within the species *C. retusus* using these 195 markers. These markers allowed a clear distinction among *C. retusus* varieties as is shown by the high bootstrap resampling coefficients obtained. Bootstrap confidence values are indicated on the dendrograms. Clusters formed groups by species with

the exception of *C. virginicus* (6007), which associated with *C. pygmaeus* based on the 57 markers analyzed as shown in Figure 2A.

Discussion

Chionanthus SSR-enriched libraries were generated using 24 oligo repeats; however, we detected twice as many repeat motifs, probably as a result of non-specific hybridization during the enrichment process. Although we detected a large number of repeats in the assembled contigs, only 394 pairs of

primers were designed under the stringent conditions used. In *C. retusus*, the frequency of repeats decreased with the increase in length of the repeat motifs (Fig. 1). This trend has also been observed in the distribution of SSRs of other eukaryotic genomes (Katti et al., 2001). We also found that in *C. retusus*, the most abundant di-, tri, and tetra-nucleotide repeat motifs were AG/AC, AAG/ACC, and ACTC/ATAC, respectively. In *Arabidopsis thaliana* (L.) Heynh., the same di- and trinucleotide motifs were the most abundant across the entire genome;

however, the predominant tetranucleotide motifs were of the type “AAA/(CTG)” (Anwar and Khan, 2005). Curiously, we found at least one (CG)_n repeat in *C. retusus*, a particular repeat motif that has not been found in the entire *A. thaliana* genome (Anwar and Khan, 2005) and that is very rare in other eukaryotic genomes (Katti et al., 2001).

Ninety-six percent of 57 SSRs that amplified across the four species of Oleaceae (*C. pygmaeus*, *C. retusus*, *C. virginicus*, and *O. americanus*) were polymorphic. Within the species *C. retusus*, 61% of 195 SSRs we developed were polymorphic. The high level of polymorphism is not unexpected in this study, because we chose representative samples of *C. retusus* that covered two extremes of its range in Asia with *C. retusus* CR1 and CR3 originating from the Yunnan province in China and CR4 and CR6 from Japan. *Chionanthus retusus* CR2 represents the typical

form in cultivation in the United States and is likely derived from the same parental stock as CR5, a plant we originally received as *C. retusus* var. *serrulatus* (Hayata) Koidz. from the Arnold Arboretum, but is of uncertain provenance. The varietal epithet *serrulatus* was first applied to a new species of *Chionanthus* from Taiwan, which was later reduced to a variety of *C. retusus* (Fogg, 1960). A larger survey of North American nurseries is planned to determine the level of genetic diversity of *C. retusus* in cultivation. Only a handful of *C. virginicus* and *C. pygmaeus* accessions were available for sampling and used to test cross-amplification of *C. retusus*-derived SSRs. A broader sampling, encompassing a greater range of *C. virginicus* in eastern North America and larger sampling in Florida, where it is in close proximity to *C. pygmaeus*, is needed to determine population structure and similarities between these species.

The percentage of heterozygous loci found in *O. americanus* (68.8%) was twofold higher than in *C. retusus* (30.1% to 39.8%). In addition, *O. americanus* shared 29% of detected alleles with *C. retusus*, which had a genetic similarity of 0.54. Very few alleles were shared between *O. americanus* and the species *C. virginicus* and *C. pygmaeus*. The amplification and sharing of alleles between *Chionanthus* sp. and *O. americanus* is indicative of a recent divergence and supports the conclusions of Wallander and Albert (2000) who reported *O. americanus* to be closely related to *Chionanthus*. The greater number of polymorphic loci in *O. americanus* may also be attributed to genome size. Whereas *Chionanthus* species used in this study are presumed diploid with $2n = 46$, *Osmanthus americanus* is a hexaploid with $2n = 138$ (Taylor, 1945).

BLASTx screening of DNA sequences containing microsatellites showed significant hits on some interesting genes. For example, the sequence that originated marker StvCHR_508a had similarity to SORBIDRAFT of *Sorghum bicolor* that is related to the development of inflorescences (Polegri et al., 2010). The DNA sequence corresponding to marker StvCHR_680a had similarity to proteasome subunit alpha; both 26S proteasome and the *REV* (*REVOLUTA*) genes are required for the maintenance of root apical meristem and shoot apical meristem (Zhang et al., 2010), and it has been shown that plants with double mutations on both isoforms of the 26S proteasome are often non-viable (Gallois et al., 2009). Several DNA sequences had similarity to chromatin remodeling, DNA binding, and control of transcription such as those from markers StvCHR_356a, StvCHR_430b, StvCHR_777a, and StvCHR_943a. The DNA sequences associated to marker StvCHR_709a had similarity to the brahma-related gene (*BRG1*), which is implicated in tumor suppressor function (Hendricks et al., 2004). Further research taking in

Fig. 1. Motifs and frequency of repeats detected in *Chionanthus retusus* simple sequence repeat (SSR)-enriched libraries. Frequency of motif repeats detected in an SSR-enriched library after screening 1072 contigs. Another 43 repeat motifs with frequencies lower than four were not included in the plot. Embedded graph shows the number of repeats detected in *C. retusus* arranged by repeat motif length (BP).

Table 3. Markers that amplified DNA samples from four species of Oleaceae (*Chionanthus retusus*, *C. virginicus*, *C. pygmaeus*, and *Osmanthus americanus*) selected by their performance in terms of peak quality and distribution across species.

Marker	UPIC	Marker	UPIC	Marker	UPIC	Marker	UPIC	Marker	UPIC
StvChR_945a	12	StvChR_209b	6	StvChR_793a	5	StvChR_656a	4	StvChR_447a	4
StvChR_350a	10	StvChR_120a	6	StvChR_343a	5	StvChR_740a	4	StvChR_430b	4
StvChR_57a	10	StvChR_504b	6	StvChR_321b	5	StvChR_832b	4	StvChR_124a	4
StvChR_35a	10	StvChR_494b	6	StvChR_185b	5	StvChR_896a	4	StvChR_195a	4
StvChR_9a	8	StvChR_881b	6	StvChR_689a	5	StvChR_351sk	4	StvChR_1017a	4
StvChR_540b	8	StvChR_301a	6	StvChR_76a	5	StvChR_141a	4	StvChR_437a	4
StvChR_925a	8	StvChR_131a	6	StvChR_352a	5	StvChR_99a	4	StvChR_312a	4
StvChR_112a	8	StvChR_221a	6	StvChR_546a	4	StvChR_786a	4	StvChR_44a	4
StvChR_11b	8	StvChR_955sk	6	StvChR_440a	4	StvChR_825a	4	StvChR_989a	3
StvChR_66a	8	StvChR_198a	6	StvChR_184b	4	StvChR_314a	4	StvChR_304a	3
StvChR_62a	8	StvChR_490a	6	StvChR_468a	4	StvChR_745sk	4	StvChR_411a	3
StvChR_251b	7	StvChR_935a	6	StvChR_961b	4	StvChR_894a	4	StvChR_480a	3
StvChR_603a	7	StvChR_150a	6	StvChR_73a	4	StvChR_680a	4	StvChR_563b	3
StvChR_566a	7	StvChR_461a	6	StvChR_284a	4	StvChR_254a	4	StvChR_1025a	3
StvChR_316a	7	StvChR_580a	6	StvChR_94a	4	StvChR_1006a	4	StvChR_33b	3
StvChR_19a	7	StvChR_605a	6	StvChR_1016a	4	StvChR_481a	4	StvChR_598a	3
StvChR_795a	6	StvChR_13b	6	StvChR_967a	4	StvChR_327a	4	StvChR_711a	3
StvChR_836c	6	StvChR_371a	6	StvChR_507a	4	StvChR_159a	4	StvChR_77a	3
StvChR_710sk	6	StvChR_169a	5	StvChR_752a	4	StvChR_15a	4	StvChR_544b	3
StvChR_341a	6	StvChR_357a	5	StvChR_730a	4	StvChR_950a	4	StvChR_326b	3

UPIC values correspond to the number of DNA samples that were discriminated by the marker out of the 12 lines tested from Table 1. Combinations of these markers can detect a number of unique patterns or alleles equal to the sum of their UPIC values.

UPIC = Unique Pattern Informative Combination.

Fig. 2. Clusters calculated using the unweighted paired group method using arithmetic averages (UPGMA) of NTSYSpc 2.2. Confidence levels from bootstrap analysis (5000 replicates) are indicated at the nodes. (A) Cluster analysis of 12 DNA samples of *Chionanthus* and *Osmanthus* using 57 simple sequence repeat (SSR) markers that amplified across species; (B) cluster analysis of 6 DNA samples of *C. retusus* using 195 SSR markers that amplified this species.

consideration the gene functions potentially affected by the polymorphism of the markers presented here would provide useful insight for plant breeding programs.

Considering that transferability of molecular markers is feasible within plant families (Barreneche et al., 2004; White and Powell, 1997; Yasodha et al., 2005), the markers we developed for *Chionanthus* could be used to examine other economically important members of the Oleaceae such as *Fraxinus* and *Olea*. Although a number of microsatellites are available for *Olea* (Cipriani et al., 2002; Omrani-Sabbaghi et al., 2007; Stambuk et al., 2007), between 200 and 500 additional markers are necessary for 80% to 95% of the *Olea* genome to lie within 10 cM of a marker (Wu et al., 2004).

In the present study, we have characterized a large number of SSR markers for *Chionanthus*-related species: 55 that discriminate among the four species of Oleaceae tested, 162 that discriminate within the species *C. retusus*, 35 polymorphic ones for *C. virginicus*, and 18 polymorphic markers for *C. pygmaeus*. These markers could aid in identifying genetic diversity of *Chionanthus* germplasm and allow verification of hybrids, pedigrees, and cultivars for botanical characterization and ornamental tree breeding programs.

Literature Cited

Abajian, C. 1994. Sputnik. Aug. 2010. <<http://espressoftware.com/sputnik/index.html>>.
Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.
Anwar, T. and A.U. Khan. 2005. Mapping and analysis of simple sequence repeats in the *Arabidopsis thaliana* genome. *Bioinformation* 1:64–68.

Arias, R.S., L.E. Ballard, and B.E. Scheffler. 2009. UPIC: Perl scripts to determine the number of SSR markers to run. *Bioinformation* 3:353–361.
Barreneche, T., M. Casasoli, K. Russell, A. Akkac, H. Meddour, C. Plomion, F. Villani, and A. Kremer. 2004. Comparative mapping between *Quercus* and *Castanea* using simple-sequence repeats (SSRs). *Theor. Appl. Genet.* 108:558–566.
Brownstein, M.J., J.D. Carpten, and J.R. Smith. 1996. Modulation of non-templated nucleotide addition by Taq DNA polymerase: Primer modifications that facilitate genotyping. *Biotechniques* 20:1004–1006, 1008–1010.
Chang, M.-C., L.C. Chiu, Z. Wei, and P.S. Green. 1996. Oleaceae: *Chionanthus*, p. 293–295. In: Wu, Z. and P.H. Raven (eds.), *Flora of China*. Vol. 15. Science Press, Beijing, China.
Cipriani, G., M.T. Marrazzo, R. Marconi, A. Cimato, and R. Testolin. 2002. Microsatellite markers isolated in olive (*Olea europaea* L.) are suitable for individual fingerprinting and reveal polymorphism within ancient cultivars. *Theor. Appl. Genet.* 104:223–228.
Dirr, M.A. 1998. *Manual of woody landscape plants*. Stipes Publishing, Champaign, IL.
Efron, B., E. Halloran, and S. Holmes. 1996. Bootstrap confidence levels for phylogenetic trees. *Proc. Natl. Acad. Sci. USA* 93:13429–13434.
Felsenstein, J. 1985. Confidence-limits on phylogenies—An approach using the bootstrap. *Evolution* 39:783–791.
Fogg, J.M.J. 1960. *Chionanthus* in the Philadelphia area. *Morris Arboretum Bul.* 11:3–6.
Gallois, J.L., A. Guyon-Debast, A. Lecureuil, D. Vezon, V. Carpentier, S. Bonhomme, and P. Guerche. 2009. The arabidopsis proteasome RPT5 subunits are essential for gametophyte development and show accession-dependent redundancy. *Plant Cell* 21:442–459.
Green, P.S. 2004. Oleaceae, p. 296–306. In: Kubitzki, K. (ed.), *The families and genera of vascular plants*. 7th Ed. Springer-Verlag, Berlin, Germany.

Hendricks, K.B., F. Shanahanand, and E. Lees. 2004. Role for BRG1 in cell cycle control and tumor suppression. *Mol. Cell. Biol.* 24:362–376.
Katti, M.V., P.K. Ranjekar, and V.S. Gupta. 2001. Differential distribution of simple sequence repeats in eukaryotic genome sequences. *Mol. Biol. Evol.* 18:1161–1167.
Nicholson, R.G. 1990. The fringe tree and its far-flung cousins. *Arnoldia* 50:24–31.
Omrani-Sabbaghi, A., M. Shahriari, M. Falahati-Anbaran, S.A. Mohammadib, A. Nankalic, M. Mardia, and B. Ghareyazie. 2007. Microsatellite markers based assessment of genetic diversity in Iranian olive (*Olea europaea* L.) collections. *Sci. Hort.* 112:439–447.
Polegri, L., O. Calderini, S. Arcioni, and F. Pupilli. 2010. Specific expression of apomixis-linked alleles revealed by comparative transcriptomic analysis of sexual and apomictic *Paspalum simplex* Morong flowers. *J. Expt. Bot.* 61: 1869–1883.
Rozen, S. and H. Skaletsky. 2000. Bioinformatics methods and protocols. In: *Methods in molecular biology* Totowa, NJ.
Sharopova, N., M.D. McMullen, L. Schultz, et al. 2002. Development and mapping of SSR markers for maize. *Plant Mol. Biol.* 48:463–481.
Stambuk, S., D. Sutlovic, P. Bakaric, S. Petricevic, and S. Andelinovic. 2007. Forensic botany: Potential usefulness of microsatellite-based genotyping of Croatian olive (*Olea europaea* L.) in forensic casework. *Croat. Med. J.* 48: 556–562.
Taylor, H. 1945. Cyto-taxonomy and phylogeny of the Oleaceae. *Brittonia* 5:337–367.
Teichen, N., R.S. Arias, Z. Pan, I. Khan, and B.E. Scheffler. 2010. Optimized construction of microsatellite-enriched libraries. *Mol. Ecol. Resources* 10:508–515.
Ueda, K. 1996. Androdioecism in *Chionanthus retusus* (Oleaceae). *J. Phytotax. Taxon.* 44:91–92.
Waldbieser, G.C., S.M. Quiniou, and A. Karsi. 2003. Rapid development of gene-tagged microsatellite markers from bacterial artificial chromosome clones using anchored TAA repeat primers. *Biotechniques* 35:976–979.
Wallander, E. and V.A. Albert. 2000. Phylogeny and classification of Oleaceae based on rps16 and trnL-F sequence data. *Amer. J. Bot.* 87: 1827–1841.
White, G. and W. Powell. 1997. Cross-species amplification of SSR loci in the Meliaceae family. *Mol. Ecol.* 6:1195–1197.
Wu, S.B., G. Collins, and M. Sedgley. 2004. A molecular linkage map of olive (*Olea europaea* L.) based on RAPD, microsatellite, and SCAR markers. *Genome* 47:26–35.
Yap, I.V. and R.J. Nelson. 1996. Winboot: A program for performing bootstrap analysis of binary data to determine the confidence limits of UPGMA based dendrograms. IRRI Discussion Paper Series No. 14. International Rice Research Institute, Manila. May 2009. <<http://www.irri.org/science/software>>.
Yasodha, R., M. Ghosh, R. Sumathi, and K. Gurumurthi. 2005. Cross-species amplification of eucalyptus SSR markers in Casuarinaceae. *Acta Botanica Croatica* 64:115–120.
Zhang, Z., H. Wang, D. Luo, M. Zeng, H. Huang, and X. Cui. 2010. Convergence of the 26S proteasome and the REVOLUTA pathways in regulating inflorescence and floral meristem functions in *Arabidopsis*. *J. Expt. Bot.* doi: 10. 1093/jxb/erq277.