

‘Ray Ruby’ Grapefruit Affected by Huanglongbing II. Planting Density, Soil, and Foliar Nutrient Management

Dinesh Phuyal

University of Florida, Institute of Food and Agricultural Sciences, Indian River Research and Education Center, 2199 South Rock Road, Fort Pierce, FL 34945

Thiago Assis Rodrigues Nogueira

São Paulo State University, School of Engineering, Department of Plant Protection, Rural Engineering and Soils, 56 Brasil Sul Avenue, Ilha Solteira, SP 15385-000, Brazil

Arun D. Jani

University of Florida, Institute of Food and Agricultural Sciences, Indian River Research and Education Center, 2199 South Rock Road, Fort Pierce, FL 34945

Davie M. Kadyampakeni

University of Florida, Institute of Food and Agricultural Sciences, Citrus Research and Education Center, 700 Experiment Station Road, Lake Alfred, FL 33850

Kelly T. Morgan

University of Florida, Institute of Food and Agricultural Sciences, Southwest Florida Research and Education Center, 2685 State Road 29 North, Immokalee, FL 34142

Rhuanito Soranz Ferrarezi

University of Florida, Institute of Food and Agricultural Sciences, Indian River Research and Education Center, 2199 South Rock Road, Fort Pierce, FL 34945

Additional index words. citrus greening, foliar application, fruit quality, fruit yield, high-density, micronutrients

Abstract. Since the arrival of Huanglongbing (HLB) disease in Florida, several management approaches, including modification of orchard architecture design and nutritional therapy, have been explored. High-density plantings anticipate early economic returns from HLB-affected orchards. With no cure available for HLB, balanced nutrient application through soil and foliar spraying can mitigate the disease. A 2-year study was conducted to investigate the effects of three grapefruit (*Citrus paradisi*) planting densities [single-row (300 and 440 trees per ha), and double-row high-density (975 trees per ha)], two controlled-release fertilizer (CRF) blends, and foliar-applied micronutrients (FAM) (a blend of B, Mn, and Zn at 0, 1.5, 3, and 6 times the recommended rates) on grapefruit growth and fruit yield, physiological parameters, and foliar nutrient concentrations in an HLB-affected orchard. All the trees tested positive for HLB based on real-time quantitative polymerase chain reaction (qPCR) test. The highest planting density resulted in the lowest trunk diameter and canopy volume. Despite lower yield per tree in 2019–20, 975 trees per ha planting induced the greatest fruit and solid yields per ha. Also, the fruit produced from 975 trees per ha planting tended to be acidic with the deposition of more soluble solids. Use of CRF with higher micronutrients increased canopy volume with the expense of reduced fruit number in 2019–20. FAM did not affect cycle threshold (Ct) value and tree growth parameters. Fruit yield, photosynthesis rate, and stomatal conductance (g_s) decreased, and all leaf nutrient concentrations except B increased in 2019–20 with all FAM rates tested. In conclusion, our study showed that high-density planting optimizes yield under HLB-endemic conditions. In addition, supplemental soil and foliar micronutrient application do not enhance yield of HLB-affected trees over a 2-year timeframe, warranting further research for confirmation of results.

Growers have been using advanced horticultural practices and modifying orchard architecture design to cope with the Huanglongbing (HLB) epidemic in Florida. High-

density planting is one modification that can potentially anticipate early economic returns (Roka et al., 2009; Singerman et al., 2018; Stover et al., 2008), which is important

because of the high probability of HLB-induced fruit yield reductions in later years. Previous work has shown that high-density plantings can increase fruit yield in mandarin (*Citrus nobilis* × *Citrus deliciosa*), sweet orange (*Citrus sinensis*), lime (*Citrus aurantifolia*), and grapefruit (*C. paradisi*) (Dalal et al., 2013; Ladaniya et al., 2020; Moreira et al., 2019; Phuyal et al., 2020; Singerman et al., 2018). The past studies were limited to tree density only, and scarce literature is available about the effect of high-density plantings with conjunction of soil and foliar nutrient application on grapefruit tree health, fruit yield, and fruit quality.

Citrus tree physiological function is negatively impacted by HLB (Etxeberria et al., 2009; Gonzalez et al., 2012). Nwugo et al. (2013) found that HLB-affected grapefruit showed alteration of several foliar proteins due to nutrient deficiency. Conversely, nutrient management can improve physiological function in HLB-affected citrus. Li et al. (2014) showed that a higher application rate of zinc (Zn) can increase photosynthetic rates and g_s of HLB-affected grapefruit seedlings under hydroponic conditions. In addition, Yang et al. (2012) observed reduced carbon dioxide (CO₂) assimilation in magnesium (Mg)-deficient pomelo (*Citrus grandis*), possibly due to the decrease in photosynthetic electron transport capacity.

As an essential constituent of chlorophyll, Mg is involved in photosynthesis, enzyme activation, protein synthesis, carbohydrate partitioning, and plant growth, playing an important role in resistance mechanisms against bacterial and fungal diseases (Huber and Jones, 2013; Marschner, 2012). Magnesium deficiency in citrus results in the impairment of photosynthesis in leaves, alterations of gas exchange, and metabolite imbalance in the tree (Li et al., 2017; Tang et al., 2012). Application of Mg in Mg-deficient soil can improve the quality of horticultural crop products, especially when the quality traits are dependent on photosynthesis and assimilate translocation processes regulated by Mg within the plant system (Gerendás and Führs, 2013).

Foliar nutrient application is important during citrus reproductive stages to enhance fruit yield and quality. When soil properties are not conducive to nutrient uptake (temperature extremes, low or excess soil moisture, pH, and salinity), foliar applications can meet nutrient demand. Compared with soil application, foliar nutrient application can potentially reduce nutrient runoff, thereby reducing negative environmental impacts (Lovatt, 1999, 2013). FAM is especially important for citrus trees grown in calcareous soils, because they typically show micronutrient deficiency symptoms, namely for iron (Fe) and Zn. High soil pH limits the uptake of micronutrients, and foliar application accelerates the process of nutrient acquisition to correct deficiencies (Ibrahim et al., 2007).

Several studies have demonstrated the benefits of FAM in citrus orchards. Labanauskas (1963) reported that manganese (Mn) and Zn deficiency in grapefruit can be

corrected by foliar spray of those nutrients. Morgan and Kadyampakeni (2020) suggested Zn deficiency should be corrected by only foliar spray on calcareous soils as the alkaline pH halts soil available Zn. Boaretto et al. (1997) showed that foliar boron (B) application increased leaf B concentration of sweet orange to an adequate range without affecting fruit yield or quality. Similar findings with Zn were reported by Boaretto et al. (2002). A study by Morgan et al. (2016) found that spraying of B, Mn, and Zn increased the leaf concentrations of those micronutrients in 6-year-old 'Valencia' sweet orange trees on Swingle citrumelo rootstock (*C. paradisi* × *Poncirus trifoliata*).

Foliar application alone or in combination with other substances such as hormones and insecticides has been shown to increase citrus fruit yield. Several authors noted higher fruit yield of 'Valencia' sweet orange with the application of insecticides and foliar nutrients (Stansly et al., 2014; Tansey et al., 2017). Morgan et al. (2016) reported that foliar application containing 3× the recommended rate of Mn increased fruit yield in 6-year-old 'Valencia' sweet orange in a five-year study. Rouse et al. (2017) observed that increasing foliar nutrient application rates enhanced 'Valencia' sweet orange fruit yield, while Al-Obeed et al. (2018) showed that foliar application of B and Zn increased mandarin (*Citrus reticulata*) fruit yield and quality.

Foliar nutrient application can also mitigate disease expression. Pustaka et al. (2008) observed reduced disease symptoms on mandarin trees affected with HLB with foliar nutrient application. Shen et al. (2013) observed increased disease resistance, higher Ct value, and improved canopy size in 'Valencia' orange with long-term application of foliar nutrients along with salicylic acid and phosphite.

Balanced application of nutrients to correct deficiencies is an important tool to control biotic and abiotic stresses as well as to improve nutrient use efficiency (Zekri and




Fig. 1. Monthly accumulated rainfall precipitation, mean, minimum, and maximum temperatures recorded in 2018–19 and 2019–20.

Obreza, 2018, 2019). The efficiency of enhanced nutritional programs on HLB-affected trees is site- and variety-specific with some studies showing no effect of those substances for mitigating disease, increasing fruit yield, and improving fruit quality (Gottwald et al., 2012). There is limited information available about high plant density and the joint use of soil and foliar nutrients on HLB-affected grapefruit. Nevertheless, it is hypothesized that nutrients, especially Mg and micronutrients, applied to grapefruit affected by HLB can increase fruit yield and reduce disease incidence by improving physiological parameters.

The objective of this study was to evaluate the effects of tree planting density, soil Mg and micronutrient application, and FAM on grapefruit growth and fruit yield parameters, physiological response, and HLB disease.

Materials and Methods

Study site. A large-scale field trial was conducted from 2018 to 2020 at the University of Florida Institute of Food and Agricultural Sciences (UF/IFAS) Indian River Research and Education Center research farm in Fort Pierce, FL (lat. 27°26'08.2" N, long. 80°26'43.2" W, and elevation of 8 m). Weather data were collected by using the Florida Automated Weather Network (FAWN) station in St. Lucie West (Fig. 1). Soil moisture was monitored during the study using soil moisture sensors (TDT ACC-SENSDI; Acclima Inc., Meridian, ID) placed

0.6 m away from the main trunk and 15 cm below the soil surface, where maximum root activity was observed. Sensors were connected to a datalogger (CR 205; Campbell Scientific Inc., Logan, UT) programmed to collect data at 15-min intervals. The datalogger was connected to a rechargeable battery (6FM7; Toyo, Tonawanda, NY) and solar panel (HY015-12P; Acopower, Walnut, CA) mounted on a pole. The data were averaged according to the treatments (Fig. 2).

Plant material. Five-year-old 'Ray Ruby' grapefruit trees on Kuharske citrange (*C. sinensis* × *P. trifoliata*) rootstock planted in Sept. 2013 were used as planting material. The study was conducted for two seasons (2018–19 and 2019–20).

Experimental design and treatments. The experiment was conducted using a split-split-plot design with four replications. We tested three planting densities, two CRF blends, and four FAM rates. Planting densities were as follows: 1) 300 trees per ha, single-row low-density (SR/LD); 2) 440 trees per ha, single-row high-density (SR/HD); and 3) 975 trees per ha, double-row high-density (DR/HD).

In 2018–19, we tested two CRF blends [16 nitrogen (N)–1.31 phosphorus (P)–16.6 potassium (K) and 12N–1.31P–7.47K], while in 2019–20, both CRF blends were adjusted according to University of Florida/Institute of Food and Agricultural Science (UF/IFAS) recommendation, and made [12–3–14 (12N–1.31P–11.62 K) with 70% of N as a CRF and other micronutrients as sulfates; and 12–3–14

Received for publication 24 June 2020. Accepted for publication 25 June 2020.

Published online 31 July 2020.

This research was funded by the UF/IFAS Citrus Initiative and in part by the U.S. Department of Agriculture (USDA)-APHIS-MAC Agreement No. AP19PPQSandT00C116.

We thank Dr. Mark Ritenour, Dr. Jude Grosser, Tom James, Randy Burton, Hernan Soto, Cristina Gil, Andres Gonzalez, Clarence King, Donald Davis, Taylor Meadows, Megan Eckman, Benjamin Holly, and Peter Spyke for the treatment application, research field maintenance, and data collection. We also thank Southern Gardens Diagnostic Laboratory (Mike Irely) and Citrus Research and Development Foundation for the real-time quantitative polymerase chain reaction (qPCR) testing.

R.S.F. is the corresponding author. E-mail: rferrarezi@ufl.edu.

This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 2. Volumetric water content reading during the experiment. Two different controlled-release fertilizer (CRF) blends [12–3–9 (12N–1.31P–7.47K) with all micronutrients as sulfur-coated products at higher rates; and 16–3–20 (16N–1.31P–16.6K) with all micronutrients as sulfates] in 2018–19 and two CRF blends [12–3–14 (12N–1.31P–11.62 K) with 2 times of Mg and 2.5 times of recommended rates; and 12–3–14 (12N–1.31P–11.62 K) with the recommended rate for Mg and all micronutrients] in 2019–20 (A), three planting density [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)] (B), and four foliar treatments (0, 1.5, 3, and 6 times) of recommended rates of B, Mn, and Zn (C). Each point represents mean \pm SE.

Table 1. Nutrient application rates for controlled-release fertilizer (CRF) blends used in the study.

	CRF 1 amount (kg·ha ⁻¹)		CRF 2 amount (kg·ha ⁻¹)	
	2018–19	2019–20	2018–19	2019–20
N	180	180	180	180
P ₂ O ₅	45	44.82	34	44.82
K ₂ O	135	209.17	225	209.17
Ca	30	14.94	11	14.94
Mg	35	36.07	16	18.08
S	223	227.83	73	194.34
B	1.09	1.69	10.67	0.67
Cu	0.60	1.49	0	0.60
Fe	15.58	14.72	5.88	5.89
Mn	18.90	21.04	17.82	8.41
Mo	0.09	0.22	0	0
Zn	7.32	14.72	5.88	5.89

(12N–1.31P–11.62 K) with 2 times Mg and 2.5 times the recommended rates of micronutrients with 70% of N as a CRF and other micronutrients as sulfates] (Table 1). Both seasons, CRF blends were applied in February, July, and October, with the amount based on recommended N rates for mature grapefruit trees in Florida (Morgan and Kadyampakeni, 2020).

Four micronutrient rates (a blend of B, Mn, and Zn) were used: 0, 1.5, 3, and 6 times the recommended UF/IFAS rates (B = 0.28, Mn = 4.48, and Zn = 5.60 kg·ha⁻¹) (Morgan and Kadyampakeni, 2020). Water-soluble fertilizers {[manganese sulfate monohydrate (MnSO₄·H₂O) with 32% of Mn], zinc sulfate monohydrate [(ZnSO₄·H₂O) with 36% of Zn], and disodium octaborate tetrahydrate

[(Na₂B₈O₁₃·4H₂O) with 21% of B]} were sprayed in March, May, and September during growth flushes as suggested by Morgan et al. (2016).

Field preparation and orchard layout.

The orchard architecture design and cultural practices followed were the same as indicated by Phuyal et al. (2020) because the studies were conducted in sequence in the same area. There were 96 experimental units each measuring 15.24 × 29.12 m.

Concentration of *Candidatus Liberibacter asiaticus* DNA in plant leaf tissue. Eight random trees were chosen excluding border trees to collect leaf samples. The samples were collected in April and analyzed by real-time qPCR. The processes to collect and analyze leaf samples are detailed in Phuyal et al. (2020). Ct value was assessed in which less than 32 is considered as and HLB-positive tree.

Tree growth measurement. Four trees were selected randomly in each plot excluding the border trees to measure the tree height, canopy diameter and trunk diameter following Phuyal et al. (2020). Canopy volume was calculated by using geometric prolate spheroid equation: [(4/3) (π) (tree height/2) (average canopy diameter)²] (Obreza and Rouse, 1993).

Fruit yield and fruit size. We selected four random trees from each plot to measure fruit yield (excluding the border trees). Fruit yield per tree, fruit yield per unit area, and fruit diameter were determined by using an optical sizer (Autoline, Reedley, CA) following Phuyal et al. (2020). Fruit size was categorized based on commercial categories of grapefruit (number of fruit per carton) and grouped in size 48 and smaller (<100 mm), size 40–27 (100–117 mm), and size 23 and larger (>117 mm) (Ferrarezi et al., 2019).

Fruit quality and yield of solids. The fruit acidity, soluble solids, ratio, and yield of solids were determined. The methods are available in Phuyal et al. (2020).

Plant photosynthesis measurement. Photosynthesis rate, *g_s*, intercellular CO₂, and transpiration rate were measured in April 2019 on two randomly selected trees per plot by using an infrared gas analyzer (LI-COR 6400XT; LI-COR Inc., Lincoln, NE). Two measurements were taken one week after FAM from 10:00 AM to 2:00 PM on sunny days from each sampled tree on different branches. Leaves were selected from a sunlit portion of the canopy. Immature, diseased, insect-damaged, mechanically injured, or dead leaves were avoided. Measurements were carried out in fully expanded mature citrus leaves as suggested by Iglesias et al. (2002). A constant flow rate of 500 μmol·s⁻¹, at 50 to 80% relative humidity, ambient reference CO₂, and photosynthetically active radiation was adjusted in the leaf chamber of the gas analyzer. The leaf was clamped in the leaf chamber until the steady-state photosynthesis level was reached.

Leaf sampling for nutrient analysis. Eight random trees were chosen excluding border trees to collect leaf samples. The leaf samples

were collected in August to analyze for leaf nutrient concentration. The protocol to collect, preserve, and analyze leaf samples are available in Phuyal et al. (2020).

Soil sampling and analysis. One 20-cm deep soil core/tree in 15 different locations/plot was taken in February to access the available soil nutrients and pH. Analysis followed methods available in Phuyal et al. (2020) and Mehlich (1984).

Statistical analysis. Statistical analysis was performed using SAS 9.4 (SAS Institute Inc., Cary, NC). Data were analyzed by year for two consecutive seasons. A generalized linear mixed model (Proc GLIMMIX) was used to analyze error variance where treatments were entered as fixed effect and block as a random effect. The data were checked for assumptions of the linear model. Fruit yield and yield of solids

data were not normally distributed, and were transformed to square root and log, respectively, to meet the assumptions of normality and homoscedasticity. Tukey test was used for multiple mean comparisons ($P < 0.05$). Correlations were calculated to find the relationship between the amount of foliar application with the total number of fruit yield per tree and per area (Proc CORR).

Table 2. Cycle threshold (Ct) value of *Candidatus Liberibacter asiaticus* (CLAs) DNA as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Ct value of CLAs DNA (unitless)		P value
	2018–19	2019–20	
CRF ^z			
CRF 1	24.95 ± 0.24 ^w	29.88 ± 0.17	
CRF 2	24.94 ± 0.26	29.96 ± 0.16	
PD ^y			
SR/LD	25.12 ± 0.39	29.83 ± 0.22	
SR/HD	24.91 ± 0.27	30.10 ± 0.20	
DR/HD	24.82 ± 0.28	29.82 ± 0.19	
FAM ^x			
0.0×	25.07 ± 0.34	29.98 ± 0.22	
1.5×	25.13 ± 0.41	30.01 ± 0.24	
3.0×	24.64 ± 0.34	30.03 ± 0.25	
6.0×	24.96 ± 0.34	29.66 ± 0.24	
Sources of variation			
CRF	NS	NS	
PD	NS	NS	
CRF*PD	NS	NS	
FAM	NS	NS	
CRF*FAM	NS	NS	
PD*FAM	NS	NS	
CRF*PD*FAM	NS	NS	

^zIn 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test. NS = nonsignificant.

Results

Candidatus Liberibacter asiaticus infection.

Planting density, CRF application, and FAM did not influence Ct values of *Candidatus Liberibacter asiaticus* (CLAs) DNA (Table 2, $P > 0.05$). All sampled trees were infected since values were below 32 (Albrecht and Bowman, 2011; Gottwald et al., 2012; Shin and van Bruggen, 2018).

Tree growth. Canopy volume and trunk diameter were both affected by planting density ($P < 0.0001$). Depending on season, canopy volume was 31% to 37% lower under DR/HD than under SR/LD and SR/HD (Table 3). Similarly, trunk diameter was 13% to 17% smaller under DR/HD than under SR/HD and SR/LD. Canopy volume, but not trunk diameter, was also affected by CRF blends. We found that CRF 1 resulted in 10% and 8% greater canopy volume than CRF 2 in 2018–19 and 2019–20, respectively (Table 3). Canopy volume was unaffected by foliar micronutrient application ($P > 0.05$).

Fruit yield and fruit size. For both seasons, the greatest fruit yields were obtained with DR/HD, which was 136% to 190% higher than under SR/LD (Table 4). Higher micronutrient application through CRF blends reduced the number of fruit in 2019–

Table 3. Trunk diameter and canopy volume as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Trunk diam (mm)		Canopy volume (m ³)	
	2018–19	2019–20	2018–19	2019–20
CRF ^z				
CRF 1	82.82 ± 1.16 ^w	87.69 ± 1.25	8.33 ± 0.28 a	8.63 ± 0.30 a
CRF 2	81.86 ± 1.08	86.49 ± 1.33	7.59 ± 0.25 b	7.98 ± 0.32 b
PD ^y				
SR/LD	88.81 ± 0.79 a	93.26 ± 1.22 a	9.18 ± 0.28 a	9.83 ± 0.32 a
SR/HD	84.88 ± 0.68 b	90.26 ± 0.95 a	8.56 ± 0.23 a	8.93 ± 0.26 b
DR/HD	73.33 ± 0.65 c	77.75 ± 0.92 b	6.13 ± 0.20 b	6.14 ± 0.19 c
FAM ^x				
0.0×	82.31 ± 1.69	87.63 ± 1.99	8.01 ± 0.44	8.59 ± 0.58
1.5×	81.42 ± 1.48	85.60 ± 2.00	7.75 ± 0.35	7.85 ± 0.41
3.0×	82.64 ± 1.69	88.17 ± 1.56	7.95 ± 0.36	8.31 ± 0.32
6.0×	83.01 ± 1.53	86.96 ± 1.77	8.12 ± 0.39	8.45 ± 0.43
Sources of variation				
CRF	NS	NS	**	**
PD	***	***	***	***
CRF*PD	NS	NS	NS	NS
FAM	NS	NS	NS	NS
CRF*FAM	NS	NS	NS	NS
PD*FAM	NS	NS	NS	NS
CRF*PD*FAM	NS	NS	NS	NS

^zIn 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, **, ***Nonsignificant or significant at $P < 0.01$ or 0.001, respectively.

Table 4. Fruit diameter, fruit yield per tree, and fruit yield per area as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Fruit diam (mm)		Fruit yield per tree (kg)		Fruit yield per area (kg·ha ⁻¹)	
	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20
CRF ²						
CRF 1	96.46 ± 0.60 ^w	98.39 ± 0.28	7.60 ± 0.40 ^w	16.97 ± 0.87	4,189.76 ± 270.08	8,634.24 ± 463.90
CRF 2	96.08 ± 0.77	98.89 ± 0.25	7.41 ± 0.31	18.53 ± 0.69	3,994.16 ± 212.45	9,724.25 ± 456.64
PD ³						
SR/LD	95.82 ± 0.83	100.16 ± 0.35 a	7.72 ± 0.48	18.69 ± 0.95 a	2,263.83 ± 142.00 c	5,482.64 ± 279.57 c
SR/HD	97.16 ± 0.37	99.03 ± 0.26 b	8.01 ± 0.44	21.18 ± 1.06 a	3,436.16 ± 189.41 b	9,080.97 ± 455.94 b
DR/HD	95.82 ± 1.15	96.73 ± 0.27 c	6.78 ± 0.37	13.37 ± 0.70 b	6,575.89 ± 363.70 a	12,974.12 ± 675.11 a
FAM ⁴						
0.0×	96.28 ± 1.11	98.87 ± 0.39 ab	8.54 ± 0.57 a	21.96 ± 1.40 a	4,533.22 ± 362.48 a	11,002.44 ± 763.59 a
1.5×	97.01 ± 0.44	97.81 ± 0.39 b	7.78 ± 0.51 ab	17.39 ± 1.07 b	4,248.30 ± 339.13 ab	8,991.36 ± 609.61 ab
3.0×	96.24 ± 1.12	98.69 ± 0.33 ab	7.59 ± 0.50 ab	17.33 ± 0.98 b	4,187.28 ± 374.52 ab	9,140.85 ± 636.23 ab
6.0×	95.52 ± 1.08	99.18 ± 0.36 a	6.11 ± 0.40 b	14.32 ± 0.75 b	3,399.05 ± 284.75 b	7,582.33 ± 542.77 b
Sources of variation				<i>P</i> value		
CRF	NS	NS	NS	NS	NS	NS
PD	NS	***	NS	***	***	***
CRF*PD	NS	NS	NS	NS	NS	NS
FAM	NS	*	**	***	*	***
CRF*FAM	NS	NS	NS	NS	NS	NS
PD*FAM	NS	*	NS	NS	NS	NS
CRF*PD*FAM	NS	NS	NS	NS	NS	NS

²In 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

³SR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

⁴Foliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05, 0.01, \text{ or } 0.001$, respectively.




Fig. 3. Fruit diameter under three planting density (PD) [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)]; and four foliar-applied micronutrient rates (FAM) (0, 1.5, 3, and 6 times) as a blend of B, Mn, and Zn according to the UF/IFAS year recommended rates. Means ± SE followed by different letters (uppercase compare FAM and lowercase compare PD) are significantly different by Tukey's test at $P < 0.05$.

20. Foliar nutrient application was negatively correlated with fruit yields in 2018–19 ($r = -0.122, P = 0.0168$) and 2019–20 ($r = -0.176, P = 0.0005$) and total fruit number (Table 5). In 2019–20, an interaction between planting densities and FAM micronutrient applications was observed, with SR/LD producing the largest-diameter fruit compared with other treatments (Fig. 3).

Fruit quality and yield of solids. Fruit from trees planted at DR/HD contained 5% more soluble solids than fruit from the SR/LD configuration in both seasons (Table 6). In addition, fruit acidity was 13% and 6% greater under DR/HD than other treatments in 2018–19 and 2019–20, respectively. In 2018–19, the acidic nature of the juice induced 7% lower ratio in the DR/HD planting.

In both seasons, fruit from the DR/HD configuration produced 130% to 200% greater yield of solids than other treatments (Table 6). In 2018–19, the CRF blend did not affect fruit quality (Table 6); however, in 2019–20 CRF 1 induced 10% higher soluble solids, 7% higher ratio, and 86% greater yield of solids than CRF 2. In 2019–20, an interaction was observed between CRF blends and planting density ($P = 0.0179$) whereby CRF 1 use in DR/HD planting produced higher yield of solids than other treatments (Fig. 4). In 2019–20, fruit increased acidity with an increase in the FAM rate applied (Table 6, $P = 0.0101$).

Plant physiology. Photosynthetic rates under DR/HD were 12% and 21% lower than under SR/LD and SR/HD, respectively. The SR/HD resulted in 16% higher g_s than DR/HD planting. Use of CRF 1 increased photosynthesis (17%), intercellular CO_2 (10%), g_s (45%), and transpiration (46%) compared with CRF 2 (Table 7); however, FAM application reduced the assimilation rate, intercellular CO_2 , and g_s . A significant interaction of intercellular CO_2 was observed between tree planting density and CRF blends. The CRF 1 resulted in better intercellular CO_2 in all the planting densities (Fig. 5).

Leaf nutrient concentration. Leaf calcium (Ca) and N concentrations ranged from 3.0 to 4.9 and 2.5 to 2.7 g·kg⁻¹ under all treatments, respectively (Table 8), which are within the optimal ranges defined by Koo et al. (1984). In addition, leaf P, Fe, and Mg concentration decreased over time in all the treatments. Leaf copper (Cu) concentration was excessively high (>20 mg·kg⁻¹) under all treatments due to frequent application of fungicides containing Cu to control citrus

canker disease (*Xanthomonas axonopodis* pv. citri). In 2018–19, leaf Mn concentration was optimal (25–100 mg·kg⁻¹) for all treatments, except 6 times foliar spray of recommended rate, where it exceeded the optimum range. In 2018–19, leaf sulfur (S) concentration showed an interaction between planting density and CRF blends, in which CRF 1 showed higher leaf S concentration (Fig. 6). It could be due to the higher amount of S-coated

fertilizer present in CRF 1 blend. In 2019–20, leaf B, Mn, and Zn concentration showed an interaction between planting and CRF blends (Fig. 7).

Soil nutrient concentrations and soil pH. There was only a planting density effect on soil nutrient concentrations in 2019–20 (Tables 9 and 10). DR/HD plots had higher soil concentrations of K, Ca, Mg, S, and B, but lower P concentration in comparison to

other treatments. Higher availability of nutrients in soil sampled from high-density plantings could be due to less uptake by trees due to reduced root volume from HLB damage.

The CRF blends resulted in changes in soil pH and nutrient availability (Tables 9 and 10). In 2018–19, CRF 2 (CRF blend with lower micronutrient rates) increased soil pH by 6%, soil B concentration by 100%, and decreased Mn concentration by 55%. In 2019–20, the use of CRF 2 increased soil pH by 12%, B, Mg, and Zn concentrations by 25%, 13%, and 19%, respectively, whereas S concentrations fell by 42% compared with use of CRF 1. Interestingly, despite the lower B concentration in CRF 2, B availability in soil was higher relative to CRF 1 (Table 11). Both CRF blends resulted in optimum soil pH for most of the nutrients as suggested by Morgan and Kadyampakeni (2020). Using the soil test interpretation outlined by Morgan and Kadyampakeni (2020), the Mehlich 3 acid extraction showed a medium to high level of soil P concentrations, whereas Mg concentrations were low in 2018–19 but medium to high in 2019–20. In 2019–20, FAM only affected soil Mn and Zn concentrations (Table 10), which was probably caused by the spray solution reaching the soil. Boaretto et al. (2002) also observed a similar result with a foliar spray of Zn in orange trees.




Fig. 4. Yield of solids under two controlled-release fertilizers (CRF) blends [CRF 1 (12N–1.31P–7.47K with 2.5 times micronutrients and 2 times Mg) and CRF 2 (16N–1.31P–16.6K)]; and three planting densities [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)]. Means ± SE followed by different letters (uppercase compare CRF and lowercase compare PD) are significantly different by Tukey's test at $P < 0.05$.

Discussion

Clas infection. Our study showed no treatment effect on Ct value, which differs from earlier work by Zhang et al. (2016) that showed Zn application on 2-year-old

Table 5. Fruit size and total number of fruit as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Small size fruit (less than 100 mm)		Medium size fruit (100 mm–117 mm)		Large size fruit (above 100 mm)		Total no. of fruit	
	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20
CRF^z								
CRF 1	18.64 ± 0.98 ^w	41.74 ± 1.93	6.04 ± 0.41	14.36 ± 1.28 b	0.46 ± 0.08	0.82 ± 0.12	25.15 ± 1.25	56.93 ± 2.67 b
CRF 2	18.12 ± 0.85	46.11 ± 1.63	6.42 ± 0.37	18.23 ± 1.08 a	0.42 ± 0.06	0.72 ± 0.08	24.94 ± 1.04	65.06 ± 2.17 a
PD^y								
SR/LD	18.23 ± 1.12	40.35 ± 1.87 b	6.58 ± 0.54	20.73 ± 1.64 a	0.40 ± 0.10	1.02 ± 0.14 a	25.21 ± 1.46	62.1 ± 2.83 a
SR/HD	19.63 ± 1.21	50.93 ± 2.49 a	6.70 ± 0.47	19.43 ± 1.53 a	0.46 ± 0.08	0.98 ± 0.16 a	26.79 ± 1.47	71.34 ± 3.35 a
DR/HD	17.27 ± 1.03	40.51 ± 2.05 b	5.41 ± 0.40	8.73 ± 0.85 b	0.46 ± 0.07	0.30 ± 0.06 b	23.13 ± 1.28	49.55 ± 2.46 b
FAM^x								
0.0×	21.72 ± 1.57 a	48.74 ± 2.75 a	6.82 ± 0.58	22.10 ± 2.31 a	0.43 ± 0.09	0.84 ± 0.14	28.97 ± 1.89 a	71.69 ± 4.16 a
1.5×	18.93 ± 1.19 ab	47.53 ± 2.90 a	6.39 ± 0.50	14.47 ± 1.60 b	0.48 ± 0.08	0.65 ± 0.17	25.79 ± 1.48 ab	62.65 ± 3.56 a
3.0×	17.97 ± 1.26 ab	44.75 ± 2.31 a	6.77 ± 0.62	15.03 ± 1.37 b	0.57 ± 0.13	0.67 ± 0.12	25.31 ± 1.65 ab	60.45 ± 3.14 ab
6.0×	14.90 ± 1.04 b	34.70 ± 1.80 b	4.93 ± 0.48	13.58 ± 1.12 b	0.28 ± 0.07	0.92 ± 0.16	20.1 ± 1.32 b	49.20 ± 2.42 b
Sources of variation								
CRF	NS	NS	NS	*	NS	NS	NS	*
PD	NS	***	NS	***	NS	***	NS	***
CRF*PD	NS	NS	NS	NS	NS	NS	NS	NS
FAM	**	***	NS	***	NS	NS	**	***
CRF*FAM	*	NS	NS	NS	NS	NS	NS	NS
PD*FAM	NS	NS	NS	*	NS	NS	NS	NS
CRF*PD*FAM	NS	*	NS	NS	NS	NS	NS	NS

^zIn 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05$, 0.01, or 0.001, respectively.

Table 6. Soluble solids content, acidity, ratio, and yield of solids as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Soluble solids content (%)		Titratable acidity (% citric acid)		Ratio		Yield of solids (kg·ha ⁻¹)	
	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20
CRF²								
CRF 1	7.96 ± 0.10 ^w	8.31 ± 0.11 a	1.07 ± 0.02	0.89 ± 0.01	7.51 ± 0.10	9.33 ± 0.15 a	159.72 ± 16.68	885.62 ± 100.57 a
CRF 2	7.74 ± 0.12	7.52 ± 0.14 b	1.08 ± 0.01	0.87 ± 0.01	7.21 ± 0.12	8.72 ± 0.24 b	130.55 ± 12.57	476.34 ± 73.57 b
PD³								
SR/LD	7.67 ± 0.10 b	7.63 ± 0.17 b	1.03 ± 0.02 b	0.86 ± 0.02 b	7.62 ± 0.11 a	8.93 ± 0.27	85.79 ± 7.03 b	343.20 ± 30.98 b
SR/HD	7.77 ± 0.11 ab	7.9 ± 0.16 ab	1.02 ± 0.01 b	0.87 ± 0.01 b	7.46 ± 0.11 a	9.14 ± 0.25	117.08 ± 9.17 b	472.60 ± 50.81 b
DR/HD	8.11 ± 0.18 a	8.22 ± 0.17 a	1.16 ± 0.01 a	0.92 ± 0.01 a	6.99 ± 0.15 b	8.99 ± 0.23	232.53 ± 22.24 a	1,227.14 ± 145.33 a
FAM⁴								
0.0×	8.05 ± 0.16	7.98 ± 0.22	1.07 ± 0.02	0.87 ± 0.01 b	7.57 ± 0.12	9.22 ± 0.26	166.73 ± 22.83	683.14 ± 125.97
1.5×	7.80 ± 0.13	8 ± 0.2	1.07 ± 0.02	0.86 ± 0.02 b	7.32 ± 0.12	9.46 ± 0.35	150.46 ± 19.68	663.23 ± 116.45
3.0×	7.60 ± 0.18	7.8 ± 0.2	1.08 ± 0.02	0.89 ± 0.01 ab	7.12 ± 0.21	8.84 ± 0.25	142.47 ± 23.75	662.11 ± 123.44
6.0×	7.94 ± 0.15	7.88 ± 0.19	1.07 ± 0.02	0.92 ± 0.01 a	7.43 ± 0.13	8.57 ± 0.25	120.89 ± 17.35	715.43 ± 161.23
Sources of variation								
CRF	NS	***	NS	NS	NS	*	NS	***
PD	*	*	***	**	***	NS	***	***
CRF*PD	NS	NS	NS	NS	NS	NS	NS	*
FAM	NS	NS	NS	*	NS	NS	NS	NS
CRF*FAM	NS	NS	NS	NS	NS	NS	NS	NS
PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS
CRF*PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS

²In 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

³SR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

⁴Foliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05, 0.01, \text{ or } 0.001$, respectively.

Table 7. Assimilation rate, intercellular CO₂, stomatal conductance (g_s), and transpiration rate as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Assimilation rate (μmol CO ₂ /m ² /s)	Intercellular CO ₂ (μmol CO ₂ /mol)	g _s (mol H ₂ O/m ² /s)	Transpiration rate (mmol H ₂ O/m ² /s)
CRF²				
CRF 1	11.24 ± 0.23 a ^w	264.36 ± 2.55 a	0.16 ± 0.01 a	2.67 ± 0.08 a
CRF 2	9.61 ± 0.26 b	238.97 ± 2.51 b	0.11 ± 0.01 b	1.83 ± 0.07 b
PD³				
SR/LD	10.46 ± 0.31 b	248.41 ± 3.45	0.14 ± 0.01 ab	2.20 ± 0.09
SR/HD	11.62 ± 0.26 a	258.09 ± 3.55	0.15 ± 0.01 a	2.39 ± 0.11
DR/HD	9.19 ± 0.30 c	248.49 ± 2.77	0.12 ± 0.01 b	2.16 ± 0.09
FAM⁴				
0.0×	11.34 ± 0.38 a	254.57 ± 3.59 ab	0.16 ± 0.01 a	2.41 ± 0.11
1.5×	10.24 ± 0.33 ab	244.4 ± 3.39 b	0.13 ± 0.01 b	2.20 ± 0.11
3.0×	9.83 ± 0.37 ab	250 ± 4.12 ab	0.13 ± 0.01 b	2.16 ± 0.11
6.0×	10.28 ± 0.30 b	257.67 ± 3.98 a	0.13 ± 0.01 b	2.23 ± 0.13
Sources of variation				
CRF			<i>P</i> value	
CRF	***	***	***	***
PD	***	NS	***	NS
CRF*PD	NS	*	NS	NS
FAM	**	*	**	NS
CRF*FAM	NS	NS	NS	NS
PD*FAM	NS	NS	NS	NS
CRF*PD*FAM	NS	NS	NS	NS

²In 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

³SR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

⁴Foliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05, 0.01, \text{ or } 0.001$, respectively.

grapefruit seedlings lowered Ct values. The absence of a soil-applied micronutrient or Mg effect on Ct values may have been due to the relatively short length of our study. As with soil-applied nutrients, we found that FAMs did not affect Ct values. Gottwald et al. (2012) similarly observed no visual differences or disease expression in HLB-affected 'Valencia' trees treated with different micronutrient applications. A longer trial period may be needed to observe dif-

ferences in disease expression (Shen et al., 2013) and as was demonstrated in sweet orange by Morgan et al. (2016) in their 5-year study. Stansly et al. (2014) noted that nutritional treatments can improve tree growth, which increases opportunities for psyllid feeding and ultimately causes higher bacterial titer or lower Ct value. These results suggest that FAMs may be counterproductive unless psyllid control measures are effective.

Tree growth. The decrease in trunk diameter and canopy volume as planting density increased was expected because high-density planting increases competition for water, nutrients, and solar radiation. As with CRF 1 (blend with higher nutrient application), Morgan and Kadyampakeni (2020) indicated the higher nutrient application results in greater tree growth with the expense of yield, as seen in reduced number of fruit in 2019–20 in our study (Table 4). FAM did not affect

Table 8. Leaf macronutrient concentration as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrient (FAM).

Treatments	N (g·kg ⁻¹)			P (g·kg ⁻¹)			K (g·kg ⁻¹)			Ca (g·kg ⁻¹)			Mg (g·kg ⁻¹)			S (g·kg ⁻¹)		
	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20		
CRF ^z																		
CRF 1	2.83 ± 0.03 ^w	2.73 ± 0.02	0.18 ± 0.0025	0.15 ± 0.0013	1.22 ± 0.03 b	0.98 ± 0.01	4.79 ± 0.03 a	3.20 ± 0.03	0.34 ± 0.0034	0.26 ± 0.0027 b	0.4 ± 0.0046 a	0.31 ± 0.0038 a						
CRF 2	2.84 ± 0.03	2.76 ± 0.03	0.18 ± 0.0022	0.15 ± 0.0009	1.44 ± 0.03 a	1.02 ± 0.02	4.35 ± 0.05 b	3.24 ± 0.04	0.34 ± 0.01	0.27 ± 0.0026 a	0.37 ± 0.01 b	0.29 ± 0.0032 b						
PD ^y																		
SR/LD	2.80 ± 0.03	2.74 ± 0.04	0.18 ± 0.0029	0.15 ± 0.0015	1.28 ± 0.04	1.01 ± 0.02	4.57 ± 0.06	3.20 ± 0.04	0.34 ± 0.01	0.26 ± 0.0033	0.39 ± 0.01	0.30 ± 0.0047						
SR/HD	2.81 ± 0.03	2.76 ± 0.03	0.18 ± 0.0026	0.15 ± 0.0013	1.32 ± 0.03	0.99 ± 0.02	4.59 ± 0.04	3.26 ± 0.03	0.33 ± 0.0043	0.27 ± 0.0028	0.38 ± 0.0036	0.29 ± 0.003						
DR/HD	2.89 ± 0.04	2.73 ± 0.02	0.19 ± 0.0032	0.15 ± 0.0014	1.40 ± 0.04	1.00 ± 0.03	4.55 ± 0.07	3.20 ± 0.06	0.34 ± 0.01	0.27 ± 0.0035	0.39 ± 0.01	0.30 ± 0.01						
FAM ^x																		
0.0x	2.81 ± 0.04	2.75 ± 0.03	0.18 ± 0.0038	0.15 ± 0.0017	1.29 ± 0.05	0.95 ± 0.02 b	4.56 ± 0.09	3.29 ± 0.05	0.34 ± 0.01	0.27 ± 0.0032	0.38 ± 0.01	0.30 ± 0.0042						
1.5x	2.86 ± 0.04	2.71 ± 0.02	0.18 ± 0.0025	0.15 ± 0.0016	1.32 ± 0.04	0.96 ± 0.02 b	4.57 ± 0.07	3.26 ± 0.04	0.34 ± 0.01	0.27 ± 0.0039	0.38 ± 0.01	0.30 ± 0.0046						
3.0x	2.85 ± 0.04	2.72 ± 0.03	0.18 ± 0.0033	0.15 ± 0.0015	1.34 ± 0.04	1.03 ± 0.03 ab	4.60 ± 0.07	3.23 ± 0.05	0.34 ± 0.01	0.26 ± 0.0039	0.39 ± 0.01	0.30 ± 0.01						
6.0x	2.82 ± 0.04	2.80 ± 0.04	0.18 ± 0.0037	0.16 ± 0.0017	1.39 ± 0.04	1.08 ± 0.02 a	4.55 ± 0.07	3.10 ± 0.05	0.34 ± 0.01	0.26 ± 0.0034	0.39 ± 0.01	0.29 ± 0.01						
Sources of variation																		
CRF	NS	NS	NS	NS	*	NS	***	NS	NS	*	***	***						
PD	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS						
CRF*PD	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	*	NS						
FAM	NS	NS	NS	NS	NS	***	NS	NS	NS	NS	NS	NS						
CRF*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS						
PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS						
CRF*PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS						

^zIn 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, ***, Nonsignificant or significant at $P < 0.01$ or 0.001 , respectively.




Fig. 5. Intercellular CO₂ under two controlled-release fertilizer (CRF) blends [CRF 1 (12N–1.31P–7.47K with 2.5 times micronutrients and 2 times Mg) and CRF 2 (16N–1.31P–16.6K)]; and three planting density (PD) [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)]. Means ± SE followed by different letters (uppercase compare CRF and lowercase compare PD) are significantly different by Tukey's test at $P < 0.05$.




Fig. 6. Leaf S concentration under two controlled-release fertilizers (CRF) blends [CRF 1 (12N–1.31P–7.47K with higher amount of micronutrient) and CRF 2 (16N–1.31P–16.6K)]; and three planting density (PD) [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)]. Means ± SE followed by different letters (uppercase compare CRF and lowercase compare PD) are significantly different by Tukey's test at $P < 0.05$.

canopy volume in HLB-affected orange trees as in Rouse et al. (2017). A long-term study could manifest differences in canopy volume with the application of foliar treatments, such as in Morgan et al. (2016).

Fruit yield and fruit size. High-density planting with UF/IFAS-recommended nutrient application increased the fruit yield but reduced the total number of fruit. The higher soil nutrient application tended to increase

canopy growth at the expense of fruit yield. In addition, in our study, FAM showed negative effect on fruit yield and fruit size (Table 5). The effect of foliar nutrient application on citrus fruit yield is a topic of controversy. Several studies have shown that foliar application alone or in combination with plant hormones and insecticides can increase fruit yields in HLB-affected orchards (Al-Obeed et al., 2018; Morgan

et al., 2016; Stansly et al., 2014). However, other studies indicated foliar nutrient application is not effective for enhancing fruit yield and quality (Boaretto et al., 1997, 2002; Gottwald et al., 2012). In our study, the negative correlation between foliar nutrient application and fruit yield may have been due in part to vegetative growth, as was also reported in sweet orange by Morgan et al. (2016).

Fruit quality and yield of solids. The fruit quality was affected by planting density, as in Phuyal et al. (2020), suggesting that high-density planting results in the accumulation of more sugars with acidic juice. As with CRF, we obtained greater soluble solids from higher micronutrient application. With all the treatments, the ratio was higher than the standard ratio outlined for grapefruit grade B which is 7:1 (U.S. Department of Agriculture, 2012). In our study, we observed an increase in fruit acidity with FAM application possibly due to increase in leaf K and Zn concentration with the application of higher rates of micronutrients, like Nasir et al. (2016) observed in mandarin.

Plant physiology. Because photosynthesis is a physiological process requiring sunlight, high-density planting could have impaired photosynthetic rates as a result of shading.

Goldman (2010) noted that micronutrients are important for plant photosynthesis, particularly Mn for chlorophyll formation, and Zn as an important cofactor for enzyme carbonic anhydrase. To our knowledge, the effect of soil elevated micronutrient application on grapefruit plant physiology under field conditions has never been tested before. We observed elevated soil micronutrient concentrations can improve plant photosynthesis, thereby enhancing canopy growth. Net photosynthesis and g_s were negatively affected by increasing FAM rates, contradicting earlier findings by Ilyas et al. (2015) that revealed an increase in net photosynthetic rate, g_s , and transpiration with foliar application of B, Cu, and Zn. Our findings show that FAM may have caused plant stress or excessive growth due to luxurious nutrient supply (Table 9). Morgan and Connolly (2013) suggested that excessive micronutrient accumulation can damage plant cells by generating reactive oxygen species.

Leaf nutrient concentration. There was a general trend showing that use of CRF 1 resulted in similar or higher leaf micronutrient concentrations than using CRF 2 for both seasons. This outcome was expected because CRF 1 contained higher amounts of micronutrients than CRF 2. As with CRF, FAM application also affected foliar micronutrient concentrations. There was a positive correlation between FAM and leaf B, Mn, and Zn concentration in 2018–19. In 2019–20, this trend was limited to Mn and Zn. There was also an increase in K and decrease in Cu concentrations, respectively. B, Zn, or their combination may have facilitated K uptake, as was previously noted by Mengel and Kirkby (1980). The decrease in Cu concentration with increase in foliar rate was




Fig. 7. The concentration of B (A), Mn (B), and Zn (C) in the leaves under two controlled-release fertilizers (CRF) blends [CRF 1 (12N–1.31P–7.47K with 2.5 times micronutrients and 2 times Mg) and CRF 2 (16N–1.31P–16.6K)]; and three planting density (PD) [single-row low-density (SR/LD, 300 trees per ha), single-row high-density (SR/HD, 440 trees per ha), and double-row high-density staggered in diamond setting (DR/HD, 975 trees per ha)]. Means \pm SE followed by different letters (uppercase compares CRF and lowercase compare PD) are significantly different by Tukey's test at $P < 0.05$.

possibly because of an antagonistic effect of Zn. Studies have shown the antagonisms of Zn on Cu nutrition in wheat (*Triticum aestivum*) and rice (*Oryza sativa*) (Chaudhry et al., 1973; Imtiaz et al., 2003).

Soil nutrient concentrations and soil pH.

In our study, soil samples from CRF 2 treatment showed higher B concentration in the soil even though we applied less B than CRF 1. Goldberg et al. (2000) reported that soil B availability depends on several factors, including soil texture, moisture, temperature, and oxide content along with soil pH because low pH favors B uptake. Zhu et al. (2007) reported that the B concentration in soils is generally related to the amount of B taken up by plants. In our study, CRF 1 decreased soil pH which could have facilitated B uptake by the trees. In addition, the increase in S content with CRF 1 is probably from the higher application of S-coated fertilizer. The application of higher amount of Mg via CRF 1 shows 11% lower Mg concentration in soil (Table 10) potentially because of the interaction between nutrient and soil pH as Mg is highly mobile element in soil and shows its maximum availability in the soil pH range of 7 to 8 (Gransee and Fühns, 2013; Truog, 1947). Prior studies showed that Mg availability reduces drastically in the soil with a

small increase in soil pH (Hailes et al., 1997; Sumner et al., 1978). In our study, CRF 1 resulted in a low soil pH (5.63), which was 11% lower than CRF 2. The lower pH in CRF 1 could have decreased the Mg availability in the soil.

Conclusions

The planting density did not affect Ct value. High-density planting resulted in smaller trunk diameter and canopy volume. Despite lower yield per tree in 2019–20, high-density planting induced the greatest fruit yields on an area basis. Also, fruit produced under high-density plantings had higher soluble solids and was more acidic. Higher micronutrient application did not affect fruit yield but did improve tree physiological parameters and canopy volume. FAM did not affect HLB incidence, tree growth, or fruit quality. However, fruit yield and physiological parameters decreased as FAM increased. Over the 2-year period, our study showed that excessive supplemental nutrient application may increase vegetative growth at the cost of reduced yield and may not be beneficial for grapefruit trees affected by HLB, whereas high-density plantings can improve yield. Moving forward, manage-

ment costs, along with long-term yield response to higher-density plantings should be determined before recommendations can be made to growers. In addition, information is needed on nutritional treatment effects on other HLB-tolerant rootstocks.

Literature Cited

- Albrecht, U. and K.D. Bowman. 2011. Tolerance of the trifoliolate citrus hybrid US-897 (*Citrus reticulata* Blanco \times *Poncirus trifoliata* L. Raf.) to Huanglongbing. *HortScience* 46:16–22.
- Al-Obeed, R.S., M.A.A. Ahmed, H.A. Kassem, and A.M. Al-Saif. 2018. Improvement of “Kinnow” mandarin fruit productivity and quality by urea, boron, and zinc foliar spray. *J. Plant Nutr.* 41:609–618.
- Boaretto, A.E., C.S. Tiritan, and T. Muraoka. 1997. Effects of foliar applications of boron on citrus fruit and on foliage and soil boron concentration, p. 121–123. In: R.W. Bell and B. Rekasem (eds.). *Boron in soils and plants*. Springer, Dordrecht.
- Boaretto, A.E., R.M. Boaretto, T. Muraoka, V.F. Nascimento Filho, C.S. Tiritan, and F.A.A. Mourão Filho. 2002. Foliar micronutrient application effects on citrus fruit yield, soil, and leaf Zn concentrations and ⁶⁵Zn mobilization within the plant. *Acta Hort.* 594: 203–209.
- Chaudhry, F.M., M. Sharif, A. Latif, and R.H. Qureshi. 1973. Zinc-copper antagonism in the nutrition of rice (*Oryza sativa*). *Plant Soil* 38:573–580.
- Dalal, R.P.S., A.K. Sangwan, B.S. Beniwal, and S. Sharma. 2013. Effect of planting density on canopy parameter, yield and water use efficiency of Kinnow mandarin. *Indian J. Hort.* 70:587–590.
- Etcheberria, E., P. Gonzalez, D. Achor, and G. Albrigo. 2009. Anatomical distribution of abnormally high levels of starch in HLB-affected Valencia orange trees. *Physiol. Mol. Plant Pathol.* 74:76–83.
- Ferrarezi, R.S., J.A. Qureshi, A.L. Wright, M.A. Ritenour, and N.P. Macan. 2019. Citrus production under screen as a strategy to protect grapefruit trees from Huanglongbing disease. *Frontiers in Plant Sci.* 10:1598.
- Gerendás, J. and H. Fühns. 2013. The significance of magnesium for crop quality. *Plant Soil* 368:101–128.
- Goldberg, S., S.M. Lesch, and D.L. Suarez. 2000. Predicting boron adsorption by soils using soil chemical parameters in the constant capacitance model. *Soil Sci. Soc. Amer. J.* 64:1356–1363.
- Goldman, C.R. 2010. Micronutrient elements (Co, Mo, Mn, Zn, Cu), p. 378–382. In: *Biogeochemistry of inland waters*. Academic Press, San Diego, CA.
- Gonzalez, P., J. Reyes-De-Corcuera, and E. Etcheberria. 2012. Characterization of leaf starch from HLB-affected and unafflicted-girdled citrus trees. *Physiol. Mol. Plant Pathol.* 79:71–78.
- Gottwald, T.R., J.H. Graham, M.S. Irey, T.G. McCollum, and B.W. Wood. 2012. Inconsequential effect of nutritional treatments on Huanglongbing control, fruit quality, bacterial titer, and disease progress. *Crop Prot.* 36:73–82.
- Gransee, A. and H. Fühns. 2013. Magnesium mobility in soils as a challenge for soil and plant analysis, magnesium fertilization and root uptake under adverse growth conditions. *Plant Soil* 368:5–21.

Table 9. Leaf micronutrient as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrient (FAM) in 2018–19 and 2019–20.

Treatments	B (mg·kg ⁻¹)		Mn (mg·kg ⁻¹)		Zn (mg·kg ⁻¹)		Cu (mg·kg ⁻¹)		Fe (mg·kg ⁻¹)	
	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20	2018–19	2019–20
CRF ^z										
CRF 1	148.42 ± 3.07 a ^w	166.48 ± 5.06 a	93.04 ± 6.36 a	181.46 ± 8.49 a	74 ± 6.16 a	162.44 ± 8.21 a	248.03 ± 12.71	208.67 ± 8.27 a	99.95 ± 3.01 a	61.25 ± 1.24 a
CRF 2	117.13 ± 3.27 b	145.02 ± 3.35 b	62.14 ± 4.93 b	139.67 ± 7.67 b	51.74 ± 3.64 b	137.79 ± 7.99 b	242.80 ± 9.26	174.81 ± 7.84 b	77.43 ± 2.49 b	56.02 ± 1.05 b
PD ^y										
SR/LD	130.84 ± 4.38	143.84 ± 4.30 a	83.03 ± 7.70	137.97 ± 8.31 b	66.66 ± 6.65	133.69 ± 8.59 b	243.87 ± 16.29	188.50 ± 9.22	92.24 ± 4.56	58.41 ± 1.78
SR/HD	129.90 ± 3.82	158.47 ± 4.77 ab	67.16 ± 7.31	170.75 ± 10.25 a	68.17 ± 6.84	163.38 ± 9.56 a	253.70 ± 11.46	197.41 ± 10.93	88.98 ± 3.93	56.44 ± 1.31
DR/HD	137.59 ± 5.87	164.94 ± 6.80 b	82.58 ± 7.24	172.97 ± 11.87 a	53.77 ± 5.79	153.28 ± 11.56 ab	238.67 ± 12.76	189.31 ± 10.82	84.85 ± 3.14	61.06 ± 1.21
FAM ^x										
0.0x	122.69 ± 4.51 b	156.08 ± 7.40	47.77 ± 3.92 c	113 ± 9.26 b	38.92 ± 2.85 c	100.29 ± 9.15 b	248.46 ± 15.75	216.25 ± 11.86 a	86.93 ± 4.02	61.33 ± 1.64
1.5x	128.16 ± 4.26 ab	154.17 ± 4.69	62.71 ± 4.07 bc	172 ± 9.09 b	50.22 ± 2.90 bc	170.92 ± 9.05 a	248.83 ± 14.92	209.63 ± 11.29 a	87.74 ± 3.74	57.08 ± 1.45
3.0x	138.23 ± 5.14 ab	159.63 ± 7.11	86.13 ± 7.06 b	175.71 ± 12.64 b	71.22 ± 7.38 ab	168.83 ± 11.67 a	251.71 ± 16.81	182 ± 11.26 ab	91.01 ± 4.69	57.71 ± 1.94
6.0x	142.01 ± 7.02 a	153.13 ± 6.49	113.76 ± 10.77 a	181.54 ± 12.22 a	91.11 ± 9.50 a	160.42 ± 10.71 a	232.66 ± 15.80	159.08 ± 9.76 b	89.08 ± 5.64	58.42 ± 1.73
Sources of variation										
CRF	***	***	***	***	***	NS	NS	NS	***	**
PD	NS	*	NS	**	NS	*	NS	NS	NS	NS
CRF*PD	NS	*	NS	*	NS	*	NS	NS	NS	NS
FAM	*	NS	***	***	***	***	NS	NS	NS	NS
CRF*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
CRF*PD*FAM	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

^zIn 2018–19, CRF 1 = 12N–1.31P–7.47K with higher micronutrient content; and CRF 2 = 16N–1.31P–16.6K. In 2019–20, CRF 1 = 12N–1.31P–11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N–1.31P–11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, ** ***Nonsignificant or significant at $P < 0.05$, 0.01, or 0.001, respectively.

Hailes, K.J., R.L. Aitken, and N.W. Menzies. 1997. Magnesium in tropical and subtropical soils from north-eastern Australia. II. Response by glasshouse-grown maize to applied magnesium. *Austral. J. Soil Res.* 35:629–642.

Huber, D.M. and J.B. Jones. 2013. The role of magnesium in plant disease. *Plant Soil* 368:73–85.

Ibrahim, M., N. Ahmad, S.A. Anwar, and T. Majeed. 2007. Effect of micronutrients on citrus fruit yield growing on calcareous soils, p. 179–182. In: F. Xu, H.E. Goldbach, P.H. Brown, et al. *Advances in plant and animal boron nutrition*. Springer, Dordrecht, The Netherlands.

Iglesias, D.J., I. Lliso, F.R. Tadeo, and M. Talon. 2002. Regulation of photosynthesis through source: Sink imbalance in citrus is mediated by carbohydrate content in leaves. *Physiol. Plant.* 116:563–572.

Ilyas, A., M.Y. Ashraf, M. Hussain, M. Ashraf, R. Ahmed, and A. Kamal. 2015. Effect of micronutrients (Zn, Cu and B) on photosynthetic and fruit yield attributes of *Citrus reticulata* Blanco var. kinnow. *Pak. J. Bot.* 47:1241–1247.

Imtiaz, M., B.J. Alloway, K.H. Shah, S.H. Siddiqui, M.Y. Memon, M. Aslam, and P. Khan. 2003. Zinc nutrition of wheat: II: Interaction of zinc with other trace elements. *Asian J. Plant Sci.* 2:156–160.

Koo, R.C.J., C.A. Anderson, I. Stewart, D.P.H. Tucker, D.V. Calvert, and H.K. Wutscher. 1984. Recommended fertilizers and nutritional sprays for citrus. Florida Agricultural Expt. Sta. Bul. 536D.

Labanauskas, C. 1963. Foliar sprays correct manganese deficiencies on desert grapefruit. *Calif. Agr.* 17:14–15. 1 Apr. 2020. <<http://hilgardia.ucanr.edu/fileaccess.cfm?article=174679&p=SRNIQO>>.

Ladaniya, M.S., R.A. Marathe, A.K. Das, C.N. Rao, A.D. Huchche, P.S. Shirgure, and A.A. Murkute. 2020. High density planting studies in acid lime (*Citrus aurantifolia* Swingle). *Scientia Hort.* 261:108935.

Li, C.P., Y. Qi, J. Zhang, L. Yang, D.H. Wang, X. Ye, N.W. Lai, L.L. Tan, D. Lin, and L.S. Chen. 2017. Magnesium-deficiency-induced alterations of gas exchange, major metabolites and key enzymes differ among roots, and lower and upper leaves of *Citrus sinensis* seedlings. *Tree Physiol.* 37:1564–1581.

Li, S.L., Z.G. Li, and Z.L. He. 2014. Effect of enhanced zinc nutrition on mitigation of Huanglongbing (HLB)-affected citrus. *J. Citrus Pathol.* 1:7.26 P (abstr.).

Lovatt, C.J. 1999. Timing citrus and avocado foliar nutrient applications to increase fruit set and size. *HortTechnology* 9:607–612.

Lovatt, C.J. 2013. Properly timing foliar-applied fertilizers increases efficacy: A review and update on timing foliar nutrient applications to citrus and avocado. *HortTechnology* 23:536–541.

Marschner, H. 2012. *Marschner's mineral nutrition of higher plants*. 2nd ed. Academic Press, Cambridge, MA.

Mehlich, A. 1984. Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant. *Commun. Soil Sci. Plant Anal.* 15:1409–1416.

Mengel, K. and E.A. Kirkby. 1980. Potassium in crop production, p. 59–110. In: N.C. Brady, ed. *Advances in agronomy*. Academic Press, Cambridge, MA.

Moreira, A.S., E.S. Stuchi, P.R.B. Silva, R.B. Bassanezi, E.A. Girardi, and F.F. Laranjeira. 2019. Could tree density play a role in

Table 10. Soil pH and soil macronutrient concentration as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatments	Soil pH			P (mg·kg ⁻¹)			K (mg·kg ⁻¹)			Ca (mg·kg ⁻¹)			Mg (mg·kg ⁻¹)			S (mg·kg ⁻¹)			
	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	
CRF ^z																			
CRF 1	6.00 ± 0.12 b ^w	5.63 ± 0.08 b	15.88 ± 1.06	35.77 ± 1.14	20.29 ± 1.75	41.75 ± 1.25	350.92 ± 20.37	429.76 ± 17.82 b	19.29 ± 1.03	36.28 ± 1.13 b	18.08 ± 2.40	18.27 ± 0.78 a							
CRF 2	6.38 ± 0.12 a	6.36 ± 0.07 a	13.67 ± 1.50	36.20 ± 0.89	21.00 ± 1.46	42.09 ± 1.57	422.13 ± 56.55	521.35 ± 21.59 a	24.08 ± 1.20	41.16 ± 1.22 a	13.75 ± 2.07	10.60 ± 0.44 b							
PD ^y																			
SR/LD	6.00 ± 0.19	5.58 ± 0.11 b	16.44 ± 1.74	34.39 ± 1.28 a	19.25 ± 1.51	38.84 ± 1.40 b	403.06 ± 89.50	386.64 ± 24.8 b	20.38 ± 1.42	32.75 ± 1.13 b	15.19 ± 3.20	12.94 ± 0.99 b							
SR/HD	6.40 ± 0.12	6.22 ± 0.10 a	14.31 ± 1.33	39.75 ± 1.00 a	21.25 ± 2.11	40.77 ± 1.59 ab	379.88 ± 28.36	512.55 ± 23.93 a	23.25 ± 2.26	41.30 ± 1.55 a	15.38 ± 2.73	14.50 ± 0.90 ab							
DR/HD	6.18 ± 0.13	6.19 ± 0.09 ab	13.56 ± 1.75	33.81 ± 1.21 b	21.44 ± 2.28	46.16 ± 1.93 a	376.63 ± 13.73	527.48 ± 20.31 a	21.44 ± 0.79	42.11 ± 1.21 a	17.19 ± 2.76	15.88 ± 1.16 a							
FAM ^f																			
0.0x	N/A	6.00 ± 0.13	N/A	37.46 ± 1.48	N/A	43.58 ± 1.68	N/A	482.94 ± 21.76	N/A	40.06 ± 1.42	N/A	15.96 ± 1.46							
1.5x	N/A	5.98 ± 0.12	N/A	35.69 ± 1.40	N/A	43.23 ± 2.15	N/A	491.35 ± 35.67	N/A	40.00 ± 2.15	N/A	14.96 ± 1.20							
3.0x	N/A	6.03 ± 0.14	N/A	35.08 ± 1.39	N/A	40.48 ± 1.72	N/A	484.50 ± 36.17	N/A	37.94 ± 1.87	N/A	13.46 ± 0.87							
6.0x	N/A	5.97 ± 0.14	N/A	35.71 ± 1.54	N/A	40.40 ± 2.36	N/A	443.44 ± 21.28	N/A	36.88 ± 1.38	N/A	13.38 ± 1.15							
Sources of variation																			
CRF	*	***	NS	NS	NS	NS	NS	***	NS	**	NS	***							
PD	NS	***	NS	**	NS	***	NS	***	NS	***	NS	*							
CRF*PD	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS							
FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS							
CRF*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS							
PD*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS							
CRF*PD*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS							

^zIn 2018-19, CRF 1 = 12N-1.31P-7.47K with higher micronutrient content; and CRF 2 = 16N-1.31P-16.6K. In 2019-20, CRF 1 = 12N-1.31P-11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N-1.31P-11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05$, 0.01, or 0.001, respectively. N/A = data not available.

managing Citrus Huanglongbing epidemics? Trop. Plant Pathol. 44:268-274.

Morgan, J.B. and E.L. Connolly. 2013. Plant-soil interactions: nutrient uptake. Nature Educ. Knowledge 4:2. 20 Apr. 2020. <<https://www.nature.com/scitable/knowledge/library/plant-soil-interactions-nutrient-uptake-105289112/>>.

Morgan, K.T. and D.M. Kadyampakeni. 2020. Nutrition of florida citrus trees. 3rd ed. SL 253. Gainesville: Univ. of Florida IFAS. 5 May 2020. <<http://edis.ifas.ufl.edu/pdffiles/SS/SS47800.pdf>>.

Morgan, K.T., R.E. Rouse, and R.C. Ebel. 2016. Foliar applications of essential nutrients on growth and yield of 'Valencia' sweet orange infected with Huanglongbing. HortScience 51:1482-1493.

Nasir, M., A.S. Khan, S.A. Basra, and A.U. Malik. 2016. Foliar application of moringa leaf extract, potassium and zinc influence yield and fruit quality of 'Kinnow' mandarin. Scientia Hort. 210:227-235.

Nwugo, C.C., H. Lin, Y. Duan, and E.L. Civerolo. 2013. The effect of 'Candidatus Liberibacter asiaticus' infection on the proteomic profiles and nutritional status of pre-symptomatic and symptomatic grapefruit (*Citrus paradisi*) plants. BMC Plant Biol. 13:59.

Obreza, T.A. and R.E. Rouse. 1993. Fertilizer effects on early growth and yield of Hamlin orange trees. HortScience 28:111-114.

Phuyal, D., T.A.R. Nogueira, A.D. Jani, D.M. Kadyampakeni, K.T. Morgan, and R.S. Ferrarezi. 2020. 'Ray Ruby' grapefruit affected by Huanglongbing I. Planting density and soil nutrient management. HortScience, doi: 10.21273/HORTSCI115111-20.

Pustaka, A.B., S. Subandiyah, P. Holford, G.A.C. Beattie, T. Iwanami, and Y. Masaoka. 2008. Interactions between plant nutrition and symptom expression in mandarin trees infected with the disease Huanglongbing. Australas. Plant Dis. Notes 3:112-115.

Roka, F., R. Muraro, R.A. Morris, P. Spyke, K. Morgan, A. Schumann, W. Castle, and E. Stover. 2009. Citrus production systems to survive greening: Economic thresholds. Proc. Florida State Hort. Soc. 122:122-126.

Rouse, R.E., M. Ozores-Hampton, F.M. Roka, and P. Roberts. 2017. Rehabilitation of Huanglongbing-affected citrus trees using severe pruning and enhanced foliar nutritional treatments. HortScience 52:972-978.

Shen, W., J.M. Cevallos-Cevallos, U. Nunes da Rocha, H.A. Arevalo, P.A. Stansly, P.D. Roberts, and A.H.C. van Bruggen. 2013. Relation between plant nutrition, hormones, insecticide applications, bacterial endophytes, and *Candidatus Liberibacter Ct* values in citrus trees infected with Huanglongbing. Eur. J. Plant Pathol. 137:727-742.

Shin, K. and A.H.C. van Bruggen. 2018. Bradyrhizobium isolated from Huanglongbing (HLB) affected citrus trees reacts positively with primers for *Candidatus Liberibacter asiaticus*. Eur. J. Plant Pathol. 151:291-306.

Singerman, A., M. Burani-Arouca, and S.H. Futch. 2018. The profitability of new citrus plantings in Florida in the era of Huanglongbing. HortScience 53:1655-1663.

Stansly, P.A., H.A. Arevalo, J.A. Qureshi, M.M. Jones, K. Hendricks, P.D. Roberts, and F.M. Roka. 2014. Vector control and foliar nutrition to maintain economic sustainability of bearing citrus in Florida groves affected by Huanglongbing. Pest Mgt. Sci. 70:415-426.

Stover, E., W.S. Castle, and P. Spyke. 2008. The citrus grove of the future and its implications

Table 11. Soil micronutrient concentration as a function of controlled-release fertilizer (CRF), planting density (PD), and foliar-applied micronutrients (FAM).

Treatment	B (mg·kg ⁻¹)		Mn (mg·kg ⁻¹)		Zn (mg·kg ⁻¹)		Cu (mg·kg ⁻¹)		Fe (mg·kg ⁻¹)	
	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20	2018-19	2019-20
CRF ^z										
CRF 1	0.19 ± 0.01 b ^w	0.39 ± 0.01 b	4.96 ± 0.42 a	16.48 ± 0.87	9.81 ± 0.61	15.11 ± 0.66 b	13.52 ± 0.56	29.22 ± 0.85	8.33 ± 0.74	81.56 ± 3.80
CRF 2	0.38 ± 0.07 a	0.49 ± 0.02 a	2.42 ± 0.19 b	18.83 ± 1.30	9.78 ± 0.91	17.97 ± 0.95 a	14.07 ± 0.70	30.56 ± 0.97	8.83 ± 0.84	83.58 ± 2.89
PD ^y										
SR/LD	0.31 ± 0.11	0.35 ± 0.02 b	4.13 ± 0.81	16.13 ± 1.46	10.09 ± 1.16	14.59 ± 0.84 b	13.95 ± 0.83	9.19 ± 0.85	79.36 ± 3.88 b	9.19 ± 0.85
SR/HD	0.25 ± 0.02	0.50 ± 0.03 a	3.38 ± 0.43	19.17 ± 1.43	9.31 ± 0.93	18.07 ± 1.25 a	13.74 ± 0.48	9.75 ± 0.89	95.28 ± 3.61 a	9.75 ± 0.89
DR/HD	0.29 ± 0.05	0.47 ± 0.02 a	3.56 ± 0.55	17.67 ± 1.16	9.99 ± 0.76	16.97 ± 0.86 ab	13.69 ± 1.00	6.81 ± 0.88	73.08 ± 3.93 b	6.81 ± 0.88
FAM ^x										
0.0x	N/A	0.44 ± 0.02	N/A	14.85 ± 1.07 b	N/A	14.31 ± 0.88 b	N/A	30.46 ± 1.00	N/A	86.13 ± 5.18
1.5x	N/A	0.46 ± 0.03	N/A	17.27 ± 1.38 ab	N/A	16.44 ± 1.27 ab	N/A	31.11 ± 1.55	N/A	83.52 ± 4.84
3.0x	N/A	0.43 ± 0.03	N/A	17.29 ± 1.61 ab	N/A	16.58 ± 1.01 ab	N/A	29.83 ± 1.45	N/A	79.02 ± 3.85
6.0x	N/A	0.44 ± 0.04	N/A	21.21 ± 1.91 a	N/A	18.85 ± 1.39 a	N/A	28.16 ± 1.08	N/A	81.63 ± 5.20
Sources of variation										
CRF	*	***	**	NS	NS	*	NS	NS	NS	NS
PD	NS	**	NS	NS	NS	*	NS	NS	NS	***
CRF*PD	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
FAM	N/A	NS	N/A	*	N/A	*	N/A	NS	N/A	NS
CRF*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS
PD*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS
CRF*PD*FAM	N/A	NS	N/A	NS	N/A	NS	N/A	NS	N/A	NS

^zIn 2018-19, CRF 1 = 12N-1.31P-7.47K with higher micronutrient content; and CRF 2 = 16N-1.31P-16.6K. In 2019-20, CRF 1 = 12N-1.31P-11.62K with 2 times Mg and 2.5 times micronutrients; and CRF 2 = 12N-1.31P-11.62K.

^ySR/LD = single-row low-density (300 trees per ha). SR/HD = single-row high-density (440 trees per ha). DR/HD = double-row high-density (975 trees per ha).

^xFoliar applied micronutrients using the recommended rates of B, Mn, and Zn per year.

^wMean ± SE followed by different lowercase letters are significantly different at $P < 0.05$ by Tukey's test.

NS, *, **, ***Nonsignificant or significant at $P < 0.05$, 0.01, or 0.001, respectively. N/A = value not available.

for Huanglongbing management. Proc. Florida State Hort. Soc. 121:155-159.

Sumner, M.E., P.M.W. Farina, and V.J. Hurst. 1978. Magnesium fixation—a possible cause of negative yield responses to lime applications. Commun. Soil Sci. Plant Anal. 9:995-1007.

Tang, N., Y. Li, and L.S. Chen. 2012. Magnesium deficiency-induced impairment of photosynthesis in leaves of fruiting Citrus reticulata trees accompanied by up-regulation of antioxidant metabolism to avoid photo-oxidative damage. J. Plant Nutr. 175:784-793.

Tansey, J.A., P. Vanaeloch, C. Monzo, M. Jones, and P.A. Stansly. 2017. Costs and benefits of insecticide and foliar nutrient applications to Huanglongbing-infected citrus trees. Pest Mgt. Sci. 73:904-916.

Truog, E.M.I.L. 1947. Soil reaction influence on availability of plant nutrients. Soil Sci. Soc. America J. 11:305-308.

U.S. Department of Agriculture. 2012. United States standards for grades of grapefruit Juice. U.S. Dept. Agr., Washington, DC. 20 June 2020. <https://www.ams.usda.gov/sites/default/files/media/Canned_Grapefruit_Juice_Standard%5B1%5D.pdf>.

Yang, G.H., L.T. Yang, H.X. Jiang, Y. Li, P. Wang, and L.S. Chen. 2012. Physiological impacts of magnesium-deficiency in citrus seedlings: Photosynthesis, antioxidant system and carbohydrates. Trees 26:1237-1250.

Zekri, M. and T. Obreza. 2018. Micronutrient deficiencies in citrus: Iron, zinc, and manganese. SL 204. Univ. of Florida IFAS, Gainesville. 20 Apr. 2020. <<https://edis.ifas.ufl.edu/pdffiles/SS/SS42300.pdf>>.

Zekri, M. and T. Obreza. 2019. Magnesium (Mg) for citrus trees. SL 380. Univ. of Florida IFAS, Gainesville. 20 Apr. 2020. <<https://edis.ifas.ufl.edu/pdffiles/SS/SS58200.pdf>>.

Zhang, M.Q., Y. Guo, C.A. Powell, M.S. Doud, C.Y. Yang, H. Zhou, and Y.P. Duan. 2016. Zinc treatment increases the titer of 'Candidatus Liberibacter asiaticus' in Huanglongbing-affected citrus plants while affecting the bacterial microbiomes. J. Appl. Microbiol. 120:1616-1628.

Zhu, D., W. Juan, S. Liao, and W. Liu. 2007. Relationship between plant availability of boron and the physico-chemical properties of boron in soils, p. 345-354. In: F. Xu, H.E. Goldbach, P.H. Brown, et al. Advances in plant and animal boron nutrition. Springer, Dordrecht, The Netherlands.