

financial support, often rotating the support among several sources. Often, the funding source is permitted to place a small selected educational advertisement in the newsletter.

One of the concerns expressed over the involvement of industry in the production of educational materials is the neutrality of the resulting information and presentation of that material. We have not faced any conflict-of-interest issues pertaining to the presentation of educational materials in an unbiased fashion. We have allowed appropriate and educational commercial advertising in the industry-sponsored publications.

Extension has a large responsibility to provide education to our stakeholders, however, the costs, the need for high quality, and the need for short turnaround time of projects are challenging for the extension service. We must continually search for innovative methods to fund and carry out our extension programs. We have found considerable success in Florida with partnering with commodity organizations and allied industries and their associations. Other states are beginning to use this approach in covering the costs of their extension publications. In the southeastern U.S. (North Carolina, South Carolina, Alabama, Mississippi, and Georgia) state's commercial vegetable recommendations will be produced with industry support.

### Literature cited

- Lamb, E. and G. McAvoy (eds.). 2000. Florida agricultural conference and trade show proceedings. Fla. Coop. Ext. Serv. PRO 516.
- Maynard, D.N. and S.M. Olson (eds.). 2000. Vegetable production guide for Florida. Fla. Coop. Ext. Serv. SP 170.
- Pernezny, K. and T. Kucharek. 1998. Scouting diseases of tomatoes and peppers. Fla. Coop. Ext. Serv. Circ. 1225.
- Vavrina, C.S. (ed.). 2000. Florida tomato institute proceedings. Fla. Coop. Ext. Serv. PRO 517.

## A Survey of Professional Turfgrass Managers in Southern California Concerning Their Use of Turfgrass Best Management Practices

Grant J. Klein<sup>1</sup> and Robert L. Green

**ADDITIONAL INDEX WORDS.** survey, best management practices, turfgrass industry

**SUMMARY.** Turfgrass management best management practices (BMPs) encompass a wide variety of activities, including fertilization, irrigation, mowing, pest control, and soil management. Little attention is given to determining just how effective information regarding BMPs is being assimilated and used by professional turfgrass managers. The objectives of this study were to assess the current perception and implementation of selected turfgrass BMPs and to determine whether or not those perceptions and implementations differed 1) between turfgrass advisors and managers and 2) between general and sports turfgrass managers. Professionals from the turfgrass industry, with an average of 13 years of experience and largely comprised of decision-makers (88%), were surveyed at the University of California, Riverside, Turfgrass Research Conference and Field Day in Fall 1998 and 1999. Turfgrass managers, especially sports turfgrass managers, were found to be the most committed to implementing the BMPs in the survey. Overall, survey respondents considered BMPs to be important and not highly difficult to implement. Limitations to the adoption of BMPs were a lack of financial backing, employee education, and necessary time—all of which could be remedied with a sufficient commitment of resources by the turfgrass industry.

There are numerous definitions of the phrase best management practice, which vary depending on the specific context involved, as well as the currently accepted standards and goals of agronomic management. In general, BMPs are considered to be a set of guidelines or procedures which have been determined, as part of an overall program, to be an effective and practical (technically, socially and economically) method for reducing, preventing, or controlling undesirable effects of management; promoting or maintaining beneficial effects of management; and/or protecting the environment or natural habitat (Campbell, 1996; Hubbard et al., 1998; King County, Wash. Dept. of Transportation, 2000; Lindsey et al., 1998; Logan, 1990; Ohio State Univ. Ext., 1996; One Plan, 2001; South Carolina Forestry Commission, 2001; U.S. Environmental Protection Agency, 2000; Yergert et al., 1993).

Turfgrass-related BMPs encompass a wide variety of activities, including fertilization, irrigation, mowing, pest control, and soil management. There is a great deal of

Department of Botany and Plant Sciences, University of California, Riverside, CA 92521-0124.

<sup>1</sup>To whom request for reprints should be addressed; e-mail grant.klein@ucr.edu.

literature concerning proper implementation strategies or evaluating the efficacy of specific BMPs. However, little attention has been given to determining just how effective information regarding BMPs is being assimilated and used by professional turfgrass managers.

BMP information needs to be made readily available to those making turfgrass management decisions. This function is often served by educational outreach and extension programs that disseminate and promote BMPs. A logical starting point for such programs is to determine what BMPs turfgrass managers (and their advisors) are aware of, and which BMPs,

if any, are currently being implemented. The objectives of this study were to assess the current perception and implementation of selected turfgrass BMPs and to determine whether or not those perceptions and implementations differed 1) between turfgrass advisors and managers and 2) between general and sports turfgrass managers.

## Materials and methods

Professionals from the turfgrass industry were surveyed at the University of California, Riverside, Turfgrass Research Conference and Field Day in Fall 1998 and 1999. These

**Table 1. Survey questions given at the University of California, Riverside, Turfgrass Research Conference and Field Day in Fall 1998 and 1999.**

### SECTION 1

1. Did you complete this survey in 1998?

*Given in 1999 survey only. Choices: yes; no; or not sure.*

2a. Which of the following indicates how you are primarily involved with turfgrass (choose only one)?

*Choices: golf course management; professional consulting/horticulture advising; manufacture/sales of turfgrass-related products; public (government/public property) site management; private (commercial/residential) site management; sports turfgrass management; sod production; turfgrass seed production; turfgrass research; or other.*

2b. Number of years you have been involved with the activity checked above?

2c. What is the county and state where you primarily perform this activity (please specify one county)?

3. How often are you responsible for making decisions or recommendations about turfgrass management practices (turfgrass selection, irrigation practices and system maintenance, fertilization programs, mowing, pesticide applications, etc.)?

*Choices: always; usually; rarely; or never.*

4. If you manage turfgrass, please indicate the turfgrass species at your site (check all that apply).

*Choices: bermudagrass (Cynodon spp.); tall fescue (Festuca arundinacea); creeping bentgrass (Agrostis palustris); perennial ryegrass (Lolium perenne); kikuyugrass (Pennisetum clandestinum); annual bluegrass (Poa annua); St. augustinegrass (Stenotaphrum secundatum); kentucky bluegrass (Poa pratensis); zoysiagrass (Zoysia spp.); and other(s).*

### SECTION 2

5. Examine each of the best management practices (BMPs) described below and note whether or not you consider it to be an important management practice, whether or not you are currently performing the practice, whether or not you are likely to either continue or initiate this practice in the future, and whether or not you feel the practice is easy or very difficult to carry out.

*Responses for each BMP (see below) given on 1 to 5 scale for the following: importance (1 = not, 5 = very); doing this now? (1 = never, 5 = always); likely to continue/start? (1 = not likely, 5 = very likely); difficulty level (1 = easy, 5 = very hard).*

The BMPs:

a) water conservation [reference evapotranspiration (ET<sub>0</sub>)-based water budgets, seasonal adjustments of irrigation clocks, irrigation system checks, etc.]

b) fertility program development (fertilization based on plant species, type of use, and seasonal and climatic requirements, soil type; use of appropriate fertilizer type, amount, and frequency of application)

c) turfgrass selection [choose species and cultivars that, for example, require less water, possess more tolerance to stress (including pests) or possess other traits that would result in the successful management of turfgrass, etc.]

d) mowing program development (mowing height/frequency based on species/cultivar requirements, plant growth and/or stress, etc.)

e) integrated pest management (managing the most healthy turfgrass as possible via sound agronomic principles as the best prevention to pests, defining threshold pest activity/amount prior to pesticide applications, etc.)

f) protecting ground water and surface water from potential contamination from turfgrass chemicals and fertilizers

g) protecting nontarget plants, animals and humans from the potential toxic effects of turfgrass chemicals

h) protecting native habitats during turfgrass construction and maintenance

6. What factor(s) have limited your ability to adopt BMPs in the past (check all that apply)?

*Choices: cost/financial limitations; availability of BMP information; lack of organization/planning; government regulations; employee skill level; time; BMPs not considered important; no personal authority to implement BMPs; client/owner/public unaware of or disinterested in BMPs; and other(s)*

7. Which of the following fertilization practice(s) do you consistently perform (check all that apply)?

*Given in 1999 survey only. Choices: apply appropriate amount of nitrogen (N) specific for turfgrass species and requirements of turfgrass use; apply different combinations of slow- and fast-release N sources according to seasonal growth and expected rainfall; apply N based on seasonal growth patterns and need; apply phosphorus and potassium relative to annual N applied; conduct soil fertility tests every 1 to 2 years; and avoid fertilizing prior to rain.*

8. Which of the following irrigation practice(s) do you consistently perform (check all that apply)?

*Given in 1999 survey only. Choices: irrigate according to weather station/soil moisture sensor data; check system operating pressures; adjust irrigation clocks at least every 3 months; check irrigation systems for proper function; cycle irrigation on slopes to prevent runoff; and size nozzles for balanced precipitation on rotor systems.*

conferences were attended by professional turfgrass managers, personnel working in the turfgrass industries, educators, and consultants from the southern California region. The participants were asked to complete the surveys during a 10-min period following a short presentation regarding BMPs for managing tall fescue (*Festuca arundinacea*). There were 191 surveys collected from the 1998 conference and 190 surveys collected from the 1999 conference, for a total of 381 surveys. In 1999, the survey asked the participants if they had answered the survey in the previous year. Responses of yes or not sure eliminated 76 surveys, leaving a total of 305 surveys as part of the data set. Although this was a sample of convenience (Guilford, 1978; Iversen and Norpoth, 1976; Kish, 1965; Vogt, 1999), which does require caution in terms of generalizability of the data, we believe the survey respondents represented a cross-section of the decision-makers in the turfgrass industry in southern California. A recent survey conducted by Templeton et al. (2000) on the environmental horticulture industry in California included a number of comparable job categories. The distribution they found was similar to our sample.

The survey instrument consisted of two sections (Table 1): the first section requesting basic information about the conference participants, and the second section gauging their activities and perceptions of eight BMPs.

Most of the survey focused on gauging the respondent's activities and perceptions of eight BMPs, including 1) water conservation; 2) fertility program development; 3) turfgrass selection; 4) mowing program development; 5) integrated pest management (IPM); 6) protecting groundwater and surface water from potential contamination from turfgrass chemicals and fertilizers; 7) protecting nontarget plants, animals and humans from the potential toxic effects of turfgrass chemicals; and 8) protecting native habitats during turfgrass construction and maintenance. The respondents indicated, for each BMP, 1) their perceptions regarding its importance, 2) if it was being currently implemented, 3) if they were likely to continue or to start the practice, and 4) their perceptions regarding its difficulty level. The survey also asked the respondents to identify the factors which have limited their ability to

adopt BMPs and which of six fertilization and six irrigation practices they consistently performed.

There were several types of questions in the survey, including 1) questions which asked for simple, basic information about the respondents themselves, which was then summarized or averaged (questions 1, 2b, and 2c); 2) questions which asked for a single response from a list of possible responses for which the percentages of respondents which indicated each possible answer were determined (questions 2a, 3, 9, and 10); 3) questions which asked survey respondents to check all that apply from a list of possible responses for which the percentages of respondents which indicated each choice were determined (questions 4, 6, 7, and 8); and 4) questions which included a series of BMPs which asked respondents to note, for each BMP, on a 1 to 5 scale its importance (1 = not important, 5 = very important), how often it was being currently implemented (1 = never, 5 = always), if they were likely to continue or start the BMP (1 = not likely, 5 = very likely), and its difficulty level (1 = easy, 5 = hard) (questions 5a-h).

To facilitate statistical analysis by the chi-square statistic, turfgrass industry job classifications were coded into advisors and managers. The managers were further coded into general and sports turfgrass managers (Table 2). Sod and turfgrass seed producers were not included in any of the coding or chi-square analyses because their management practices are substantially different from typical turfgrass managers. Respondents who indicated "other" or who indicated multiple turfgrass industry job classifications were also not included.

The responses to two sets of questions were then analyzed with chi-square tests of independence in a 2 × 2 frequency table for both advisors versus managers and general versus sports turfgrass managers. The first set of questions to be analyzed in this manner were regarding the factors limiting the adoption of BMPs and the implementation of specified fertilization and irrigation practices, with each factor or practice coded as yes (if checked) or no (not checked). The second set of questions were regarding the series of eight BMPs, with the responses from the original 1 to 5 scales being coded into "low to moderate" (1 to 3) and "high" (4 to 5).

**Table 2. Information concerning respondents surveyed over 2 years at the University of California, Riverside, Turfgrass Research Conference and Field Day.**

Turfgrass industry job classification	Respondents		Category <sup>y</sup>		Avg no. years experience	Frequency of decisionmaking <sup>z</sup>	
	(no.)	(%)	A/M	G/S		A/U	R/N
Public (government/public property) site management	124	41	M	G	13	89 <sup>x</sup>	11
Golf course management	59	19	M	S	14	85	15
Private (commercial/residential) site management	24	8	M	G	11	96	4
Manufacture/sales of turfgrass-related products	18	6	A	---	14	83	17
Professional consulting/horticultural advising	14	5	A	---	16	93	7
Sports turfgrass management	13	4	M	S	14	92	8
Sod production	12	4	---	---	8	75	25
Turfgrass research	4	1	A	---	18	100	0
Turfgrass seed production	4	1	---	---	15	75	25
Other	7	2	---	---	10	100	0
Multiple classifications	25	8	---	---	15	84	16
Total <sup>w</sup>	305	100	---	---	13	88	12

<sup>z</sup>A/U = always/usually, R/N = rarely/never.

<sup>y</sup>A/M = advisor/manager, G/S = general/sports turfgrass.

<sup>x</sup>Percentages under always/usually and rarely/never total 100% of respondents for a given turfgrass industry job classification.

<sup>w</sup>Includes one survey respondent that did not answer the question regarding turfgrass industry job identification.

**Table 3. Influence of job and turfgrass management categories on the perception of factors limiting the adoption of best management practices (BMPs) of survey respondents.**

Limiting factor	Job category <sup>z</sup> (% yes) <sup>x</sup>			Turfgrass management category <sup>y</sup> (% yes)			All survey respondents <sup>w</sup> (% yes)
	A	M	$\chi^2$	G	S	$\chi^2$	
Cost/financial limitations	54	59	0.27 <sup>NS</sup>	60	57	0.13 <sup>NS</sup>	58
Employee skill level	25	39	2.02 <sup>NS</sup>	38	40	0.05 <sup>NS</sup>	37
Time	25	36	1.41 <sup>NS</sup>	38	34	0.27 <sup>NS</sup>	35
Availability of BMP information	32	21	1.81 <sup>NS</sup>	21	22	0.02 <sup>NS</sup>	21
Lack of organization/planning	14	22	0.95 <sup>NS</sup>	26	14	3.94 <sup>*</sup>	21
Government regulations	7	19	2.52 <sup>NS</sup>	21	15	0.99 <sup>NS</sup>	18
No personal authority to implement BMPs	14	20	0.50 <sup>NS</sup>	23	14	2.18 <sup>NS</sup>	19
BMPs not considered important	18	6	4.63 <sup>*</sup>	9	2	3.66 <sup>+</sup>	8
Other	11	5	1.27 <sup>NS</sup>	4	9	2.84 <sup>+</sup>	7

<sup>z</sup>Total number of respondents = 234; A = advisor, M = manager.

<sup>y</sup>Total number of respondents = 206; G = general, S = sport.

<sup>x</sup>Respondents could check (yes) for as many of the listed factors limiting the adoption of BMPs as applied to their situation.

<sup>w</sup>Total number of respondents from all job classifications = 275.

<sup>NS</sup>, <sup>+</sup>, <sup>\*</sup> Nonsignificant or significant at  $P \leq 0.10$  or  $0.05$ , respectively, by the chi-square test of independence in a  $2 \times 2$  frequency table.

## Results and discussion

The respondents to the surveys were primarily managers of public (government/public property) sites (41%), golf courses (19%), private (commercial/residential) sites (8%), and manufacturers or sales representatives of turfgrass-related products (6%) (Table 2). The vast majority of the respondents were from southern California (88%) and they had an average of 13 years of experience in the turfgrass industry. The respondents managed a wide variety of turfgrasses, including bermudagrass (*Cynodon* sp., 82%), tall fescue (57%), perennial ryegrass (*Lolium perenne*, 56%), kikuyugrass (*Pennisetum clandestinum*, 40%), annual bluegrass (*Poa annua*, 27%), and creeping bentgrass (*Agrostis palustris*, 25%). Most notably, the

respondents were decision-makers, with a total of 88% indicating they were always or usually responsible for turfgrass management decisions or recommendations at their site. This is precisely the audience that educational outreach and extension programs, which are vital for conveying what is learned from scientific research to the turfgrass industry, will need to reach in order to have an impact on turfgrass management practices.

### Factors limiting adoption of BMPs

The single most common factor which limited the ability of all survey respondents to adopt BMPs was cost or financial limitations (58%) (Table 3). About a third of all respondents also chose employee skill level and time (37%

**Table 4. Influence of job and turfgrass management categories on the reported frequency of consistently performing selected fertilization best management practices (BMPs) by survey respondents.**

Fertilization BMPs	Job category <sup>z</sup> (% yes) <sup>x</sup>			Turfgrass management category <sup>y</sup> (% yes)			All survey respondents <sup>w</sup> (% yes)
	A	M	$\chi^2$	G	S	$\chi^2$	
Apply appropriate amount of nitrogen specific for turfgrass species and requirements of turfgrass use	33	60	0.89 <sup>NS</sup>	46	83	11.95 <sup>***</sup>	61
Apply nitrogen based on seasonal growth patterns and need	33	59	0.81 <sup>NS</sup>	54	69	2.01 <sup>NS</sup>	59
Apply different combinations of slow- and fast-release nitrogen sources according to seasonal growth and expected rainfall	67	52	0.26 <sup>NS</sup>	43	66	4.51 <sup>*</sup>	53
Conduct soil fertility tests every 1 to 2 years	0	36	1.68 <sup>NS</sup>	23	57	10.73 <sup>***</sup>	37
Apply phosphorus and potassium relative to annual nitrogen applied	33	25	0.10 <sup>NS</sup>	20	34	2.44 <sup>NS</sup>	26
Avoid fertilizing before rain	0	11	0.37 <sup>NS</sup>	9	14	0.63 <sup>NS</sup>	12

<sup>z</sup>Total number of respondents = 94 (data available for 1999 survey only); A = advisor, M = manager.

<sup>y</sup>Total number of respondents = 91 (data available for 1999 survey only); G = general, S = sport.

<sup>x</sup>Respondents could check (yes) for as many of the listed fertilization BMPs as applied to their situation.

<sup>w</sup>Total number of respondents from all job classifications = 107 (data available for 1999 survey only).

<sup>NS</sup>, <sup>\*</sup>, <sup>\*\*\*</sup> Nonsignificant or significant at  $P \leq 0.05$  or  $0.001$ , respectively, by the chi-square test of independence in a  $2 \times 2$  frequency table.

and 35%, respectively) as important limitations. Notably only 8% of all respondents indicated that not considering BMPs to be important was a limiting factor for adopting BMPs. However, advisors (18%) were more likely than managers (6%) and general turfgrass managers (9%) more likely than sports turfgrass managers (2%) to indicate this as a limiting factor. Also, lack of organization or planning was more of a limiting factor for general turfgrass managers (26%) than sports turfgrass managers (14%).

**FERTILIZATION PRACTICES.** Most respondents indicated that they consistently apply appropriate amounts of nitrogen specific for turfgrass species and requirements of turfgrass use (61%), apply nitrogen based on seasonal growth patterns and need (59%), and apply different combinations of slow- and fast-release nitrogen sources according to seasonal growth and expected rainfall (53%) (Table 4). More than a third (37%) of all respondents also indicated that they conduct soil fertility tests every 1 to 2 years. The least common practices were applying phosphorus and potassium relative to annual nitrogen applied (26%) and avoiding fertilization prior to rain (12%).

There were no differences between advisors and managers for any of the fertilization practices. However, 23% to 37% more sports turfgrass managers than general turfgrass managers indicated that they were consistently performing several of the practices.

**IRRIGATION PRACTICES.** Most respondents (86%) indicated that they consistently check irrigation systems for proper function (Table 5). About two-thirds of all respondents also indicated that they consistently adjust irrigation clocks at least every 3 months (68%) and size nozzles for balanced precipitation on rotor systems (62%). About half of all respondents consistently cycle irrigation on slopes to prevent runoff (55%) and irrigate according to weather station or soil moisture sensor data (49%). Only 41% of all respondents indicated they consistently check system operating pressures.

There were no differences between advisors and managers for any of the irrigation practices. However, more general turfgrass managers (81%) than sports turfgrass managers (49%) indicated they consistently adjusted irrigation clocks at least every 3 months. Also, 20% more sports turfgrass manag-

ers than general turfgrass managers noted they irrigated according to weather station or soil moisture sensor data.

**Perceptions and commitment to BMPs in terms of different job categories**

Overall, most respondents considered all eight of the BMPs to be highly important (i.e., rating them 4 or 5 on the 1 to 5 scale, with 5 being highest), ranging from 64% for protecting native habitats during maintenance and construction to 83% for fertility program development (Table 6). The respondents also considered the BMPs to not be highly difficult—only 22% to 37% rated the BMPs with a 4 or 5 (on the 1 to 5 scale, with 5 being most difficult). Despite the recognition of the importance of the BMPs and the fact that they are generally not highly difficult to implement, only about half to two-thirds of all respondents were conducting the practices with high frequency (ratings of 4 or 5 on the 1 to 5 scale, with 5 being most frequent), or were highly likely to continue or start a practice (ratings of 4 or 5 on the 1 to 5 scale, with 5 being most likely). For all eight of the BMPs in the survey, there were differences between the responses of advisors and managers and between general and sports turfgrass managers, as noted below.

**WATER CONSERVATION.** More turfgrass managers than advisors considered water conservation to be highly important (83% to 69%, respectively). Managers were also more likely to be implementing water conservation with a high frequency than advisors (60% and 44%, respectively) and to continue or start the practice (73% and 53%, respectively). There were no differences between general and sports turfgrass managers in terms of their perceptions and commitment to water conservation.

**FERTILITY PROGRAM DEVELOPMENT.** There were no differences between advisors and managers in terms of their perceptions and commitment to fertility program development. However, more sports turfgrass managers than general turfgrass managers considered this practice to be highly important (90% to 78%, respectively). Sports turfgrass managers, when compared to general turfgrass managers, were also more frequently implementing (76% to 48%, respectively) and more likely to continue or start the practice (79% to 59%, respectively).

**TURFGRASS SELECTION.** Managers were somewhat more

**Table 5. Influence of job and turfgrass management categories on the reported frequency of consistently performing selected irrigation best management practices (BMPs) by survey respondents.**

Irrigation BMPs	Job category <sup>z</sup> (% yes) <sup>x</sup>			Turfgrass management category <sup>y</sup> (% yes)			All survey respondents <sup>w</sup> (% yes)
	A	M	$\chi^2$	G	S	$\chi^2$	
Check irrigation systems for proper function	67	88	1.13 <sup>NS</sup>	87	89	0.05 <sup>NS</sup>	86
Adjust irrigation clocks at least every 3 months	100	69	1.36 <sup>NS</sup>	81	49	10.67 <sup>***</sup>	68
Size nozzles for balanced precipitation on rotor systems	33	61	0.90 <sup>NS</sup>	67	51	2.07 <sup>NS</sup>	62
Cycle irrigation on slopes to prevent runoff	67	58	0.08 <sup>NS</sup>	61	54	0.41 <sup>NS</sup>	55
Irrigate according to weather station/soil moisture sensor data	33	51	0.34 <sup>NS</sup>	43	63	3.49 <sup>*</sup>	49
Check system operating pressures	33	40	0.06 <sup>NS</sup>	39	43	0.14 <sup>NS</sup>	41

<sup>z</sup>Total number of respondents = 92 (data available for 1999 survey only); A = advisor, M = manager.

<sup>y</sup>Total number of respondents = 89 (data available for 1999 survey only); G = general, S = sport.

<sup>x</sup>Respondents could check (yes) for as many of the listed irrigation BMPs as applied to their situation.

<sup>w</sup>Total number of respondents from all job classifications = 103 (data available for 1999 survey only).

<sup>NS,\*,\*\*\*</sup>Nonsignificant or significant at  $P \leq 0.10$  or 0.001, respectively, by the chi-square test of independence in a 2 x 2 frequency table.

**Table 6. Influence of job and turfgrass management categories on the perception and commitment to eight best management practices (BMPs).**

Perception/commitment to BMPs	Job category <sup>z</sup> (% high) <sup>x</sup>			Turfgrass management category <sup>y</sup> (% high)			All survey respondents <sup>w</sup> (% high)
	A	M	$\chi^2$	G	S	$\chi^2$	
Importance							
Water conservation	69	83	3.53*	84	81	0.35 <sup>NS</sup>	79
Fertility program development	81	82	0.06 <sup>NS</sup>	78	90	4.70*	83
Turfgrass selection	61	75	3.26*	74	79	0.80 <sup>NS</sup>	75
Mowing program development	53	82	15.07***	81	83	0.16 <sup>NS</sup>	79
Integrated pest management (IPM)	58	73	3.09*	70	78	1.38 <sup>NS</sup>	73
Protecting water sources from chemicals and fertilizers	69	77	1.04 <sup>NS</sup>	77	78	0.01 <sup>NS</sup>	75
Protecting nontarget plants, animals and humans from chemicals	69	84	4.16*	84	82	0.22 <sup>NS</sup>	82
Protecting native habitats during construction/maintenance	53	66	2.49 <sup>NS</sup>	61	76	4.82*	64
Frequency of current implementation							
Water conservation	44	60	3.26*	57	67	1.73 <sup>NS</sup>	56
Fertility program development	64	57	0.56 <sup>NS</sup>	48	76	15.98***	61
Turfgrass selection	44	45	0.01 <sup>NS</sup>	42	53	2.31 <sup>NS</sup>	47
Mowing program development	39	67	10.73***	64	74	0.16 <sup>NS</sup>	64
Integrated pest management (IPM)	42	49	0.60 <sup>NS</sup>	41	65	11.86***	49
Protecting water sources from chemicals and fertilizers	56	54	0.03 <sup>NS</sup>	48	67	6.82**	52
Protecting nontarget plants, animals and humans from chemicals	58	74	3.80*	72	78	0.76 <sup>NS</sup>	71
Protecting native habitats during construction/maintenance	36	48	1.68 <sup>NS</sup>	43	57	3.64*	45
Likelihood to continue/start implementation							
Water conservation	53	73	5.85*	74	69	0.58 <sup>NS</sup>	68
Fertility program development	64	66	0.06 <sup>NS</sup>	59	79	8.37**	69
Turfgrass selection	53	55	0.06 <sup>NS</sup>	54	57	0.16 <sup>NS</sup>	55
Mowing program development	39	68	11.12***	66	71	0.47 <sup>NS</sup>	65
Integrated pest management (IPM)	42	58	3.23*	52	69	6.02**	57
Protecting water sources from chemicals and fertilizers	64	62	0.06 <sup>NS</sup>	61	64	0.19 <sup>NS</sup>	62
Protecting nontarget plants, animals and humans from chemicals	58	75	4.06*	74	75	0.01 <sup>NS</sup>	71
Protecting native habitats during construction/maintenance	36	55	4.64*	50	67	5.45*	53
Difficulty level							
Water conservation	33	30	0.12 <sup>NS</sup>	34	24	2.37 <sup>NS</sup>	30
Fertility program development	25	22	0.18 <sup>NS</sup>	22	22	0.01 <sup>NS</sup>	22
Turfgrass selection	39	34	0.31 <sup>NS</sup>	30	43	3.83*	33
Mowing program development	31	23	1.04 <sup>NS</sup>	25	18	1.33 <sup>NS</sup>	24
Integrated pest management (IPM)	39	37	0.06 <sup>NS</sup>	36	38	0.02 <sup>NS</sup>	37
Protecting water sources from chemicals and fertilizers	36	36	0.00 <sup>NS</sup>	39	31	1.33 <sup>NS</sup>	35
Protecting nontarget plants, animals and humans from chemicals	28	30	0.10 <sup>NS</sup>	33	25	1.50 <sup>NS</sup>	30
Protecting native habitats during construction/maintenance	28	31	0.14 <sup>NS</sup>	26	40	4.40*	32

<sup>z</sup>Total number of respondents = 256; A = advisor, M = manager.

<sup>y</sup>Total number of respondents = 220; G = general, S = sport.

<sup>x</sup>Responses originally on a 1 to 5 scale (5 = highest) with 1 to 3 coded as "moderate/low" and 4 to 5 as "high."

<sup>w</sup>Total number of respondents from all job classifications = 305.

<sup>NS</sup>, \*, \*\*, \*\*\* Nonsignificant or significant at  $P \leq 0.10$ , 0.05, 0.01, or 0.001, respectively, by the chi-square test of independence in a  $2 \times 2$  frequency table.

likely than advisors to view turfgrass selection as highly important (75% to 61%, respectively). However, only about half of all respondents were either performing this practice with high frequency (47%) or likely to start or continue this practice (55%). Overall, only 33% of the respondents considered the practice to be highly difficult, although more sports turfgrass managers (43%) than general turfgrass managers (30%) considered it to be highly difficult.

**MOWING PROGRAM DEVELOPMENT.** Only 53% of advisors considered mowing program development to be highly important as compared to 82% of managers. Similarly, managers were conducting this practice with higher frequency and were more likely to continue or start the practice than were advisors (differences of 28% and 29%, respectively). Overall, 79% considered the practice to be highly important, and only 24% considered it to be highly difficult to implement. There were no differences between general and sports turfgrass managers in regards to this BMP.

**INTEGRATED PEST MANAGEMENT.** IPM was viewed by managers to be more highly important than it was by advisors (73% and 58%, respectively), and managers were more highly likely to continue or start the practice than advisors (58% and 42%, respectively). Less than half of both managers and advisors practiced IPM with high frequency (49% and 42%, respectively). Most of the general and sports turfgrass managers considered IPM to be highly important (70% and 78%, respectively), but sports turfgrass managers were more frequently conducting IPM and more highly likely to continue or start IPM than general turfgrass managers (a difference of 24% and 17%, respectively).

**PROTECTING WATER SOURCES FROM CHEMICALS AND FERTILIZERS.** There were no differences between advisors and managers when it came to protecting groundwater and surface water sources from potential contamination from turfgrass chemicals and fertilizers, and only one difference between general and sports turfgrass managers (19% more sports turfgrass managers were more highly likely to be frequently conducting the BMP than general turfgrass managers).

**PROTECTING NONTARGETS FROM CHEMICALS.** Managers were more attentive to protecting nontarget plants, animals and humans from the potential toxic effects of turfgrass chemicals than were advisors. There were 15% more managers than advisors who considered the BMP to be highly important, and 16% to 17% were more highly likely to be conducting this practice or likely to continue or start this practice, respectively. There were no differences between general and sports turfgrass managers in regards to this BMP.

**PROTECTING NATIVE HABITATS DURING CONSTRUCTION AND MAINTENANCE.** The only difference between managers and advisors in regards to protecting native habitats during turfgrass construction and maintenance was that more managers than advisors were highly likely to continue or start the practice (55% and 36%, respectively). However, sports turfgrass managers and general turfgrass managers considered the BMP quite differently. Sports turfgrass managers considered the practice to be more highly important than general turfgrass managers (76% and 61%, respectively) and were more highly likely to be frequently conducting the BMP or highly likely to continue or start the BMP (a difference of 14% and 17%, respectively). Further, 40% of sports turfgrass managers, compared to 26% of general turfgrass managers considered protecting native habitats to be highly difficult to implement.

## Conclusion

Overall, the survey respondents considered BMPs to be important, although turfgrass managers, especially sports turfgrass managers, were the most likely to be committed to the BMPs listed in the survey. The survey respondents also did not generally consider the BMPs to be difficult to implement. Together, this suggests that what is needed is a greater commitment to provide the financial backing, employee education, and necessary time which has previously limited the adoption of important BMPs. Educational outreach and extension programs will play an important role in this effort, and in order to most effectively create such programs, more in-depth and rigorous surveys of the turfgrass industry are necessary.

## Literature cited

- Campbell, C.R. 1996. Great tools for implementing best management practices. In: Proc. 6th Annu. S.E. N.C. Prof. Turfgrass Conf. Field Tour, 11-13 Sept. 1996, Sunset Beach and Raleigh, N.C. N.C. Coop. Ext. Serv. 14 June 2001. <<http://www.agr.state.nc.us/agronomi/crc996.htm>>.
- Guilford, J.P. 1978. *Fundamental statistics in psychology and education*. 4th ed. McGraw Hill, New York.
- Hubbard, W., C. Latt, and A. Long. 1998. Forest terminology for multiple-use management. Univ. Fla. Coop. Ext. Serv. Inst. of Food and Agr. Sci., Fla. Forest Stewardship Program. Fla. Div. For. School of For. Resources and Conservation Publ. SS-FOR-11. 11 Apr. 2001. <<http://www.sfr.ufl.edu/Extension/ssfor11.htm>>.
- Iversen, G.R. and H. Norpoth. 1976. *Analysis of variance*. Sage Univ. Paper series on quantitative application in the social sciences 07-001. Sage Publ., Inc., Beverly Hills, Calif.
- King County, Wash., Department of Transportation. 2000. Regional road maintenance Endangered Species Act program guidelines (final draft), p. 3.1. King County, Wash., Dept. Transportation, Reg. Road Maintenance Tech. Working Group. 11 Apr. 2001. <<http://www.metrokc.gov/roadcon/bmp/pdf/glossary.pdf>>.
- Kish, L. 1965. *Survey sampling*. Wiley, New York.
- Lindsey, B.D., K.J. Breen, M.D. Bilger, and R.A. Brightbill. 1998. Water quality in the lower Susquehanna river basin, Pennsylvania and Maryland, 1992-95. U.S. Geol. Survey Circ. 1168. 11 Apr. 2001. <<http://water.usgs.gov/pubs/circ1168>>.
- Logan, T.J. 1990. Agricultural best management practices and groundwater protection. *J. Soil Water Conservation* 45(2):201-206.
- Ohio State University Extension. 1996. Best management practices. Ohio State Univ. Ext. Bul. 472. 14 June 2001. <<http://www2.ag.ohio-state.edu/~ohioline/b472/manage.html>>.
- One Plan. 2001. Best management practices: A list of components. The One Plan. 11 Apr. 2001. <<http://www.oneplan.org/allBMP.htm>>.
- South Carolina Forestry Commission. 2001. Best management practices. S.C. For. Comm. 11 Apr. 2001. <<http://www.state.sc.us/forest/rbg.htm>>.
- Templeton, S.R., C. Brown, G.E. Goldman, S.J. Yoo, and V.S. Pradham. 2000. An economic analysis of environmental horticulture with a focus on California. *HortScience* 35:987-992.
- U.S. Environmental Protection Agency. 2000. Urban watershed management branch glossary. U.S. EPA Urban Watershed Mgt. Branch. 11 Apr. 2001. <<http://www.epa.gov/ednrmrl/main/abc/b.htm>>.
- Vogt, W.P. 1999. *Dictionary of statistics and methodology: A nontechnical guide for the social sciences*. 2nd ed. Sage Publ., Thousand Oaks, Calif.
- Yergert, M., B. Austin, and R. Waskom. 1993. Best management practices for turfgrass production: Turf BMP fact sheet. Colo. State Univ. Coop. Ext. and Colo. Dept. Agr. 14 June 2001. <[http://www.vilnius.lt/vanduo/dw\\_eng/bmp\\_turfbmp.htm](http://www.vilnius.lt/vanduo/dw_eng/bmp_turfbmp.htm)>.