Performance of Primocane-fruiting Experimental Blackberry Cultivars in the Southern Appalachian Mountains

Gina E. Fernandez¹ and James R. Ballington

ADDITIONAL INDEX WORDS. *Rubus*, synthetic spun-bonded cover, polyester, polypropylene, production systems, season extension

SUMMARY. Yield performance of primocane-fruiting experimental blackberry (*Rubus* spp.) cultivars from two breeding programs were evaluated in the southern Appalachian Mountains. The trial consisted of the nine experimental cultivars: NC 533, NC 534, NC 535, NC 537, NC 538, NC 539, APF-27, APF-46, and Prime-Jim®. In addition, rowcovers were applied to split plots in the spring to determine if harvest dates could be advanced. In 2007, five experimental cultivars had comparable high total yields (APF-27, APF-46, NC 535, NC 537, and NC 539), yet only three of these had comparable marketable yields (APF 27, APF-46, and NC 537). In 2008, APF-27 and APF-46 had significantly higher total yield (2234.5 and 2257.3 g/plant, respectively) and marketable yield (1611.2 and 1654.2 g/plant, respectively) than all other experimental cultivars. Rowcover treatments did not increase or decrease either total or marketable yield in any of the experimental cultivars in either year. In addition, rowcovers did not advance the date of 5% or 50% harvest in either year. On the basis of the conditions imposed in these studies, we determined that “APF” selections evaluated in this trial have higher second-year yield than “NC” selections, primocane-fruiting experimental cultivar yields are lower than yields commonly found with florican-fruiting blackberries, and rowcovers as applied in this study are not a viable option for advancing harvest season of primocane-fruiting blackberries in the southern Appalachian Mountains.

Commercial blackberry production is increasing in the southern and eastern United States (Strik et al., 2007). The majority of the production in this region is based on the use of florican-fruiting cultivars, where fruiting occurs on second-year canes and the harvest season commences in June and ends in mid-August. In 1995, the University of Arkansas and North Carolina State University jointly released NC 194, a primocane-fruiting blackberry selection that was intended for use as parental germplasm (Ballington and Moore, 1995). Ten years later, the University of Arkansas released the first primocane-fruiting

cultivars Prime-Jim® and Prime-Jan® (Clark et al., 2005). Although small in scope, the North Carolina State University blackberry breeding program continued to devote a portion of their resources to the breeding of primocane-fruiting blackberries since the release of NC 194 and have had several lines in observational trials in multiple locations in the state.

In North Carolina, we have observed low productivity when primocane-fruiting experimental cultivars from both breeding programs were grown in observational plots at research stations in the Piedmont region (elevation 300–400 ft). However, when primocane-fruiting experimental blackberry cultivars were grown in Mills River, NC (elevation 2000 ft), fruit

production was much higher; both fruit size and number of fruit were greater than those observed in the Piedmont region. Although fruit production potential was much greater in the higher elevation, fall frosts often damaged fruit and ended the harvest season before the crop could reach its full potential (J.R. Ballington, personal communication). In Clarksville, AR, primocane-blackberries produce only a small amount of fruit in the late summer, likely as a result of lack of heat tolerance of the flowers (Clark, 2008; Stanton et al., 2007). However, when these same experimental cultivars were grown in Aurora, OR, harvest commenced in August and extended into October (Strik et al., 2008).

Season extension techniques are used to advance or delay the fruiting season of many small fruit crops including raspberry (*Rubus idaeus*) and blackberry (Pritts, 2008; Pritts et al., 1992; Strik and Thompson, 2009; Strik et al., 2008). When rowcovers were applied over the rows of primocane-fruiting raspberries in the early spring before primocane emergence, fruit production occurred up to 3 weeks earlier than in rows without rowcovers. Strik et al. (2008) were able to advance bloom by 14 d when rowcovers were placed over dormant plots of primocane-fruiting experimental blackberry cultivars in Oregon. No studies have been conducted to evaluate the efficacy of rowcover use in promoting earlier fruiting in primocane-fruiting blackberries in the high elevations of the southern Appalachian region.

The objectives of this research were 2-fold. First, we wanted to document yield of primocane-fruiting experimental blackberry cultivars from the North Carolina State University and University of Arkansas breeding programs in the southern Appalachian Mountains. Second, we wanted to examine the efficacy of rowcover use in promoting earlier fruiting and higher

Department of Horticultural Science, North Carolina State University, Box 7609, Raleigh, NC 27695

Research support provided in part by funds from the Golden LEAF Foundation.

We thank Tomas Moreno, Absalom Shank, and Larry Wohlers for their technical assistance.

Numbered experimental cultivars used in this study were tested in replicated yield trials from the North Carolina State University and University of Arkansas blackberry breeding programs. These selections were under evaluation at a number of locations to determine merits of each individual experimental cultivar. Therefore, experimental cultivars used in this study may or may not be available to the general public as a public cultivar release in the future.

¹Corresponding author. E-mail: Gina_Fernandez@ncsu.edu.

Units

To convert U.S. to SI, multiply by	U.S. unit	SI unit	To convert SI to U.S., multiply by
0.4047	acre(s)	ha	2.4711
0.3048	ft	m	3.2808
2.54	inch(es)	cm	0.3937
1.1209	lb/acre	kg·ha ⁻¹	0.8922
28.3495	oz	g	0.0353
33.9057	oz/yd ²	g·m ⁻²	0.0295
(°F – 32) ÷ 1.8	°F	°C	(1.8 × °C) + 32

yields in nine primocane-fruiting experimental blackberry cultivars.

Materials and methods

This study was conducted at the Upper Mountain Research Center in Laurel Springs, NC (lat. 36°23'60"N, long. 86°17'24"W, elevation 867 m; average last/first freeze date 19 Apr./15 Oct.). The soil was a Toxaway loam. The plants received standard recommended cultural practices for blackberries in this region, with the addition of rowcovers placed on plots in the spring as described below (Fernandez and Krewer, 2008). The plantings consisted of nine primocane-fruiting experimental cultivars (NC 533, NC 534, NC 535, NC 537, NC 538, NC 539, APF-27, APF-46) and the cultivar Prime-Jim®. APF-27, APF-46, and 'Prime-Jim®' were provided courtesy of the University of Arkansas. Plants were set in the field on 24 May 2006 into raised beds that were covered with black plastic. Plants were spaced at 4 ft between plants and 10 ft between rows. Plants were allowed to grow in the first year with minimal training and were pruned to the ground in December in all years of the study.

The experiment was a randomized complete block design with split plots. The experimental cultivar was the main plot, and presence or absence of rowcover (+/–RC) was the split plot. Each split plot consisted of two plants. The treatment combinations were replicated four times. Plots were 20 ft long, and each split plot was 10 ft long. Spunbound polypropylene rowcovers [1.2 oz/yard² (Typar®; Reemay, Old Hickory, TN)] were placed on the designated split plot on 3 Mar. 2007 and 1 Apr. in 2008, before primocane emergence. The rowcovers remained on the plots for 56 d in 2007 and 42 d in 2008. At this time, newly emerging canes were at least 6 inches tall. Cane heights were recorded on three dates in 2007. Plants were soft-tipped (removal of 0.75–2 inches) in all plots when the majority of canes in a plot reached 1 m. Soft-tipping occurred on a weekly basis for ≈3 weeks in July depending on experimental cultivar; however, dates were not recorded.

Harvested fruit from each plot was separated into marketable and non-marketable berries. Marketable berries were blemish free, weighed 5 g or greater, and did not have any visible deformities such as white drupelets.

Individual fruit weight was determined using berries randomly sampled before separating fruit into marketable and non-marketable berries. Determination of the effects of experimental cultivar and rowcover on yield parameters was analyzed statistically as a split-plot design with four replications using appropriate analysis of variance procedures (SAS version 9.1; SAS Institute, Cary, NC). Means were separated using Fisher's protected least significant difference at $P \leq 0.05$.

Results and discussion

Year, experimental cultivar, and the interaction between the two variables had significant effect on total yield, marketable yield, and fruit weight. Therefore, data for the 2 years (consecutive) could not be combined (Table 1). Total and marketable yield was lower in most experimental cultivars in 2007 than in 2008 (Table 2). This year-to-year difference could be due to heat stress or age of the planting. Previous research has shown that heat stress could result in this type of variability. Stanton et al. (2007) found that viability of floral parts (pollen germination and stigma receptivity) in primocane-fruiting blackberry cultivars was reduced when temperatures exceeded 84 °F. Bloom commenced on 12 June 2007 and 24 June 2008 and lasted ≈3 weeks for all of the experimental cultivars. The maximum daily temperatures during bloom were 86 to 87 °F in June and July of both years (State Climate Office of North Carolina, 2010). Therefore, if damage from heat stress to flowers occurred during bloom, it

likely occurred each year and had comparable negative effects on yield.

Although temperatures during bloom could have reduced yield, the age of the planting was more likely the major factor in lower yield in 2007. The plots in this study were established in 2006 and were harvested for the first time in 2007. We have found that first-year yields from our florican-fruiting blackberry trials are lower than those in subsequent years (Fernandez and Shank, 2010). Although we have not systematically harvested primocane-fruiting blackberries in North Carolina before this study, we believe that yield in all primocane-fruiting production systems will be lower in the first year of harvest than in subsequent years.

In both years of this study, APF-27 and APF-46 had the highest total yield among all the experimental cultivars evaluated in this trial (Table 2). Yields of the "NC" experimental cultivars were not consistent from year to year. Total yield of NC 537 and NC 539 were comparable to APF-27 and APF-46 in 2007. However, in 2008, none of the "NC" selections had yields comparable to the two "APF" selections. Total yields of APF-27 and APF-46 in 2008 were 2234.5 and 2257.3 g/plant, respectively. Therefore, based on a plant density recommended for florican-fruiting cultivars (Fernandez and Krewer, 2008) of 1090 plants/acre, the potential total yield of APF-27 and APF-46 would be ≈5622 and 5413 lb/acre, respectively. This yield is lower than 'Navaho', a florican-fruiting blackberry that has produced at least 8000 g/plant (Fernandez and

Table 1. Probability values from analysis of variance for year, experimental cultivar, with and without rowcovers, and interaction of total yield, marketable yield, and fruit size of nine primocane-fruiting experimental blackberry cultivars grown in 2 years (consecutive) in Laurel Springs, NC.

Source	Probability		
	Total yield	Marketable yield	Mean fruit wt
Year	0.0001	0.0001	0.052
Replication (year)	0.5422	0.4895	0.0255
Experimental cultivar	0.0001	0.0001	0.0001
Experimental cultivar × year	0.0001	0.0001	0.0024
Replication × experimental cultivar (year)	0.9993	0.996	0.8777
Rowcover	0.538	0.489	0.7464
Rowcover × experimental cultivar	0.4776	0.4477	0.2094
Rowcover × year	0.7239	0.796	0.9228
Rowcover × experimental cultivar × year	0.3552	0.2058	0.0589

Table 2. Total yield, marketable yield, individual fruit weight, and 5% and 50% harvest dates of nine primocane-fruited experimental blackberry cultivars in 2007 and 2008.

Experimental cultivar	Total yield (g/plant) ^a		Marketable yield (g/plant)		Mean fruit wt (g)		Date of 5% harvest		Date of 50% harvest	
	2007	2008	2007	2008	2007	2008	2007	2008	2007	2008
APF 27	997.3 a ^y	2344.5 a	859.8 a	1611.2 a	7.6 a	7.9 a	7 Sept. a	28 Aug. ab	16 Sept. a	10 Sept. ab
APF 46	827.5 a	2257.3 a	667.0 ab	1654.2 a	6.1 cd	6.1 c	7 Sept. ab	28 Aug. a	19 Sept. ab	12 Sept. a
NC 533	440.0 bc	1329.0 bc	298.7 cd	690.6 b	4.1 d	4.9 de	7 Sept. a	31 Aug. abc	18 Sept. ab	13 Sept. abc
NC 534	70.5 d	441.0 d	59.1 e	262.0 c	3.1 d	5.2 de	7 Sept. a	7 Sept. d	18 Sept. ab	17 Sept. d
NC 535	751.2 ab	1513.7 b	495.3 bc	741.3 b	7.0 ab	7.1 b	7 Sept. ab	4 Sept. cd	21 Sept. bc	18 Sept. cd
NC 537	902.2 a	910.2 cd	619.5 ab	517.9 bc	5.7 c	5.9 c	10 Sept. ab	6 Sept. d	24 Sept. cd	19 Sept. d
NC 538	175.2 cd	981.5 bcd	100.8 de	540.3 bc	3.7 d	4.7 e	18 Sept. de	2 Sept. bcd	30 Sept. e	19 Sept. bcd
NC 539	822.7 a	446.1 d	501.0 bc	237.7 c	6.1 bc	5.3 d	12 Sept. bc	13 Sept. e	25 Sept. d	22 Sept. e
Prime-Jim	319.2 cd	900.9 cd	212.2 de	563.1 bc	7.6 a	6.8 b	13 Sept. cd	3 Sept. cd	29 Sept. e	25 Sept. cd

^a1 g = 0.0353 oz.

^yMean separation in columns by Fisher's protected least significance difference test at $P \leq 0.05$.

Shank, 2010). The highest yielding experimental cultivars from the North Carolina State University breeding program in 2008 were NC 533 (3187 lb/acre) and NC 535 (3629 lb/acre), which had total yields higher than 'Prime-Jim®' (2160 lb/acre).

Marketable yields for most of the experimental cultivars were much lower than total yields (Table 2). The majority of the culled fruit was considered unmarketable (too soft to ship) and therefore unsuitable for commercial production. Strik and Thompson (2009) report that quality of fruit harvested from the florican crop of 'Prime-Jim®' and 'Prime-Jan®' is lower than that of fruit harvested from other florican-fruited cultivars. Although no formal postharvest assessments of the "NC" experimental cultivars were conducted, notes from harvest indicate that the majority of the culled fruit was soft and therefore not marketable.

Fruit weight is an important component of yield potential in blackberries (Clark et al., 2007). Although much of the fruit was soft, the majority of the experimental cultivars had average fruit weight above 5 g, which was considered a marketable size in this trial (Table 2). The experimental cultivars APF-27 and NC 535 had an average fruit weight of 7 g or more for both years of this study (Table 2). This fruit weight is comparable to the newly released 'Prime-Ark® 45' from the University of Arkansas (Clark, 2010).

Initial harvest dates changed in each year of this study; however, the order in which we started to harvest each experimental cultivar was, in general, consistent (Table 2). Harvest commenced in the first week of Sept. 2007

and the last week of Aug. 2008. APF-27, APF-46, NC 533, and NC 534 were the earliest experimental cultivars and had comparable dates of 5% and 50% harvest in 2007. Only the former three experimental cultivars had comparable harvest dates in 2008.

The placement of rowcovers on plots of primocane-fruited blackberries at Laurel Springs, NC, in the spring did not increase total or marketable yield or fruit weight or advance the dates of 5% and 50% harvest (Table 3). In 2007, average cane height in plots with rowcovers was greater than for plots with no rowcovers on all dates before tipping (Fig. 1). Once the primocanes reached 1 m, they were soft-tipped. Strik et al. (2008) found that canes that were tipped at 1 m produced higher yields than canes that were not tipped. The higher yields were attributed to the higher number of branches on each main cane. However, we did not experience the same increase in yield. In this study, all canes were tipped at the same height so that tipping occurred over 3 weeks starting in mid-July during the two growing seasons. Drake and Clark (2003) found that soft-tipping late in the season reduced yield of primocane-fruited plants. Because canes in our study were tipped at the same height over 3 weeks in mid-summer, the tipping process may have negated any advantage in earliness and we may have experienced the same late season effect as experienced by Drake and Clark (2003).

This study was carried out at a site in the southern Appalachian Mountains, where the frost-free season is ≈ 144 d (State Climate Office of North Carolina, 2010). Temperatures of 31

and 28 °F were recorded in the plots on 1 Oct. 2007 and 3 Oct. 2008, respectively. Although these temperatures were slightly below freezing temperature, the fruit remaining on the plants was severely damaged and had to be culled. A combination of high tunnels and rowcovers or heat blankets could help to extend the harvest season in the future. Both et al. (2007) were able to maintain temperatures 1–2 °F above outside temperatures in high tunnels using heat blankets. Alternatively, growers may be able to avoid early frosts by planting in a location that may have a slightly longer growing season. In the future, the new U.S. Department of Agriculture Plant Hardiness Zone Map may enable growers to decide where to plant a frost-sensitive crop. The new maps, which are based on geographic information system data, will be able to elucidate the effects of elevation increases and enclosure by mountainous terrain on a finer geographic scale (K. Kaplan, personal communication).

These are the first data collected on yield and season extension of primocane-fruited experimental blackberry cultivars in the southern Appalachian Mountains. Primocane-fruited experimental cultivars from North Carolina State University had yields that were comparable to the first generation of release from the University of Arkansas, such as 'Prime-Jim®'. The newer experimental cultivars, APF-27 and APF-46, had the highest total and marketable yields of all the materials evaluated in this trial. However, none of the experimental cultivars had yields comparable to those of florican-fruited cultivars. Previous reports showed that

Table 3. Total yield, marketable yield, individual fruit weight, and 5% and 50% harvest dates of nine primocane-fruited experimental blackberry cultivars with (+RC) and without (–RC) rowcovers in 2007 and 2008.

	Total yield (g/plant) ^a				Marketable yield (g/plant)				Mean fruit wt (g)		Date of 5% harvest				Date of 50% harvest			
	2007		2008		2007		2008		2007		2007		2008		2007		2008	
	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC	+RC	–RC
Treatment mean ^y	610.1	555.0	1214.1	1144.4	438.3	387.8	734.1	688.1	5.6	5.6	5.9	6.0	10 Sept.	8 Sept.	21 Sept.	24 Sept.	9 Sept.	9 Sept.

^a1 g = 0.0353 oz.

^yTreatment means (+RC and –RC) within each year were not significantly different with Fisher's protected least significant difference test at $P \leq 0.05$.

Literature cited

- Ballington, J.R. and J.N. Moore. 1995. NC 194 primocane fruiting thorny erect tetraploid blackberry germplasm. *Fruit Var. J.* 49:101–102.
- Both, A.J., E. Reiss, J.F. Sudal, K.E. Holmstrom, C.A. Wyenandt, W.L. Kline, and S.A. Garrison. 2007. Evaluation of a manual energy curtain for tomato production in high tunnels. *HortTechnology* 17:467–472.
- Clark, J.R. 2008. Primocane-fruited blackberry breeding. *HortScience* 43: 1636–1639.
- Clark, J.R. 2010. Prime-Ark 45®, a new blackberry variety. *Small Fruit News*. 22 Sept. 2010. <<http://www.smallfruits.org/Newsletter/Vol10-Issue1.pdf>>.
- Clark, J.R., J.N. Moore, J. Lopez-Medina, C. Finn, and P. Perkins-Veazie. 2005.

'Prime-Jan®' and 'Prime-Jim®' primocane-fruited blackberries. *HortScience* 40:852–855.

Clark, J.R., E.T. Stafne, H.K. Hall, and C.E. Finn. 2007. Blackberry breeding and genetics. *Plant Breed. Rev.* 29:19–144.

Drake, C.A. and J.R. Clark. 2003. Effects of pruning and cropping on field-grown primocane fruiting blackberries. *HortScience* 38:260–262.

Fernandez, G.E. and G. Krewer. 2008. 2008 Southeast Regional Bramble Production Guide. 22 Sept. 2010. <<http://www.smallfruits.org/SmallFruitsRegGuide/Guides/2008/08BrambleguideMay22.pdf>>.

Fernandez, G. and A. Shank. 2010. Blackberry and raspberry harvest data 2008 and 2009. 22 Sept. 2010. <<http://ncsu.edu/enterprises/blackberries-raspberries/production/latest-research/>>.

Pritts, M., E. Hanson, J. Fiola, and M. Kelly. 1992. Rowcovers accelerate fruiting and increase productivity in primocane-fruited red raspberries. *HortTechnology* 2:46–51.

Pritts, M. 2008. Primocane-fruited raspberry production. *HortScience* 43:1640–1641.

Stanton, M.M., J.C. Scheerens, R.C. Funt, and J.R. Clark. 2007. Floral competence of primocane-fruited blackberries Prime-Jan® and Prime-Jim® grown at three temperature regimes. *HortScience* 42:508–513.

State Climate Office of North Carolina. 2010. NC CRONOS Database, Laurel Springs (NLSP). 6 Aug. 2010. <<http://>>

Fig. 1. Average height of primocanes of nine primocane-fruited experimental blackberry cultivars on three dates in 2007 in Laurel Springs, NC. Cane heights were measured before the date when all canes were tipped to promote branching. Plots with rowcovers (●) and without rowcovers (◆). Each point is mean $\pm$ SE of the nine experimental cultivars; 1 cm = 0.3937 inch.

www.nc-climate.ncsu.edu/cronos/?station=NLSP.

Strik, B.C., J.R. Clark, C.E. Finn, and M.P. Banados. 2007. Worldwide blackberry production. *HortTechnology* 17: 205–213.

Strik, B., J.R. Clark, C. Finn, and G. Buller. 2008. Management of primo-

cane-fruited blackberry to maximize yield and extend the fruiting season. *Acta Hort.* 777:423–428.

Strik, B.C. and E. Thompson. 2009. Primocane-fruited blackberries: Potential for extending harvest season and production regions. *HortScience* 44:23–24.

Thompson, E., B.C. Strik, C.E. Finn, Y. Zhao, and J.R. Clark. 2009. High tunnel versus open field: Management of primocane-fruited blackberry using pruning and tipping to increase yield and extend the fruiting season. *HortScience* 44: 1581–1587.