Orientation of Small Hobby High Tunnels and Potential Effects on Cut Sunflowers and Fresh Herbs

Karen L. Panter^{1,4}, Timmothy M. Gergeni^{2,3},
Casey P. Seals^{2,3}, and Andrea R. Garfinkel^{2,3}

ADDITIONAL INDEX WORDS. *Allium tuberosum*, cut flowers, *Helianthus annuus*, hoop houses, *Origanum majorana*, *Origanum vulgare*

SUMMARY. High tunnels are gaining popularity for their use in horticultural crop production. However, little is known about the effect of high tunnel orientation on plant growth and development. In this set of studies, we show tunnel orientation does not necessarily affect the production of cut sunflower (*Helianthus annuus*) and culinary herbs oregano (*Origanum vulgare*), marjoram (*Origanum majorana*), and garlic chive (*Allium tuberosum*). Two high tunnels, one with the long axis oriented north-south (NS) and the other east-west (EW), were used to test the effects of high tunnel orientation on several crops over a 5-year period: cut sunflower (2012 and 2016); marjoram, oregano, and garlic chive (2013 and 2014); and garlic chive (2015). The tunnels are 12 × 16 ft, smaller than those used in commercial production. The size would be appropriate for hobby and seasonal production of horticultural crops for local markets. Cut sunflower stems were similar lengths both years in both high tunnels. Sunflower times to harvest were different between cultivars but not between high tunnels. Oregano fresh weight yields were highest in the NS tunnel in 2013 but similar between tunnels in 2014. Marjoram fresh weights were highest in 2013 in the EW tunnel but highest in 2014 in the NS tunnel. Garlic chive fresh weights were similar between tunnels all 3 years. We show that differences are more a function of innate cultivar characteristics than which way small high tunnels are oriented.

High tunnels are important alternatives to greenhouse or field production of high-value crops such as cut flowers and herbs (Carey et al., 2009; Wells and Loy, 1993; Wien, 2009b). Many cut flowers and herbs produced in the United States are grown in open field settings where year-round production is possible (Bonarriva, 2003); however, the interest in locally produced horticultural goods has driven the regional production of these crops

into greenhouses, high tunnels, and other season-extending environments (Cavins et al., 2000; Ortiz et al., 2012; Wien, 2009a). Not only do high tunnels allow for a local extended growing season, but they also play an important role in protecting cut flower crops from extreme environmental conditions, including freezing temperatures, rain, wind, hail, and in some cases, disease (Lamont, 2009; Wien, 2009a). The increased interest in domestic production of cut flowers and herbs in high tunnels and the advantages of high tunnel production create the need for region- and crop-specific research to maximize local production

potential (Ferguson et al., 2012; Lamont, 2009; Ortiz et al., 2012).

Numerous factors may affect growth of crops in high tunnels, including orientation of the structure (Sethi, 2009). General studies on high tunnel orientation indicate that growers in the northern hemisphere should orient high tunnels with the long sides running NS to maximize solar radiation (Sethi, 2009). Despite this recommendation, there is evidence to show that the benefits of high tunnel orientation and growing location within a particular orientation may be crop-specific. Taber et al. (2009) showed that tunnel orientation and planting location within a high tunnel negatively affected bean (*Phaseolus vulgaris*) yield, with decreased productivity of bean plants grown in north (in an EW-oriented tunnel) and west (in a NS-oriented tunnel) growing locations.

The sunflower will flower from 75 to 150 d after sowing (Fernández-Martínez et al., 2009), with temperature and photoperiod significantly affecting time from sowing to maturity (Fernández-Martínez et al., 2009; Schuster, 1985). Cool temperatures delay flowering, whereas warm temperatures accelerate maturation (Fernández-Martínez et al., 2009).

Due to their hardiness, garlic chive is described as being low maintenance (Folia, 2015). Chive plants do best in full sun or part shade, and well-drained but moist soil (Gardener's Network, 2016). Recommended growing temperature is 58 °F (Nau, 1999).

Oregano and marjoram require similar growing temperatures of ≈55 °F. Both are frost-sensitive and are typically grown as annuals in colder climates, unless under controlled environments (Nau, 1999).

Received for publication 9 Jan. 2019. Accepted for publication 23 Apr. 2019.

Published online 6 June 2019.

The authors appreciate funding support from the Wyoming Department of Agriculture Specialty Crops Grant Program and from the University of Wyoming Agricultural Experiment Station.

¹Plant Sciences – Department 3354, University of Wyoming, 1000 East University Avenue, Laramie, WY 82071

²Department of Plant Sciences, University of Wyoming

³Former graduate student.

⁴Corresponding author. E-mail: kpanter@uwyo.edu.

This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.21273/HORTTECH04273-19>

Units

To convert U.S. to SI, multiply by	U.S. unit	SI unit	To convert SI to U.S., multiply by
10.7639	ft	lx	0.0929
0.3048	ft	m	3.2808
0.0929	ft ²	m ²	10.7639
2.54	inch(es)	cm	0.3937
1.1209	lb/acre	kg·ha ⁻¹	0.8922
0.0254	mil(s)	mm	39.3701
1	mmho/cm	dS·m ⁻¹	1
28.3495	oz	g	0.0353
(°F – 32) ÷ 1.8	°F	°C	(°C × 1.8) + 32

Although a number of studies have assessed the potential for producing cut sunflowers and herbs in high tunnels (Cavins et al., 2000; Lamont, 2009; Lopez and Ortiz, 2012; Ortiz et al., 2012; Wien, 2006), none report on the effect of high tunnel orientation or growing location on plant growth. Thus, we set up a series of trials with the objective of determining differences in growth of cut sunflowers and fresh herbs between high tunnel orientations. The tunnels are small hobby size; therefore, these studies are appropriate for part-time growers interested in seasonal production of horticultural crops for local markets. This work represents a combination of three different graduate students' effort over 4 years; some differences in methodology occurred as a result.

Materials and methods

Site background

Experiments were conducted at the University of Wyoming Laramie Research and Extension Center greenhouse complex. The facility is located in Laramie, WY (lat. 41°19'12"N, long. 105°33'31"W, elevation of 7200 ft) in U.S. Department of Agriculture cold hardiness zone 4b (Fig. 1C).

High tunnels

The high tunnels used in this study were two 12 × 16-ft arch-roof-style tunnels (Solar Star Greenhouses; Growers Supply, Dyersville, IA) covered in a double layer of 6-mil uninflated polyethylene plastic with double-walled polycarbonate end walls. Both tunnels had manual roll-up sides, up to ≈3 ft, to provide ventilation. Sides were rolled up when temperatures exceeded 40 °F and down when temperatures dipped below 40 °F. The two tunnels were oriented in different directions, one with the long axis running NS and the other EW (Fig. 2). In Apr. 2012, 2013, and 2015, before planting, soil in the tunnels was amended with 2 inches of peatmoss and compost, spaded into a depth of 6 inches. Soil in the tunnels is described as sandy loam.

Sunflower

Two cultivars of fresh-cut sunflower were grown in 2012 and 2016. Cultivar selections were based on flower color (brown and gold, the University of Wyoming's colors) and

single-stem characteristics. Single-stem sunflowers do not branch. One seed equals one plant that produces one solitary flowering stem (Harris Seeds, Rochester, NY). The cultivars were Dafna and ProCut Bicolor (Harris Seeds) (Fig. 1A and B).

SUNFLOWER 2012. The two sunflower cultivars were arranged in a randomized complete block design with four replications of four plants each per cultivar arranged in a two-by-two square with plants on 6-inch centers. Blocks were arranged on either side of the central aisle in each high tunnel, four blocks on either side, eight blocks total per tunnel. An edge row was planted around the perimeter of all sunflower plots to mitigate edge effects.

SUNFLOWER 2016. The same two cultivars were arranged in a completely random design with 20 single-plant replicates per cultivar on 6-inch centers. Twenty plants were randomized on each side of the central aisle in each high tunnel, 40 plants total per tunnel. An edge row was planted around the perimeter of all plots to mitigate edge effects. Different students pursued these studies in 2012 and 2016; thus the differences in methodology between the 2 years.

PROPAGATION. Seeds were sown on 2 May 2012 and 26 May 2016 in 128-cell plug trays in a professional germination medium (Fafard Super Fine Germinating Mix; Conrad Fafard, Agawam, MA) and maintained in a greenhouse 70/65 °F day/night on a mist bench. Misting was controlled with an automated clock system (Schaefer Solar 3B; Pinnacle Climate Technologies, Sauk Rapids, MN). The mist system was set to trigger on for 20 s for every 20 solar units collected, where one solar unit is equal to 0.02 mol·m⁻² light. Seeds from different cultivars, Sunrich Gold (2012) and Sunbright Supreme (2016) (Harris Seeds), were sown on the same schedule and were used as edge rows.

After 15 d (17 May 2012 and 10 June 2016), seedlings were transplanted on 6-inch centers directly into the ground in the high tunnels and immediately irrigated. Thirteen d following transplant, each plant was top-dressed with 2.5 g (1076.4 kg·ha⁻¹) 15N-4.4P-9.9K controlled-release fertilizer (Osmocote Plus 15-9-12; ICL Specialty Fertilizers, Dublin, OH). Throughout production,

sunflowers were watered by hand daily. Plants were grown under natural light and daylength conditions experienced in the hobby high tunnels.

Herbs

Three species of fresh culinary herbs were chosen for study in 2013 and 2014: oregano, marjoram, and garlic chive (Fig. 1C). The latter was used in 2015. Each of the species in all 3 years was grown as 16 (2013 and 2014) and 24 (2015) single-plant replicates. Each group of one species was grown together in one plot; plots were randomly placed one on each side of the central aisle in each tunnel (Fig. 1C). Plots were completely surrounded by edge rows of a combination of lavender (*Lavandula angustifolia*) and rosemary

Fig. 1. Photos of plants used in high tunnel orientation studies. Fresh-cut sunflower cultivars grown were Dafna (A) and ProCut Bicolor (B) in 2012 and 2016. Fresh herbs grown are shown in (C), with the photo taken of crops in the high tunnel oriented north-south in 2013.

(*Rosmarinus officinalis*) plants. Again, differences in methodology reflect different students handling the work.

PROPAGATION. All plants were grown from seeds (Harris Seeds). Sowings occurred 4 Apr. 2013, 15 Apr. 2014, and 8 Mar. 2015. Seeds were sown in 128-cell plug trays in a professional germination medium (Fafard Super Fine Germinating Mix). Plug trays were placed on a mist bench using a solar mist controller (Schaefer Solar 3B) set to come on for 20 s every 20 solar counts. Following germination, seedlings were transplanted directly into the ground in the high tunnels on 12-inch centers: 30 May 2013, 24 May 2014, and 22 May 2015. Ten days following transplant, each plant was top-dressed with 2.5 g (538.2 kg·ha⁻¹) 15N-4.4P-9.9K controlled-release fertilizer (Osmocote Plus 15-9-12). Throughout production, herbs were watered by hand once daily. Plants were grown under natural light and daylength conditions.

Light and temperature measurements

Environmental monitors (HOBO Pendant Temperature/Light Data Logger 64K, UA-002-64HOBO; Onset Computer Corp., Bourne, MA) were positioned 12 inches above the ground in the center of each block of plants to record air temperature and light levels, two per high tunnel, four sensors total in 2012 through 2015.

Soil temperatures were monitored using data loggers (HOBO Pendant Temperature Data Logger 64K, UA-001-64; Onset Computer Corp.). Each was buried 4 inches deep in the soil in the center of the long axis of each tunnel, ≈4 inches from the outside wall. Thus, there were two data loggers per high tunnel, four total (Fig. 2). All data loggers were programmed to collect data every 4 h starting at midnight.

Soil analyses

Soil analyses in each tunnel were conducted in spring 2013, 2014, and 2015 (Colorado Analytical Laboratories, Brighton, CO). Results indicated minor differences in characteristics between the two tunnels. The soil texture was sandy loam. In 2013, pH and electrical conductivity (EC) were 7.9 and 1.3 dS·m⁻¹, respectively,

Fig. 2. Diagram of two high tunnels oriented perpendicular to one another. Each is 12 × 16 ft. One is oriented with the long axis north-south and the other with the long axis east-west. The high tunnels are located at the University of Wyoming Laramie Research and Extension Center in Laramie, WY (lat. 41°19'12"N, long. 105°33'31"W, elevation 7200 ft). Plants were grown on either side of the central aisle in each high tunnel. Air temperatures were monitored in the center of the growing area of each tunnel at 12 inches, and soil temperatures were monitored 4 inches deep in the center of the long side of each high tunnel; 1 ft = 0.3048 m, 1 inch = 2.54 cm.

in the NS tunnel and 7.7 and 1.5 dS·m⁻¹ in the EW tunnel. In 2014, the analysis results were pH 7.6 and EC 2.6 in the NS tunnel and 7.4 and 3.6 in the EW tunnel. In 2015, pH levels in the NS tunnel were 7.6 with EC at 1.2; EW pH was 7.7 with EC at 1.3.

Organic matter (OM) levels were below 5% in both tunnels. In 2013, the NS tunnel OM was 3.5%, and the EW tunnel was 3.1%. In 2014, the figures were 4.5% in the NS tunnel and 4.9% in the EW tunnel. In 2015, levels were NS tunnel 3.5% and EW tunnel 2.9%.

Harvesting and data collection

SUNFLOWER. Individual flowers were harvested when ray flowers were

open (Dole and Wilkins, 1999), as per industry standard. Harvest occurred approximately every other day. Stems were cut at the soil level and stem lengths were immediately measured and recorded. The length of the stem was measured from the soil level where the cut was made to the base of the flower head receptacle. Data were also collected on time to harvest from sowing (days).

HERBS. In 2013 and 2014, oregano and marjoram were harvested periodically during the growing season, just before flower initiation, at their peak culinary flavor (Morgan, 2005), as per industry standard. These were harvested by hand leaving two to three nodes on each stem for regrowth. Garlic chive was harvested

when the aboveground biomass was at least 6 inches tall, cut to 2 inches aboveground. Harvested materials were immediately weighed for fresh weight. All fresh weights were measured using a portable scale (PC 4400; Mettler-Toledo International, Greifensee, Switzerland). In 2015, garlic chive was harvested once at the end of the growing season for total fresh weight, 10 Sept., and weighed using a portable scale (model SLB50; Carolina Biological Supply Co., Burlington, NC).

Statistical analysis

Data were analyzed and statistical computations were facilitated using analysis of variance procedure in SAS (version 9.3 or 9.4; SAS Institute; Cary, NC). Means separations were conducted using Fisher's least significant difference at $\alpha = 0.05$.

Results

Sunflower: Time to harvest

2012. Time to harvest did not differ between NS and EW high tunnel orientations (Table 1). The only differences observed in time to harvest were between cultivars. Time to harvest between the two high tunnel orientations, NS vs. EW, for 'ProCut Bicolor' averaged 77 and 79 d, respectively, which was significantly fewer than 'Dafna' at 95 and 94 d, respectively (Table 1).

2016. Time to harvest again differed between the cultivars Dafna and ProCut Bicolor but there were no differences between the two high tunnel orientations. 'Dafna' averaged 80 d in both high tunnels, whereas 'ProCut Bicolor' averaged 71 d in the NS tunnel and 76 d in the EW tunnel (Table 1).

Sunflower: Stem lengths

2012. Neither 'Dafna' nor 'ProCut Bicolor' differed in stem length between cultivars or the two high tunnel orientations (Table 1).

2016. There were no differences in stem lengths between cultivars or high tunnels (Table 1).

Herbs: Time to harvest

2013, 2014, and 2015. No differences occurred in time to harvest from sowing between the two high tunnels with any crop in either 2013 or 2014, because each crop was

harvested the same days in both high tunnels (data not shown). In 2015, all garlic chives were harvested at once at the end of the growing season, so no differences in time to harvest occurred that year either (data not shown).

Herbs: Fresh weights

2013. There were no differences in fresh weight yields per plant between the two high tunnel orientations with garlic chive. An average of 13 g of garlic chive per plant was harvested from the NS tunnel and 16 g from the EW tunnel (Table 2). Marjoram and oregano per plant fresh weights differed between the two high tunnels (Table 2). Marjoram per plant fresh weight in the EW tunnel was 115 g, significantly higher than in the NS tunnel, 85 g (Table 2). Oregano showed the opposite, with higher per plant fresh weight in the NS tunnel at 117 g than the EW tunnel, 89 g (Table 2).

2014. Garlic chive and oregano per plant yields between the two high tunnels were not different (Table 2). Garlic chive produced 37 g per plant in the NS tunnel and 40 g in the EW tunnel. Oregano averaged 106 g per plant in both high tunnels in 2014; however, marjoram yields were higher in the NS tunnel (145 g) than in the EW tunnel (110 g) (Table 2).

2015. Marjoram and oregano were not grown in the tunnels in 2015. Garlic chive yields per plant were similar between the two high tunnels. The NS tunnel averaged 25 g per plant and the EW high tunnel averaged 22 g on per plant bases (Table 2).

Environment

Average high tunnel air temperatures in 2013, 2014, and 2015 were not different (Table 3). Example temperatures during the 2012 growing season ranged from a low on 20 May of 33.6 °F in both tunnels to highs on

Table 1. Results of analysis of variance and means separation in fresh-cut sunflower time to harvest from sowing and stem lengths when grown in two high tunnels in 2012 and 2016. One high tunnel was oriented with the long axis oriented north-south (NS), and the other was oriented east-west (EW).

Yr	Cultivar	High tunnel	Mean time to harvest (d)	Mean stem length (cm) ^z
2012	Dafna	NS	95 a ^y	123 a
		EW	94 a	110 a
	ProCut Bicolor	NS	77 b	103 a
		EW	79 b	109 a
2016	Dafna	NS	80 a	82 a
		EW	80 a	84 a
	ProCut Bicolor	NS	71 b	81 a
		EW	76 b	72 a

^z1 cm = 0.3937 inch.

^yDifferent letters within a year, cultivar, and column indicate statistical differences using Fisher's least significant difference means separation at $\alpha = 0.05$.

Table 2. Results of analysis of variance and means separation of fresh weight of culinary herbs garlic chive, marjoram, and oregano, grown in two high tunnels in 2013, 2014, and 2015. One high tunnel was oriented with the long axis oriented north-south (NS), and the other was oriented east-west (EW).

Yr	Mean fresh wt (g/plant) ^z			
	High tunnel	Garlic chive	Marjoram	Oregano
2013	NS	13 a ^y	85 b	117 a
	EW	16 a	115 a	89 b
2014	NS	37 a	145 a	106 a
	EW	40 a	110 b	106 a
2015	NS	22 a	Not grown	Not grown
	EW	25 a	Not grown	Not grown

^z1 g = 0.0353 oz.

^yDifferent letters within a year, herb, and column indicate statistical differences using Fisher's least significant difference means separation at $\alpha = 0.05$.

10 July of 106.8 °F in the EW tunnel and 105.8 °F in the NS tunnel. Average soil temperatures in the east and south planting locations were statistically higher than those in the west and north locations in 2012, 2013, and 2015; in 2014, east average soil temperature was higher than the other three edges (Table 3). Soil temperatures within the hobby high tunnels were consistently lower than air temperatures (Table 3).

Light levels differed between high tunnels depending on the time of day, as shown in data from 2012, a representative year (Fig. 3). Lower average light levels were recorded in the NS tunnel than the EW tunnel, particularly at 0800 and 1200 HR (Fig. 3).

Discussion

Sunflower: Time to harvest

The similarities between high tunnel orientations with regard to time to harvest were expected because all were grown under the same photoperiod. Photoperiod has been shown to be the major factor in determining flower initiation and maturation of cut sunflowers (Blacquière et al., 2002; Gimelli et al., 2003; Hayata and Imaizumi, 2000; Pallez et al., 2002; Yañez et al., 2004, 2005). Variation in time to harvest between the two cultivars tested in this study are in accordance with research that suggests that cut sunflowers are sensitive to photoperiod and that this response is cultivar-specific (Wien, 2012a, 2012b). Wien (2012a, 2012b) showed that ‘ProCut Bicolor’ flowered faster under summer, long-day conditions, whereas ‘Dafna’ is slower to bloom during long-day conditions. Fourteen hours has been shown to be the critical daylength to elicit a long-day sunflower response (Blacquière et al., 2002); daylength exceeded 14 h throughout both years of study.

Sunflower: Stem lengths

Because stem lengths were similar between the two high tunnels, lower light intensities in the NS high tunnel were apparently not different enough to cause stem lengths to differ. In our other studies conducted on sunflower responses to light (Garfinkel and Panter, 2014), we found light intensity contributed to differences in stem length of cut sunflowers when grown in a greenhouse. Soil temperature differences in our studies were statistically different but not

Table 3. Mean high tunnel air (2013, 2014, 2015) and soil temperatures (2012, 2013, 2014, and 2015). Two high tunnels were used, one with the long axis oriented north-south and the other east-west. Crops grown were fresh-cut sunflowers or fresh culinary herbs. Locations within high tunnels were as follows: north and south sections in the east-west tunnel, and east and west sections in the north-south tunnel. Temperatures were recorded every 4 h.

Yr	Location within high tunnel	Air temp (°C) ^z	Soil temp (°C)
2012	North	NA ^y	14.4 b ^x
	South	NA	15.7 a
	East	NA	15.7 a
	West	NA	14.6 b
2013	North	19.2 a	15.4 b
	South	19.5 a	17.0 a
	East	20.7 a	16.8 a
	West	20.0 a	16.6 b
2014	North	18.8 a	15.6 a
	South	18.6 a	15.8 a
	East	19.6 a	16.6 a
	West	18.7 a	15.6 a
2015	North	25.7 a	18.4 b
	South	26.1 a	20.0 a
	East	26.4 a	20.3 a
	West	25.6 a	18.3 b

^z $(1.8 \times ^\circ\text{C}) + 32 = ^\circ\text{F}$.

^yNot applicable; air temperatures were not recorded in 2012.

^xDifferent letters within a year, location, and column indicate statistical differences using Fisher’s least significant difference means separation at $\alpha = 0.05$.

Fig. 3. Plot of light intensity by high tunnel orientation and time of day. Data are from two high tunnels oriented perpendicular to one another, one with the long axis north-south [NS (red)] and the other east-west [EW (blue)]. Data were recorded every 4 h starting at midnight 17 May 2012 through 7 Oct. 2012. Each is 12 × 16 ft. The high tunnels are located at the University of Wyoming Laramie Research and Extension Center in Laramie, WY (lat. 41°19'12"N, long. 105°33'31"W, elevation 7200 ft); 1 ft = 0.3048 m, 1 lx = 0.0929 fc.

biologically, because stem lengths were the same between the two tunnels. Optimum soil temperatures for sunflowers are ≈68 °F (McMichael

and Burke, 1998). Our soil temperatures were lower than recommended, but this did not contribute to parameters measured.

Herbs

2013, 2014, AND 2015. Differences in fresh weights among the three species did not show any particular patterns among the 3 years. Fresh weights of the three species varied by year and high tunnel with no easily discernable cause. Reasons for higher fresh weights in general in 2014 probably include the human component of simply being more experienced growers. It seems differences between the tunnels in soil temperatures, light levels, and soil properties had little effect overall. Soil temperatures were slightly lower in 2014, but fresh weights were slightly higher that year. Light levels between the two tunnels were slightly higher in the EW tunnel but herb fresh weights were not always higher in that tunnel in either year. Soil temperatures were lower on the north and west sides, but only in 2013, not 2014.

Parameters for optimum growth of oregano and sweet marjoram include temperatures of 65 to 70 °F days and 55 to 62 °F nights (Nau, 2011). Temperatures in the two high tunnels were within these ranges in 2013 and 2014, indicating temperatures were not a factor in fresh weight differences. Recommended pH levels for these two herbs are between 4.9 and 8.7, with optimum at ≈ 6.9 (Savio and Robinson, 1998); ours were in this range, indicating soil properties did not lead to fresh weight differences.

Chive generally requires temperatures of 55 to 60 °F days and 40 to 55 °F nights (Nau, 2011). Our temperatures were warmer than recommended but were similar between the two tunnels, so were not factors in fresh weight differences. The recommendation for pH is between 5.8 and 7.0 (Andersen, n.d.), lower than ours. However, we saw no differences between tunnels in chive fresh weights.

Environment

Although these high tunnels are small and hobby-sized, this did not influence crop production parameters. Recommendations are for high tunnels to be oriented perpendicular to prevailing winds (Jett, 2004); however, because our two high tunnels were open with the sides rolled up during all five growing seasons, temperatures were consistent within the two. Air temperatures were higher in 2015 than the previous 2 years, but garlic chive

fresh weights were intermediate in 2015, between those from the two previous years. It seems air temperatures are not a contributing factor in herb yields or sunflower time to harvest or stem lengths; however, soil temperatures were consistently lower on the north side of the EW tunnel and the west side of the NS tunnel.

Conclusions

Our data suggest that in small tunnels with these dimensions in this location, orientation was not a primary production consideration. In this series of experiments, we found that stem lengths of two cultivars of sunflower were unaffected by high tunnel orientation; time to harvest was also unaffected. Variation in time to harvest and stem length among cultivars can likely be attributed to innate variations among cultivars and their responses to different environmental conditions including photoperiod. Therefore, cultivar selection would be more important when considering growing fresh-cut sunflowers in high tunnels. Herb fresh weights were not consistent from year to year in any location, suggesting either high tunnel orientation is acceptable for growing them. This study focused on production in small, hobby-size high tunnels. Further research would be necessary to elucidate differences in production parameters with larger tunnels.

Literature cited

- Andersen, C.R. n.d. Chives. 8 Jan. 2019. <<https://www.uaex.edu/publications/pdf/FSA-6082.pdf>>.
- Blacquire, T., N. Straver, and D. van den Berg. 2002. Possibilities for using photoperiodism to program flowering of sunflowers (*Helianthus annuus*) in the greenhouse and in the open. *Acta Hort.* 580:101–109.
- Bonarriva, J. 2003. Industry & trade summary: Cut flowers. U.S. Intl. Trade Commission Publ. 3580.
- Carey, E.E., L. Jett, W.J. Lamont, Jr., T.T. Nennich, M.D. Orzolek, and K.A. Williams. 2009. Horticultural crop production in high tunnels in the United States: A snapshot. *HortTechnology* 19:37–43.
- Cavins, T.J., J.M. Dole, and V. Stambach. 2000. Unheated and minimally heated winter greenhouse production of specialty cut flowers. *HortTechnology* 10:793–799.
- Dole, J.M. and H.F. Wilkins. 1999. Floriculture – Principles and species. Prentice Hall, Upper Saddle River, NJ.
- Ferguson, C.G., P.G. Vuppapapati, M.J. Stone, and E. Gray. 2012. Evaluation of cultivars of sunflowers (*Helianthus annuus* L.) and selected environments for production of cut flowers. *J. Ky. Acad. Sci.* 73:90–95.
- Fernndez-Martinez, J.M., B. Perez-Vich, and L. Velasco. 2009. Sunflower, p. 155–232. In: J. Vollman and I. Rajcan (eds.). *Oil crops*, Vol. 4. Springer Sci. Business Media, New York, NY.
- Folia, 2015. Garlic chive *Allium tuberosum*. Plant database/garlic chive plant care. 8 Jan. 2019. <<https://myfolia.com/plants/919-garlic-chive-allium-tuberosum>>.
- Gardener's Network. 2016. How to grow chives. 7 Feb. 2019. <<http://www.gardenersnet.com/herbs/chives.htm>>.
- Garfinkel, A.R. and K.L. Panter. 2014. Year-round greenhouse production of cut sunflowers in the Rocky Mountain West. *HortTechnology* 24:743–748.
- Gimelli, F., A. Loporati, and L. Maggiora. 2003. The all year round cultivation of cut sunflower in San Remo district. *Acta Hort.* 614:553–560.
- Hayata, Y. and Y. Imaizumi. 2000. Effect of photoperiod on flower bud development of ornamental sunflowers. *J. Jpn. Soc. Hort. Sci.* 69:708–710.
- Jett, L.W. 2004. High tunnel tomato production. Univ. Missouri Ext. Bul. M170. 8 Jan. 2019. <<https://mospace.umsystem.edu/xmlui/bitstream/handle/10355/51733/m0170-2004.pdf?sequence=1>>.
- Lamont, W.J., Jr. 2009. Overview of the use of high tunnels worldwide. *HortTechnology* 19:25–29.
- Lopez, R.G. and M.A. Ortiz. 2012. Do high tunnels make the cut? 8 Jan. 2019. <<http://www.greenhousegrower.com/article/28959/do-high-tunnels-make-the-cut>>.
- McMichael, B.L. and J.J. Burke. 1998. Soil temperature and root growth. *HortScience* 33:947–951.
- Morgan, L. 2005. Fresh culinary herb production – A technical guide to the hydroponic and organic production of commercial fresh gourmet herb crops. Suntec (NZ), Tokomaru, New Zealand.
- Nau, J. 1999. Ball culture guide. The encyclopedia of seed germination. 3rd ed. Ball Publ., Batavia, IL.
- Nau, J. (ed.). 2011. Ball RedBook. Ball Publ., West Chicago, IL.

- Ortiz, M.A., K. Hyrczyk, and R.G. Lopez. 2012. Comparison of high tunnel and field production of specialty cut flowers in the Midwest. *HortScience* 47:1265–1269.
- Pallez, L.C., J.M. Dole, and B.E. Whipker. 2002. Production and post-production studies with potted sunflowers. *HortTechnology* 12:206–210.
- Savio, Y. and C. Robinson. 1998. Oregano, winter marjoram, wild marjoram, pot marjoram. In: Specialty and minor crops handbook. 8 Jan. 2019. <https://www.ctahr.hawaii.edu/sustainag/extn_pub/veggie%20pubs/Oregano,%20Winter%20Marjoram,%20Wild%20Marjoram,%20Pot%20Marjoram.pdf>.
- Schuster, W.H. 1985. *Helianthus annuus*, p. 98–121. In: A.H. Halevy (ed.). CRC handbook of flowering, Vol. 3. CRC Press, Boca Raton, FL.
- Sethi, V.P. 2009. On the selection of shape and orientation of a greenhouse: Thermal modeling and experimental validation. *Sol. Energy* 83:21–38.
- Taber, H.G., B.J. Havlovic, and N.P. Howell. 2009. High tunnel pole bean evaluation. Iowa State Res. Farm Prog. Rpt. 468.
- Wells, O.S. and J.B. Loy. 1993. Row-covers and high tunnels enhance crop production in the northeastern United States. *HortTechnology* 3:92–95.
- Wien, C.H. 2006. Growing black-eyed susan and sunflower out of season. *Growing for Market*. 15(6):17–18.
- Wien, C.H. 2009a. Daylength response of sunflowers. Dept. Hort., Cornell Univ., Ithaca, NY.
- Wien, C.H. 2009b. Floral crop production in high tunnels. *HortTechnology* 19:56–60.
- Wien, C.H. 2012a. Cut flower cultural practice studies and variety trials. Dept. Hort., Cornell Univ., Ithaca, NY.
- Wien, C.H. 2012b. Sunflower seedling daylength response. Dept. Hort., Cornell Univ., Ithaca, NY.
- Yañez, P., H. Ohno, and K. Ohkawa. 2004. Effect of photoperiod on flowering and growth of ornamental sunflower cultivars. *Environ. Control Biol.* 42:287–293.
- Yañez, P., H. Ohno, and K. Ohkawa. 2005. Photoperiodic response and vase life of ornamental sunflower cultivars. *HortTechnology* 15:386–390.