Duration of CO₂ Enrichment Influences Growth, Yield, and Gas Exchange of Two Tomato Species

Serge Yelle, Richard C. Beeson, Jr., Marc J. Trudel, and André Gosselin

Département de Phytologie, Faculté des Sciences de l'Agriculture et de l'Alimentation, Université Laval, Québec, Que. G1K 7P4, Canada

Additional index words. *Lycopersicon esculentum*, *Lycopersicon chmielewskii*, photosynthesis, stomatal conductance, internal CO₂

Abstract. *Lycopersicon esculentum* Mill. cv. Vedettos and *Lycopersicon chmielewskii* Rick, LA 1028, were exposed to two CO₂ concentrations (330 or 900 $\mu\text{mol}\cdot\text{m}^{-3}$) for 10 weeks. The elevated CO₂ concentration increased the relative growth rate (RGR) of *L. esculentum* and *L. chmielewskii* by 18% and 30%, respectively, after 2 weeks of treatment. This increase was not maintained as the plant matured. Net assimilation rate (NAR) and specific leaf weight (SLW) were always higher in CO₂-enriched plants, suggesting that assimilates were preferentially accumulated in the leaves as reserves rather than contributing to leaf expansion. Carbon dioxide enrichment increased early and total yields of *L. esculentum* by 80% and 22%, respectively. Carbon exchange rates (CER) increased during the first few weeks, but thereafter decreased as tomato plants acclimated to high atmospheric CO₂. The relatively constant concentration of internal CO₂ with time suggests that reduced stomatal conductance under high CO₂ does not explain lower photosynthetic rates of tomato plants grown under high atmospheric CO₂ concentrations. Leaves 5 and 9 responded equally to high CO₂ enrichment throughout plant growth. Consequently, acclimation of CO₂-enriched plants was not entirely due to the age of the tissue. After 10 weeks of treatment, leaf 5, which had been exposed to high CO₂ for only 10 days, showed the greatest acclimation of the experiment. We conclude that the duration of exposure of the whole plant to elevated CO₂ concentration, rather than the age of the tissue, governs the acclimation to high CO₂ concentrations.

Carbon dioxide enrichment of greenhouses and the projected doubling of the atmospheric CO₂ concentration within the next century (Gates et al., 1983; Lemon, 1983) have stimulated interest in the long-term effects of high CO₂ levels on plant growth. The enhancement of photosynthesis, growth, and yield by increasing the concentration of CO₂ in the atmosphere has been shown for many species (Kimball, 1983). Carbon dioxide enrichment to 1000 $\mu\text{mol}\cdot\text{m}^{-3}$ increased yields and plant dry weight of tomato plants by 30% (Slack, 1986) and 36% (Yelle et al., 1987), respectively. Cucumber (Dennis, 1980), lettuce (Knecht and O'Leary, 1983), and many ornamental species (Mortensen, 1987) also responded positively to CO₂ enrichment. However, a side-effect of long-term growth in a CO₂-enriched environment is diminished photosynthetic efficiency of leaves. Declines in photosynthesis were demonstrated for tomatoes (Hurd, 1968; Hicklenton and Joliffe, 1980), cucumbers (Aoki and Yabuki, 1977; Frydrieh, 1976; Peet et al., 1986), and several other species (Mauney et al., 1978; Mortensen, 1983; Wong et al., 1979). For instance, Aoki and Yabuki (1977) reported that CO₂ enrichment increased the photosynthetic rate of cucumbers by 76% after 1 day of treatment. However, as enrichment continued, the photosynthetic rate at the high CO₂ concentration rapidly decreased below that of the control level. Although this effect is well-known, there is no consensus yet on the physiological and biochemical causes.

Several reasons have been proposed to explain plant acclimation to high CO₂ concentrations: decreases in stomatal conductance (Peet et al., 1986), reduced Rubisco activity (Peet et al., 1986; Spencer and Bowes, 1986; Vu et al., 1983), and buildup of carbohydrates (Clough et al., 1981; Sasek et al., 1985).

The present study compared the long-term response of two tomato species, *L. esculentum* and *L. chmielewskii*, to elevated atmospheric CO₂. These species differ greatly in their sink metabolism and carbohydrate assimilation. Yelle et al. (1988) showed that *L. chmielewskii* accumulated less starch than *L. esculentum*. This difference was related to a lower level of ADP glucose pyrophosphorylase and a higher level of phosphorylase in *L. chmielewskii* fruits. A low level of invertase and a nondetectable level of sucrose synthase in *L. chmielewskii* were associated with the high level of sucrose in its fruits. *L. chmielewskii* had twice the soluble solids content in the fruits as *L. esculentum*. There were three objectives in this study: 1) assess whether the sink metabolism and carbohydrate assimilation of *L. chmielewskii* would prevent the long-term loss of efficiency resulting from CO₂ enrichment; 2) verify whether the loss of efficiency was due to the age of the tissue rather than the time of exposure to high CO₂ levels; 3) determine the relationship between stomatal conductance and internal CO₂ concentrations on the carbon exchange rates associated with CO₂ acclimation.

Materials and Methods

Plant material. 'Vedettos' (*Lycopersicon esculentum*) and LA 1028 (*L. chmielewskii*) tomatoes were seeded in rockwool blocks (Grodania, Hedehusene, Denmark) on 15 Dec. 1987 and transplanted on 17 Jan. 1988 in 16 independent hydroponic systems (NFT). Tomato plants were grown in four identical glasshouse compartments (24 m²). The CO₂ concentration within each compartment was maintained at 330 $\pm$ 50 (control, ambient concentration) or 900 $\pm$ 50 $\mu\text{mol}\cdot\text{m}^{-3}$. Each CO₂ treatment was repeated twice. Both species were repeated twice in each compartment. Each experimental unit was randomly distributed in each compartment and consisted of 20 plants.

Carbon dioxide levels were monitored and controlled by an infrared gas analyzer (Priva Computers, No. APBA 251 E CO₂ Monitor). The CO₂ was supplied from a pure liquid source during the photoperiod. Day and night temperatures were maintained at a minimum of 22 $\pm$ 2C and 17 $\pm$ 2C, respectively.

Received for publication 26 Oct. 1988. We gratefully acknowledge the Québec Ministries of Agriculture and Food, and of Education and Science for their financial support. The skilled technical assistance of René Pouliot, Jacques Debroux, and Linda Gaudreau is also gratefully acknowledged. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

Forced ventilation was activated when air temperature reached 30°C. During forced ventilation, CO₂ levels dropped to 500 $\mu\text{mol}\cdot\text{m}^{-3}$. Forced ventilation accounted for $\approx 1\text{ hr/day}$ during the first half of the experiment and 2 hr/day during the second. Temperatures were measured using six thermocouples in each compartment; four for the nutrient solutions and two for air temperatures. Thermocouples were connected to a Minitrend 205 recorder (Doric Scientific). Supplemental lighting (150 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}\cdot\text{PPF}$) was supplied by high-pressure sodium lamps for a photoperiod of 16 hr.

Nutrient solutions contained (in $\mu\text{l}\cdot\text{liter}^{-1}$) 176 N, 46 P, 284 K, 140 Ca, 47 Mg, 2 Fe, 1 Mn, 0.4 Zn, 0.29 B, 0.03 Cu, and 0.06 Mo and were renewed every 5 days. Solutions were adjusted daily with H₂SO₄ to a pH of 5.8 and to an electric conductivity of 220 dS $\cdot\text{cm}^{-1}$ using a complete nutrient solution.

Vegetative growth and gas exchange. Fresh and dry weights of the leaves and stems and leaf areas were measured every 2 weeks for 10 weeks on four plants randomly sampled from each experimental unit. After each plant removal, the remaining plants were redistributed into the canopy to minimize patchy light distribution. Leaves and stems were dried for 48 hr at 70°C before measurement of the dry weights. Leaf area was measured with a LI-COR portable area meter (LI-3000, LI-COR, Lincoln, Neb.). Relative growth rate (vegetative growth), net assimilation rate, and leaf area ratio were calculated as reported by Gardner et al. (1985). Ripe tomato fruits were harvested during 4 weeks from 16 Mar. to 15 Apr. Early yields corresponded to the first 3 weeks of harvest. For the last harvest date (15 Apr.), all green and red fruits of all sizes were picked and included in the total yield. Yield of *L. chmielewskii* is not reported because of frequent fruit abscission and high variability however, it was about 100 times lower than that of *L. esculentum*, with an average fruit size of $\approx 8\text{ g}$.

Gas exchange was measured every week for 10 weeks on the 5th and 9th leaves with a portable photosynthesis apparatus (LI-6000). Leaf 9 was considered as a fully expanded leaf, acting as a source leaf, while leaf 5 was not fully expanded and was considered as an importer of carbohydrates. Leaves were counted from the top of the plant, starting with the first leaf $> 2\text{ cm}$. Photosynthetic rates were measured at the same CO₂ concentrations under which the plants were grown. The photosynthesis system monitored photosynthetic photon flux (PPF), relative humidity, leaf temperature, chamber air temperature and CO₂ concentration, initial internal leaf CO₂ concentration, stomatal conductance, and the net photosynthetic rate. The measurement period was 90 sec, with an air flow inside the 1-liter leaf chamber of 6 $\text{cm}^3\cdot\text{s}^{-1}$ and a leaf area of 28 cm^2 . Each point is the average of four series of four measurements taken randomly during the mid-day (1000 to 1400 HR) in each experimental unit.

Results and Discussion

Growth analysis. Plants grown under high CO₂ levels showed a higher relative growth rate (RGR) than the plants grown at ambient CO₂ for the first 2 weeks of treatment (Fig. 1). The difference was greater for *L. chmielewskii* (30% higher) than *L. esculentum* (18% higher) for these first 2 weeks. Thereafter, the RGR of CO₂-enriched plants declined faster than the control. The greatest decline was found for *L. chmielewskii* grown under high CO₂. After 7 weeks of treatment, the RGRs of the 900- and 330- $\mu\text{mol}\cdot\text{m}^{-3}$ -grown plants were identical for both species. After 10 weeks, the RGRs of CO₂-enriched plants were lower than the controls for *L. esculentum* and *L. chmielewskii*. Peet (1986) also observed higher RGRs for CO₂-enriched cu-

Fig. 1. Relative growth rate of *L. esculentum* (A) and *L. chmielewskii* (B) grown at 330 and 900 $\mu\text{mol}\cdot\text{m}^{-3}$ for 10 weeks. Each point represents the mean of four values $\pm$ SE.

cumber plants during the first 2 to 3 weeks. Thereafter, the RGR declined faster in CO₂-enriched plants. Mauney et al. (1978) reported that cotton, sorghum, soybean, and sunflower all had higher RGRs under CO₂ enrichment during the juvenile stage, but not during the mature stage.

Leaf areas were slightly greater for CO₂-enriched plants after 2 weeks (5.5% and 6.7% for *L. esculentum* and *L. chmielewskii*, respectively), but were similar thereafter (data not shown). The specific leaf weight (SLW) was higher under high CO₂ for all sampling dates (Fig. 2). After 2 weeks, the SLW had increased by 33% for *L. esculentum* and 27% for *L. chmielewskii*, whereas the corresponding values after 8 weeks were 51% and 53%. *L. chmielewskii* had a higher SLW at both CO₂ levels than *L. esculentum* (average of 18% higher). These results suggest that more photosynthates were stored per unit of leaf area in *L. chmielewskii* and at high CO₂. Sionit et al. (1981) and Imai and Murata (1976) previously showed that the increase in leaf area was not as important as the increase in leaf dry weight under high CO₂ levels.

Net assimilation rates (NAR) of the high-CO₂-grown plants were always higher than those of the ambient-grown plants (Fig. 3 A and B). After 2 weeks of treatment, NAR at 900 $\mu\text{mol}\cdot\text{m}^{-3}$ was 43% higher for *L. esculentum* and 58% for *L. chmielewskii* than for the controls. After 8 weeks, the NAR at high CO₂ was still significantly higher for *L. chmielewskii*, but not for *L. esculentum*. The increase in leaf dry weight (data not shown) relative to leaf area explains the higher NAR as compared to RGR under high CO₂ levels. These results suggest that assimilates were preferentially accumulated as reserves in the leaves rather than contributing to canopy enlargement. This particular partitioning, which resulted in thicker leaves under high CO₂ levels, was more evident in *L. chmielewskii* than *L. esculentum*.

L. esculentum plants grown under high CO₂ concentrations flowered 7 days earlier than those grown under ambient CO₂

Fig. 2. Specific leaf weight of *L. esculentum* (A) and *L. chmielewskii* (B) grown at 330 and 900 $\mu\text{mol}\cdot\text{m}^{-3}$ for 10 weeks. Each point represents the mean of four values $\pm$ SE.

Fig. 3. Net assimilation rate of *L. esculentum* (A) and *L. chmielewskii* (B) grown at 330 and 900 $\mu\text{mol}\cdot\text{m}^{-3}$ for 10 weeks. Each point represents the mean of four values $\pm$ SE.

(Table 1). The early (first 3 weeks of harvest) and total yields were increased by 80% and 21.5%. Slack (1986) reported that CO_2 enrichment increased total yield of greenhouse tomato by $\approx 30\%$. Calvert and Slack (1975) showed that the major effect of high CO_2 was on the earliness and pointed out that the beneficial effect of CO_2 was mainly on the first truss. They con-

Table 1. Flowering date and yields of *Lycopersicon esculentum* grown under 330 or 900 $\mu\text{mol}\cdot\text{m}^{-3}$ CO_2 for 10 weeks.^z

Criterion	CO_2 concn ($\mu\text{mol}\cdot\text{m}^{-3}$)		Increase (%)
	330	900	
Flowering date	4 Feb.	28 Jan.	
Early yields (kg/plant)	1.95 ± 0.31	3.51 ± 0.49	80
Total yields (kg/plant)	6.92 ± 0.95	8.41 ± 1.02	22
Number of fruits/plant	76.00 ± 8.41	81.00 ± 8.53	6.6
Average fruit weight (g)	91.05 ± 9.37	103.83 ± 9.89	14

^zEach value is a mean of eight plants $\pm$ SE.

cluded that the preferential increase of leaf thickness, as opposed to leaf expansion under high CO_2 , suggests that the extra assimilates were stored in the leaves rather than translocated into the developing fruits. Similarly, we found that high CO_2 increased leaf dry weight more than leaf area.

Gas exchange rate. The carbon exchange rates (CER) were significantly higher in CO_2 -enriched plants for the early sampling dates. After 1 week of treatment, CO_2 enrichment increased the CER of leaf 5 by 37% for *L. esculentum* and 38% for *L. chmielewskii* (Fig. 4 A and B). However, long-term growth at elevated CO_2 concentrations resulted in diminished CER (Fig. 4 C and D). The decrease was faster and more pronounced for *L. chmielewskii* for both leaves. After 6 weeks, no beneficial effect of CO_2 enrichment on CER was measured for either species. For both species, most of the decline in photosynthesis of CO_2 -enriched plants occurred during the first 6 weeks of treatment; thereafter, the differences stabilized. Aoki and Yabuki (1977) found that the decline of photosynthesis under CO_2 enrichment was faster in lower (older) leaves of cucumber plants. With tomato, we found a similar decline in both the younger (5th) and older (9th) leaves.

Stomatal conductance of leaves 5 and 9 of the two tomato species was lower for plants grown at 900 $\mu\text{mol}\cdot\text{m}^{-3}$ (average of 57% lower) than those grown at 330 $\mu\text{mol}\cdot\text{m}^{-3}$. The difference between the two CO_2 treatments gradually increased throughout the experiment (Fig. 5). Stomatal conductance of the 5th leaf of *L. chmielewskii* and *L. esculentum* decreased by 84% and 36% over the course of the experiment. In leaf 9, the average decline due to CO_2 enrichment was the same (54%) in both species. Our results agree with those reported by Peet et al. (1986), Havelka et al. (1984), and Spencer and Bowes (1986), who measured a significant decrease of stomatal conductance of leaves grown under high CO_2 concentration.

The stomatal conductance of high- CO_2 -grown plants showed the same trend of decline as the photosynthetic rates. These results suggest that the acclimation of plants to high CO_2 concentration may be attributed to a partial closure of the stomata. However, in contrast to photosynthesis and stomatal conductance, internal CO_2 levels were similar for both types of leaves grown in a given CO_2 concentration (Table 2). Since internal CO_2 was not reduced over the course of the experiment (data not shown), we believe that reduced stomatal conductance does not account for the decrease in photosynthesis seen in high- CO_2 -grown plants. Our results suggest that the decline of photosynthesis of high- CO_2 -grown plants results from a lower rate of use of the internal CO_2 . Spencer and Bowes (1986) also found a simultaneous decrease of photosynthesis and stomatal conductance, resulting in a constant internal CO_2 concentration during plant development. In tomatoes, even though the partial closure of the stomata did not affect the carbon exchange rate,

Fig. 4. Carbon exchange rates of two tomato species grown at 330 and 900 $\mu\text{mol}\cdot\text{m}^{-3}$ for 10 weeks. (A) Leaf 5 of *L. esculentum*. (B) Leaf 5 of *L. chmielewskii*. (C) Leaf 9 of *L. esculentum*. (D) Leaf 9 of *L. chmielewskii*. Each point represents the mean of four values $\pm$ SE.

Fig. 5. Stomatal conductance of two tomato species grown at 330 and 900 $\mu\text{mol}\cdot\text{m}^{-3}$ for 10 weeks. (A) Leaf 5 of *L. esculentum*. (B) Leaf 5 of *L. chmielewskii*. (C) Leaf 9 of *L. esculentum*. (D) Leaf 9 of *L. chmielewskii*. Each point represents the mean of four values $\pm$ SE.

CO_2 enrichment may have sufficiently reduced transpiration and lowered nutrient uptake (Madsen, 1975) to indirectly cause the acclimation of plants.

Leaves 5 and 9 responded similarly to high CO_2 enrichment throughout plant growth, which suggests that the loss of efficiency of CO_2 -enriched plants was not caused by tissue aging. Our data also demonstrated that leaf 5 was less efficient in 10-week-old plants than in 1-week-old plants, even though leaf 5 was young tissue (≈ 10 to 15 days old). Our results indicate that it is the duration of exposure of the whole plant to elevated CO_2 concentration, rather than the time of exposure of the tissue, that governs the acclimation to high CO_2 concentration.

Spencer and Bowes (1986) compared the effects of CO_2 enrichment on mature and immature tissues. Both mother and daughter plants of water hyacinth had the same temporary enhancement of photosynthesis when exposed to high CO_2 . Ac-

climation of these plants under high CO_2 level was more related to the time of exposure at high CO_2 than the age of the plant. In addition, our results suggest that the time of exposure of the mother plant to high CO_2 greatly influences the acclimation of the young plant tissues exposed to high CO_2 only for very short periods of time. The loss of photosynthetic efficiency of old leaves exposed for a long time to high CO_2 seems to be transmitted to young leaves exposed only for a short period.

As reported for many C_3 species, the beneficial effects of CO_2 enrichment on CER was most effective during short-term exposure. However, our results show that the photosynthetic efficiency was not maintained throughout plant growth. Consequently, the photosynthetic efficiency of CO_2 -enriched plants decreases more rapidly than the control. Both *L. esculentum* and *L. chmielewskii* acclimated to high CO_2 . These results suggest that the specific mechanisms involved in the sink metab-

Table 2. Internal CO₂ concentration of *Lycopersicon esculentum* and *L. chmielewskii* grown under two CO₂ concentrations.

Species	CO ₂ treatments ($\mu\text{mol}\cdot\text{m}^{-3}$)	Internal CO ₂ concn ^z ($\mu\text{mol}\cdot\text{m}^{-3}$)	
		Leaf 5	Leaf 9
<i>L. esculentum</i>	330	265 $\pm$ 7	257 $\pm$ 8
	900	766 $\pm$ 22	716 $\pm$ 31
<i>L. chmielewskii</i>	330	254 $\pm$ 7	261 $\pm$ 8
	900	701 $\pm$ 27	727 $\pm$ 26

^zEach value is a mean of four replications $\pm$ SE and the average of 10 weeks of experiment. Measurements were taken weekly.

olism and carbohydrate assimilation of *L. chmielewskii* cannot eliminate, nor attenuate, the long-term decline of photosynthesis of plants exposed to high CO₂ concentrations. However, even though tomato plants lost some of their photosynthetic efficiency under high CO₂ concentrations, early and total yields of *L. esculentum* were increased by more than 80% and 21%, respectively.

The results reported here demonstrate that stomatal conductance decreased significantly at high CO₂, while internal CO₂ remained constant throughout tomato plant growth. Consequently, the partial closure of stomata cannot explain the acclimation of plants to high CO₂ level. An increased mesophyll resistance rather than stomatal resistance seems a more likely explanation for the long-term decline of photosynthesis with extended use of high CO₂ concentration.

This study found that leaves 5 and 9 responded similarly to high CO₂ concentrations, which suggests that aging and/or differential sink : source ratios of the two leaves are not primary explanations of the long-term decline. In addition, this result implies that the time of exposure of the whole plants to high CO₂ concentration was more critical than the time of exposure of a specific tissue. The loss of photosynthetic efficiency that gradually appears as tomato plants were exposed to high CO₂ level seems to be transmitted to young tissue.

Further research is needed to verify if there is a relationship between leaf ontogeny and the time of exposure to high CO₂ concentrations. Long-term studies are required to evaluate if very young leaves become acclimated due to the acclimation of older leaves. Furthermore, fundamental research should be undertaken to specify the relationships among sugar metabolism, Rubisco activity, and the loss of photosynthetic efficiency to specify the biochemical and physiological reasons underlying the efficiency loss.

Literature Cited

Aoki, M. and K. Yabuki. 1977. Studies on the carbon dioxide enrichment for plant growth: VII. Changes in dry matter production and photosynthetic rate of cucumber during carbon dioxide enrichment. *Agr. Meteorol.* 18:475-485.

Calvert, A. and G. Slack. 1975. Effects of carbon dioxide enrichment on growth, development and yield of glasshouse tomatoes: 1. Response to controlled concentration. *J. Hort. Sci.* 50:61-71.

Clough, J. M., M.M. Peet, and P.J. Kramer. 1981. Effects of high atmospheric CO₂ and sink size rates of photosynthesis of a soybean cultivar. *Plant Physiol.* 67:1007-1010.

Dennis, D.J. 1980. Effects of carbon dioxide enrichment and temperature programme on the growth and yield of glasshouse cucumbers. *Acta Hort.* 118:205-220.

Frydich, J. 1976. Photosynthetic characteristics of cucumber seedlings

grown under two levels of carbon dioxide. *Photosynthetic* 10:335-338.

Gardner, F. P., R.B. Pearce, and R.L. Mitchell. 1985. Growth and development, p. 187-208. In: F.P. Gardner, R.B. Pearce and R.L. Mitchell (eds.). *Physiology of crop plants*, Iowa State Univ. Press, Ames.

Gates, D. M., B.R. Strain, and J.A. Weber. 1983. Ecophysiological effects of changing atmospheric CO₂ concentration, p. 503-526. In: O.L. Lange, P.S. Nobel, C.B. Osmond, and H. Ziegler (eds.). *Physiological plant ecology IV*, encyclopedia of plant physiology, New Ser. vol. 12 D. Springer-Verlag, Berlin.

Havelka, U. D., R.C. Ackerson, M.G. Boyle, and V.A. Wittenbach. 1984. CO₂ enrichment effects on soybean physiology: I. Effects of long-term CO₂ exposure. *Crop Sci.* 24:1146-1150.

Hicklenton, P.R. and P.A. Joliffe. 1980. Alterations in the physiology of CO₂ exchange in tomato plants grown in CO₂-enriched atmospheres. *Can. J. Bot.* 58:2181-2189.

Hurd, R.G. 1968. Effect of CO₂-enrichment on the growth of young plants in low light. *Ann. Bot.* 32:531-542.

Imai, K. and Y. Murata. 1976. Effect of carbon dioxide concentration on growth and dry matter production of crop plants. *Proc. Crop. Sci. Soc. Jpn.* 45:598.

Kimball, B.A. 1983. Carbon dioxide and agricultural yield: An assemblage and analysis of 430 prior observations. *Agron. J.* 75:779-788.

Knecht, G.N. and J.W. O'Leary. 1983. The influence of carbon dioxide on the growth, pigment, protein, carbohydrate, and mineral status of lettuce. *J. Plant Nutr.* 64:301-312.

Lemon, E.R. 1983. CO₂ and plants. The response of plants to rising levels of atmospheric carbon dioxide. *AAAS Selected Symp.* 84. AAAS, Washington, D.C.

Madsen, E. 1975. Effects of CO₂ enrichment on growth, development, fruit production and fruit quality of tomato plant from a physiological viewpoint. *Phytotronics* III:318-330.

Mauney, J. R., K.E. Fry, and G. Guinn. 1978. Relationship of photosynthetic rate to growth and fruiting of cotton, soybean, sorghum and sunflower. *Crop Sci.* 18:259-263.

Mortensen, L.M. 1983. Growth response of some greenhouse plants to environment: X. Long-term effect of CO₂-enrichment on photosynthesis, photorespiration, carbohydrate content and growth of *Chrysanthemum morifolium* Ramat. *Meld. Nor. Landbrukshogsk.* 62:1-11.

Mortensen, L.M. 1987. Review: CO₂ enrichment in greenhouses. *Crop responses*. *Scientia Hort.* 33:1-25.

Peet, M.M. 1986. Acclimation to high CO₂ in monoecious cucumbers: 1. Vegetative and reproductive growth. *Plant Physiol.* 80:59-62.

Peet, M. M., S. Huber and D.T. Patterson. 1986. Acclimation of high CO₂ in monoecious cucumbers: H. Carbon exchange rates, enzyme activities, and starch and nutrient concentrations. *Plant Physiol.* 80:63-67.

Sasek, T. W., E.H. DeLucia, and B.R. Strain. 1985. Reversibility of photosynthetic inhibition after long term exposure to elevated CO₂ concentrations. *Plant Physiol.* 78:619-622.

Sionit, N., B.R. Strain, and H. Hellmers. 1981. Effects of different concentrations of atmospheric CO₂ on growth and yield components of wheat. *J. Agr. Sci.* 79:335-339.

Slack, G. 1986. CO₂ enrichment of tomato crops, p. 151-163. In: H.Z. Enoch and B.A. Kimball (eds.). *Carbon dioxide enrichment of greenhouse crops*. vol. H. Physiology, yields, and economics, CRC Press, Boca Raton, Fla.

Spencer, W. and G. Bowes. 1986. Photosynthesis and growth of water hyacinth under CO₂ enrichment. *Plant Physiol.* 82:528-533.

Vu, C. V., L.H. Allen, Jr., and G. Bowes. 1983. Effects of light and elevated atmospheric CO₂ on the ribulose biphosphate carboxylase activity and ribulose biphosphate level of soybean leaves. *Plant Physiol.* 73:729-734.

Wong, S. C., I.R. Cowan, and G.D. Farquhar. 1979. Stomatal conductance correlates with photosynthetic capacity. *Nature (London)* 282:424-426.

Yelle, S., A. Gosselin, and M.J. Trudel. 1987. Effect of atmospheric

CO₂ concentration and root-zone temperature on growth, mineral nutrition and nitrate reductase activity of greenhouse tomato. J. Amer. Soc. Hort. Sci. 112(6):1036-1040.

Yelle, S., J.D. Hewitt, N.L. Robinson, S. Damon, and A.B. Bennett. 1988. Sink metabolism in tomato fruit: III. Analysis of carbohydrate assimilation in a wild species. Plant Physiol. 87:737-740.