Regulating Vegetative Growth and Flowering with Gibberellic Acid in Intact Plants and Cultured Phylloclades of 'Crimson Giant' Easter Cactus

Thomas H. Boyle¹, Michael Marcotrigiano¹, and Suzanne M. Hamlin²

Department of Plant and Soil Sciences, French Hall, University of Massachusetts, Amherst, MA 01003

Additional index words. flowering potted plant, GA₃, *Hatiora gaertneri*, ornamental plant, *Rhipsalidopsis gaertneri*

Abstract. Investigations were performed to determine the influence of gibberellic acid (GA₃) on intact plants and cultured phylloclades of 'Crimson Giant' Easter cactus [*Rhipsalidopsis gaertneri* (Regel) Moran]. Responses of intact plants depended on GA₃ concentration, number of spray applications, and application time. Single GA₃ applications delayed flowering and reduced the percentage of apical phylloclades flowering and number of flower buds per plant when applied before floral primordia formation [from 20 days before to the start of long days (LDs)], but hastened flowering and did not affect the percentage of apical phylloclades flowering or number of flower buds per plant when applied during floral bud development (20 days after the start of LDs). When sprays were applied at or before the start of LDs, increasing the GA₃ concentration resulted in fewer plants flowering, longer flowering delays, and further decreases in the number of flower buds per plant. Multiple GA₃ applications were more inhibitory to flowering than single applications. Whole plants and cultured phylloclades exhibited similar reactions to GA₃, but cultured phylloclades were more responsive to GA₃ than intact plants. Intact plants and cultured phylloclades generally produced more new phylloclades as GA₃ concentration increased. Spine growth also increased when phylloclades were cultured in a GA₃-containing medium. Flowering was accelerated by ≈5 days when GA₃ was applied to intact plants with 1- to 2-mm-long flower buds. GA₃ may be horticulturally useful for Easter cactus crop scheduling.

Easter cactus—currently *Rhipsalidopsis gaertneri* (Regel) Moran (Liberty Hyde Bailey Hortorium, 1976), but proposed by Barthlott (1987) as *Hatiora gaertneri* (Regel) Barthlott—is an epiphytic shrub with a determinate growth pattern that produces a series of flattened, two-ribbed joints (phylloclades). Areoles (modified axillary buds) are restricted to the margins and apexes of phylloclades (Liberty Hyde Bailey Hortorium, 1976). Areoles that develop below the phylloclade apex are solitary, whereas, at the apex, areoles are crowded together to form a composite areole (Barthlott, 1987). Flowers and new phylloclades develop almost invariably from composite areoles on apical phylloclades and infrequently from subapical areoles or subapical phylloclades (Boyle, 1992; Liberty Hyde Bailey Hortorium, 1976).

Areoles are a prominent anatomical feature of cacti, and each areole contains several spine primordia and an areolar meristem (Boke, 1944). Vegetative shoots (phylloclades), leaves, spines, flowers, and roots arise through the activity of the areolar meristem (Gibson and Nobel, 1986). Mauseth and Halperin (1975) were able to regulate the developmental pattern in cultured areoles of *Opuntia polyacantha* Haw. by varying the proportions of GA₃, cytokinin, and auxin in the medium. These results demonstrate that exogenously applied GA₃ may strongly influence the developmental fate of the areolar meristem.

Exogenously applied GA has modified growth and development in a wide range of plants. Growth of stems and other organs is promoted by GA and results from enhanced cell division, increased carbohydrate hydrolysis, and increased cell-wall plasticity (Sachs, 1961; Salisbury and Ross, 1978). GA also regulates or promotes flowering in several species. GA induces many long-

day (LD) plants to flower under noninductive short days (SDS) (Lang and Reinhard, 1961; Wittwer and Bukovac, 1958) and substitutes for the LD phase in several dual-photoperiod species, including long-short-day plants (LSDPs) (Bünsow and Harder, 1956) and short-long-day plants (SLDPs) (Chouard, 1960; Ketellapper and Barbaro, 1966). In short-day plants (SDPs), GA applied during inductive photoperiods generally inhibits or delays floral initiation (Vince-Prue, 1975), but GA may accelerate floral development if applied after floral initiation (Greulach and Haesloop, 1958; Stuart and Cathey, 1962). In the SDP *Schlumbergera truncata* (Haw.) Moran (Cactaceae), GA₃ inhibits flower bud formation and delays flowering when applied before or at the beginning of SDS, accelerates flowering by up to 2 weeks when applied at the visible bud stage (buds 1 to 3 mm long), and has no or little influence when applied in the late flower development stage (buds 6 to 8 mm long) (Fujihara, 1959; Halevy and Rudich, 1968; Ho et al., 1985; Runger, 1984). In a few SDPs, GA has promoted floral initiation when applied to plants under noninductive LDs (Nanda et al., 1967; Wittwer and Bukovac, 1958), but GA cannot substitute for the SD requirement in most SDPs. Further, GA₃ cannot replace the SD phase in several dual-photoperiod species (Ketellapper and Barbaro, 1966; Penner, 1960; Wellensiek, 1960).

Limited information has been published on the responses of *R. gaertneri* to growth regulators (Boyle, 1992; Boyle et al., 1988; Kaukovirta, 1979), and we are unaware of any published reports on the responses of *R. gaertneri* to GA₃. The purpose of this research was to determine the morphological responses of intact plants and cultured phylloclades of the SLDP *R. gaertneri* 'Crimson Giant' to application time, number of applications, and concentration of exogenously applied GA₃.

Materials and Methods

General procedures. Plants were propagated and grown in glasshouses at the Univ. of Massachusetts, Amherst (lat. 42°22.5'N).

Received for publication 30 Dec. 1992. Accepted for publication 11 May 1993. Publication no. 3086 of the Massachusetts Agricultural Experiment Station. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Associate professor.

²Undergraduate student.

Plants were fertilized weekly with 20N-4.3P-16.6K (12% NO₃-N, 8% NH₄-N) at 200 mg N/liter. Fertilization began after phylloclades were rooted and was discontinued 4 weeks before the start of LDs to hasten apical phylloclade maturation. Four weeks after start of LDs, fertilization was resumed and applied at 200 mg/N per liter every 2 weeks until the experiments ended

Thermostat setpoints were 18/20C (heat-vent) during all experiments. Photosynthetic photon flux (PPF) and air temperatures were monitored with a datalogger (model LI- 1000) equipped with a quantum sensor (model LI- 190SA) and an aspirated thermistor (model LI-1000- 16; LI-COR, Lincoln, Neb.). The datalogger was configured with a sampling interval of 60 sec and recorded mean PPFs and temperatures at 1-h intervals. GA₃ was applied to intact plants (Expts. 1 and 2) near midday (1200 HR); temperature, PPF, and natural photoperiod on GA₃ application dates are provided in Table 1.

Multiple GA₃ applications on intact plants (Expt. 1). Whole phylloclades were propagated on 28 Apr. 1987 in 72-cell plastic trays using one phylloclade per 35-cm³ cell. The propagation medium was a commercial soilless mix composed of sphagnum peat, perlite, and vermiculite (Fafard Mix no. 2, Conrad Fafard, Springfield, Mass.). Rooted phylloclades were transplanted singly into 520-cm³ (10-cm-diameter) plastic pots containing Fafard Mix no. 2. Plants received natural daylengths (NDs) from propagation until 10 Jan. 1988. LDs were provided from 10 Jan. until the experiment ended by supplementing ND with incandescent irradiation at 3 μmol·m⁻²·s⁻¹ (400 to 700 nm) from 1600 to 2200 HR. Actual glasshouse temperatures were 18 ± 1.5C nights/20 ± 2C days and ranged from 16 to 26C.

Treatments included an unsprayed control and GA₃ (Pro-Gibb; Abbott Labs, North Chicago) at 25 or 50 mg·liter⁻¹ in one, two, three, or four applications. GA₃ solutions contained Tween 20 (polyoxyethylene sorbitan monolaurate) at 1 ml·liter⁻¹ as a surfactant and were applied to plants with a hand sprayer at ≈ 15 ml/plant. The initial treatment was applied on 10 Jan.; the other abdication dates were 17, 24, and 31 Jan. The experiment was a completely randomized design with eight pots per treatment.

Data were collected on days to flowering from the start of LDs (10 Jan.) to full expansion of the first flower, number of flower buds per flowering apical phylloclade, number of flower buds per plant, and percentage of apical phylloclades flowering. The experiment ended on 20 Apr. 1988. Before statistical analysis, the numbers of flower buds per flowering apical phylloclade and new apical phylloclades per plant were log-transformed [$\log_{10}(X + 1)$] and percentage values were arcsin-transformed. All variables were

analyzed by SAS's (1985) General Linear Model (GLM) procedure. Single degree-of-freedom contrasts were used to test for treatment differences.

Single GA₃ applications on intact plants (Expt. 2). Whole phylloclades were propagated on 27 Mar. 1988 using procedures and materials described in Expt. 1. Rooted phylloclades were transplanted singly into 520-cm³ plastic pots containing Fafard Mix no. 2. Plants were grown under ND from propagation until 31 Jan. 1989, and were then given LDs (as described in Expt. 1) until the experiment ended. Actual glasshouse temperatures were 18 ± 1.5C nights/21 ± 2C days and ranged from 16 to 28C. Treatments included GA₃ (Pro-Gibb) at 5, 50, and 500 mg·liter⁻¹ and an unsprayed control. GA₃ solutions contained Tween 20 at 1 ml·liter⁻¹ and were applied with a hand sprayer at ≈ 15 ml/plant. Treatments were applied on 11 Jan. (20 days before the start of LDs), 21 Jan. (10 days before the start of LDs), 31 Jan. (start of LDs), 10 Feb. (10 days after start of LDs), and 20 Feb. (20 days after the start of LDs). The experiment was a completely randomized design with eight pots per treatment.

At the start of LDs, 24 mature apical phylloclades were removed from eight supplementary plants (three phylloclades per plant) that were the same age and maintained under the same conditions as the controls. Apexes were dissected under a stereomicroscope at ×50 magnification to determine the developmental stage of the areolar meristems in the composite areole. Data were also collected on the presence of visible flower buds at each application date, flower bud length, days to flowering from the start of LDs (31 Jan.) to full expansion of the first flower, number of flower buds per plant, number of new apical phylloclades per plant, percentage of apical phylloclades flowering, and percentage of apical phylloclades with new phylloclades. The experiment ended on 1 May 1989. Chi-square tests were used to test for differences in percentage of flowering plants between GA₃ treatments and controls. Other variables were analyzed by SAS's (1985) GLM procedure. Percentage variables were arcsin-transformed and numbers of flower buds and new apical phylloclades per plant were log-transformed before analysis. Single degree-of-freedom contrasts were used to test for treatment differences.

In vitro responses of phylloclades cultured on GA₃-containing media (Expt. 3). Mature apical phylloclades were harvested on 21 Feb. 1990 from glasshouse-grown stock plants. Before harvest, stock plants were exposed to 8 weeks of 8-h SDs (0800 to 1600 HR) to satisfy the SD phase for flowering (Boyle, 1991). Surface contamination was reduced by pruning the bristly spines on the composite areole with scissors. Phylloclades were surface-disinfected by im-

Table 1. Glasshouse environmental conditions recorded on GA₃ application dates.

Experiment	GA ₃ application date	Air temp (°C) ^z	Photosynthetic photon flux (μmol·m ⁻² ·s ⁻¹) ^z	Natural photoperiod ^y	
				h	min
1	10 Jan. 1988	21.5 ± 0.1	442 ± 44	9	20
	17 Jan. 1988	20.1 ± 0.1	389 ± 24	9	31
	24 Jan. 1988	21.6 ± 0.1	136 ± 26	9	44
	31 Jan. 1988	21.4 ± 0.3	607 ± 66	10	0
2	11 Jan. 1989	18.9 ± 0.5	153 ± 20	9	21
	21 Jan. 1989	17.8 ± 1.3	422 ± 107	9	39
	31 Jan. 1989	18.5 ± 0.7	259 ± 30	10	0
	10 Feb. 1989	18.4 ± 0.7	373 ± 79	10	23
	20 Feb. 1989	19.0 ± 0.1	292 ± 71	10	50

^zActual glasshouse air temperatures and photosynthetic photon flux were collected by a datalogger configured with a sampling interval of 60 sec and recording mean values at 1-h intervals. Data are averages of hourly means (±SE) for 1000 to 1359 HR.

^yDuration of daylight (interval between sunrise and sunset) at lat. 42°N. Data are from List (1951).

mersing them for 25 min in a 20% (by volume) bleach solution (1.05% NaOCl) containing 0.8% Tween 20 as a surfactant, followed by three rinses in sterile distilled water. Explants were obtained by excising a ≈ 1.5 -cm-long, wedge-shaped piece from the phylloclade apex, which included the entire composite areole. Explants were oriented vertically in 55-cm³ (25-mm-diameter) vials containing 10 ml of sterile Murashige and Skoog (1962) medium that was solidified with 7 g Difco Bacto-agar/liter and supplemented with GA₃ at 0, 0.1, 1, 10, or 100 mg-liter⁻¹. Crystalline GA₃ (Grade III; Sigma, St. Louis) was dissolved in ethanol, and, to preserve GA₃ activity, stock solutions were filter-sterilized and added after autoclaving the medium (van Bragt and Pierik, 1971). The experiment was a completely randomized design with 10 replications (vials) per treatment.

Cultures were maintained in a growth room maintained at $25 \pm 1.5^\circ\text{C}$. Irradiance was provided by cool-white fluorescent lamps ($\approx 35 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPF), and a 16-h (LD) photoperiod was maintained to induce flowering in the cultured phylloclades (Boyle, 1991). After 35 days in vitro, data were collected on the numbers of flower buds and secondary (2°) phylloclades (those developing from the composite areole), and the presence of spine growth on the composite areole. The experiment was repeated on 12 Mar. 1990 using mature apical phylloclades collected from the stock plants previously described.

Percentage data were analyzed by single degree-of-freedom chi-square tests. Two variables (flower buds per explant and 2° phylloclades per explant) were log-transformed and analyzed by SAS's (1985) GLM procedure using data combined from both experiments. Homogeneity of error variances was previously verified using a two-tailed F test (Gomez and Gomez, 1984).

Results and Discussion

Experiment 1. All plants produced flowers, regardless of GA₃

concentration or number of applications. Compared to unsprayed plants, flowering was delayed and the percentage of apical phylloclades flowering and number of flower buds per plant was decreased in plants sprayed with GA₃ (Table 2). Increasing the number of GA₃ applications resulted in a linear increase in the number of days to flowering and linear decreases in the percentage of apical phylloclades flowering and number of flower buds per plant. With GA₃ at 25 mg-liter⁻¹, the number of flower buds per flowering apical phylloclade was not affected by number of applications, but at 50 mg-liter⁻¹, the number of flower buds per flowering apical phylloclade decreased linearly as the number of applications increased. Generally, plants treated with GA₃ at 50 mg-liter⁻¹ produced fewer flowers and flowered later than plants treated with 25 mg-liter⁻¹.

Experiment 2. At the start of LDs, dome-shaped areolar meristems (150 to 175 μm in diameter) and a few immature phylloclades ($< 800 \mu\text{m}$ long) were observed in the composite areoles of apical phylloclades collected from the supplementary plants. None of the composite areoles that were examined contained floral primordia. No flower buds were visible on plants when GA₃ was applied from 20 days before to 10 days after start of LDs. At 20 days after start of LDs, flower buds were clearly visible on untreated plants and were a maximum of 1 to 2 mm long.

Flowering was either inhibited, unaffected, or accelerated by GA₃, depending on concentration and application time (Table 3). Flowering was inhibited when GA₃ at 50 or 500 mg-liter⁻¹ was applied at or before the start of LDs. Generally, one application of GA₃ at 500 mg-liter⁻¹ resulted in fewer plants flowering, greater delays in flowering, and fewer flower buds compared to a single application of GA₃ at 50 mg-liter⁻¹. Applying GA₃ 10 days after the start of LDs did not affect days to flowering, but reduced the percentage of apical phylloclades flowering and number of flower buds per plant. When GA₃ was applied 20 days after the start of LDs, flowering was hastened but the percentage of apical phyllo-

Table 2. Influence of GA₃ concentration and number of applications on flowering of 'Crimson Giant' Easter cactus (Expt. 1).

GA ₃ concn (mg-liter ⁻¹)	No. of applications	Days to flowering ^a	Apical phylloclades flowering (%)	Flower buds/ flowering apical phylloclade (no.)	Flower buds/ plant (no.)
0 (control)	---	57	67	1.7	25.3
25	1	62	54	1.5	19.9
	2	69	49	1.6	21.9
	3	69	42	1.4	16.9
	4	80	28	1.5	11.5
	1	64	56	1.6	25.1
50	2	73	35	1.9	17.8
	3	86	31	1.3	10.6
	4	86	10	1.2	4.0
Contrasts					
Control vs. GA ₃		***	***	NS	**
Control vs. 25 mg-liter ⁻¹		***	***	NS	**
Control vs. 50 mg-liter ⁻¹		***	***	NS	**
25 mg-liter ⁻¹ vs. 50 mg-liter ⁻¹		***	***	NS	***
25 mg-liter ⁻¹ , no. applications linear		**	***	NS	*
25 mg-liter ⁻¹ , no. applications quadratic		NS	NS	NS	NS
50 mg-liter ⁻¹ , no. applications linear		***	***	*	***
50 mg-liter ⁻¹ , no. applications quadratic		NS	NS	NS	NS

^aDays to flowering from start of long days (10 Jan. 1988) for all treatments.

NS, *, **, *** Nonsignificant or significant at $0.05 \geq \alpha > 0.01$, $0.01 \geq \alpha > 0.001$, or $\alpha \leq 0.001$, respectively, according to F test of contrast between treatments.

Table 3. Influence of GA₃ concentration and application time on flowering and vegetative growth of 'Crimson Giant' Easter cactus (Expt. 2).

GA ₃ concn (mg·liter ⁻¹)	Application time ^z	Plants flowering (%)	Days to flowering ^y	Apical phylloclades flowering (%)	Flower buds/ plant (no.)	Apical phylloclades with new phylloclades (%)	New apical phylloclades/ plant (no.)
0 (control)	---	100	60	82	25.3	4	0.6
5	-20	100	61	75	16.4	2	0.3
	-10	100	64	76	24.5	0	0
	0	100	62	63	17.0	9	1.8
	10	100	58	75	20.4	5	1.4
	20	100	55	82	24.8	1	0.3
50	-20	100	73	31	6.6	10	2.5
	-10	100	83	25	7.0	22	6.5
	0	50 ^x	85	6	1.5	22	7.0
	10	100	56	42	11.1	5	1.0
	20	100	56	68	19.4	3	0.6
500	-20	13 ^x	92	1	0.1	32	10.5
	-10	0 ^x	---	0	0	33	10.4
	0	13 ^x	77	1	0.3	22	5.6
	10	100	58	42	7.9	4	0.6
	20	100	55	75	22.8	2	0.5
Contrasts							
Control vs. GA ₃ , day -20			***	***	***	*	*
Control vs. GA ₃ , day -10			***	***	***	**	***
Control vs. GA ₃ , day 0			***	***	***	**	**
Control vs. GA ₃ , day 10			NS	***	***	NS	NS
Control vs. GA ₃ , day 20			*	NS	NS	NS	NS
Control vs. 5 mg·liter ⁻¹			NS	NS	NS	NS	NS
Control vs. 50 mg·liter ⁻¹			***	***	***	NS	***
Control vs. 500 mg·liter ⁻¹			w	***	***	**	***
50 mg·liter ⁻¹ vs. 500 mg·liter ⁻¹			w	***	***	*	*
5 mg·liter ⁻¹ , date linear			***	NS	NS	NS	NS
5 mg·liter ⁻¹ , date quadratic			*	*	NS	NS	NS
50 mg·liter ⁻¹ , date linear			***	***	***	*	**
50 mg·liter ⁻¹ , date quadratic			***	***	***	**	**
500 mg·liter ⁻¹ , date linear			w	***	***	***	***
500 mg·liter ⁻¹ , date quadratic			w	***	***	NS	NS

^zDays relative to before (–) or after the start of long days.^yDays to flowering from start of long days (31 Jan. 1989) for all treatments.^xTreatment significantly different from control by chi-square test, $P \leq 0.05$; chi-square = 3.841 for 1 df.^wContrast not estimable due to missing values or treatments.NS, *, **, *** Nonsignificant or significant at $0.05 \geq \alpha > 0.01$, $0.01 \geq \alpha > 0.001$, or $\alpha \leq 0.001$, respectively, according to F test of contrast between treatments.

clades flowering and number of flower buds per plant were not affected. GA₃ at 5 mg·liter⁻¹ generally did not affect vegetative growth or flowering, regardless of application time, but was as effective as 50 or 500 mg·liter⁻¹ in accelerating flowering when applied 20 days after the start of LDs.

GA₃-treated plants generally produced more new phylloclades than the controls (Table 3). There was a highly significant negative correlation between the number of flower buds per plant and new apical phylloclades per plant ($r = -0.83$, $P < 0.001$ under H_0 : $\rho = 0$ ($n=14$)). There was also a highly significant negative correlation between the percentage of apical phylloclades flowering and the percentage of apical phylloclades with new phylloclades ($r = -0.91$, $P < 0.001$ under H_0 : $\rho = 0$ ($n = 14$)). The number of new phylloclades depended on GA₃ concentration and application time. Applying GA₃ from 20 days before to the start of LDs resulted in a higher percentage of apical phylloclades with new phylloclades and more new apical phylloclades per plant compared with the controls (Table 3). There were no differences

between controls and plants treated with GA₃ at 5 mg·liter⁻¹ in the percentage of apical phylloclades with new phylloclades or the number of new apical phylloclades per plant. However, increasing GA₃ from 50 to 500 mg·liter⁻¹ resulted in a higher percentage of apical phylloclades with new phylloclades and more new apical phylloclades per plant.

The morphology of flowers and phylloclades was altered by GA₃. Relative to the controls, new phylloclades and flowers on plants treated with GA₃ at 50 or 500 mg·liter⁻¹ were thinner and elongated (data not presented). GA₃ at 5 mg·liter⁻¹ did not affect the morphology of new phylloclades, but some flowers were elongated slightly when GA₃ at 5 mg·liter⁻¹ was applied 20 days after starting LDs.

Flowering responses of intact plants depended on GA₃ concentration, application time, and number of applications. A single GA₃ application delayed flowering and reduced the percentage of apical phylloclades flowering and number of flower buds per plant when applied to plants before the formation of floral primordia,

i.e., from 20 days before to the start of LDs, but hastened flowering and did not affect the percentage of apical phylloclades flowering or number of flower buds per plant when applied to plants during floral bud development, i.e., 20 days after starting LDs (Table 3). Increasing the concentration always resulted in longer flowering delays and fewer flowers when GA₃ was applied at or before starting LDs. In addition, multiple GA₃ applications were more inhibitory than single applications and resulted in fewer flowers and longer flowering delays (Table 2). Similar results were obtained with the SDP *S. truncata* (Fujihara, 1959; Halevy and Rudich, 1968; Ho et al., 1985; Runger, 1984). *Rhipsalidopsis gaertneri* and *S. truncata* are remarkably similar in their responses to GA₃, even though *R. gaertneri* is a SLDP at 18 to 22°C (Boyle et al., 1988; Boyle, 1991) whereas *S. truncata* is a SDP at 15 to 21°C (Runger and Poole, 1985). The results of the current experiments and those with *S. truncata* demonstrate that GA₃ may inhibit or promote flowering, depending on the stage of plant development when applied.

Experiment 3. Contaminated cultures (13 in total) were discarded before data collection. Statistical analysis was performed on data collected from 87 uncontaminated cultures.

For number of flower buds per explant and 2° phylloclades per explant, there were no significant differences between the two experiments (21 Feb. and 12 Mar. 1990) nor was there a significant experiment × GA₃ interaction (Table 4). Both variables, however, were significantly affected by GA₃ concentration. The number of flower buds decreased linearly as GA₃ concentration increased,

and the percentage of explants with flower buds was significantly lower among explants cultured in GA₃-amended medium compared to those in GA₃-free medium (Table 5). The percentage of explants with 2° phylloclades and the percentage of explants with spine growth were significantly higher among those cultured in GA₃-amended medium than in those in GA₃-free medium. GA₃ increased the number of 2° phylloclades compared to the controls, but the response to GA₃ concentration was nonlinear. The number of 2° phylloclades increased as GA₃ concentration increased from 0 to 1 mg·liter⁻¹, and then decreased at concentrations >1 mg·liter⁻¹. The correlation between numbers of flower buds per explant and 2° phylloclades per explant was negative but not significant ($r = -0.71$, $P > 0.05$ under H_0 : $\rho = 0$ ($n = 5$)). Lack of significance may have been due to the inhibition of flower and phylloclade formation when explants were cultured on medium containing GA₃ at 100 mg·liter⁻¹ (Table 5).

Whole plants and cultured phylloclades exhibited similar reactions to GA₃, but cultured phylloclades were more responsive to GA₃ than intact plants (Tables 2, 3, and 5). GA₃ at 5 mg·liter⁻¹ generally did not affect the vegetative or reproductive growth of intact plants (Table 3), whereas GA₃ at 0.1 mg·liter⁻¹ inhibited flowering and promoted vegetative growth in cultured phylloclades (Table 5). Mauseth and Halperin (1975) also reported that GA₃ at concentrations as low as 0.01 mg·liter⁻¹ elicited responses in quiescent buds of *O. polyacantha* that were cultured in vitro. In Expt. 2, the maximum number of new phylloclades was obtained when plants were treated with GA₃ at 500 mg·liter⁻¹ (Table 3); in

Table 4. Analysis of variance for flower bud and phylloclade development for cultured phylloclades of 'Crimson Giant' Easter cactus (Expt. 3).

Source of variation	df	Mean squares	
		Flower buds/ explant (no.)	2° Phylloclades/ explant (no.)
Experiment (E) ^z	1	0.0132	0.0055
Replications within E	18	0.0096	0.0297
GA ₃ concentration (GA ₃)	4	0.5584***	0.2900***
E × GA ₃	4	0.0112	0.0142
Pooled Error	59	0.0181	0.0421

^zF test performed using replications within E as the error term.

***Significant at $P \leq 0.001$.

Table 5. Influence of GA₃ concentration on flowering and vegetative growth of cultured phylloclades of 'Crimson Giant' Easter cactus (Expt. 3).

GA ₃ concn (mg·liter ⁻¹)	Explants with flower buds (%)	Flower buds/ explant (no.)	Explants with 2° phylloclades (%)	2° Phylloclades/ explant (no.)	Explants with spine growth (%)
0 (control)	100	1.9	0	0	0
0.1	47 ^z	0.5	33 ^z	0.6	60 ^z
1	11 ^z	0.2	68 ^z	1.4	100 ^z
10	22 ^z	0.2	61 ^z	1.1	100 ^z
100	12 ^z	0.1	29 ^z	0.4	100 ^z
Contrasts					
Control vs. GA ₃		***		***	
GA ₃ linear		***		NS	
GA ₃ quadratic		NS		NS	

^zTreatment significantly different from control by chi-square test, $P \leq 0.05$; chi-square = 3.841 for 1 df.

NS,*,**,*NSignificant or significant at $0.05 \geq \alpha > 0.01$, $0.01 \geq \alpha > 0.001$, or $\alpha \leq 0.001$, respectively, according to F test of contrast between treatments.

Expt. 3, the number of 2° phylloclades was maximal when explants were cultured on media containing GA₃ at 1 mg-liter⁻¹, and concentrations >1 mg-liter⁻¹ inhibited phylloclade formation (Table 5). Differences between cultured phylloclades and intact plants in responsiveness to GA₃ may have been due to greater hormone uptake in the former, possibly due to direct contact between the GA₃-containing medium and the wounded surface of the explant.

In Expt. 3, old spines were shorter (due to trimming) and paler (due to bleaching) than new spines; as a consequence, spine growth on the composite areole was evaluated readily. Spine growth increased when phylloclades were cultured in medium containing GA₃ at >0.01 mg-liter⁻¹ (Table 5). GA₃ also increased the growth of spines in *Chamaecereus silvestri* (Speg.) Britt. & Rose (Sanderson et al., 1986), *Mammillaria elongata* DC. (Sanderson et al., 1986), and *O. microdasys* (Lehm.) Pfeiff. (White et al., 1978). Exogenously applied GA₃ may have induced spine growth in 'Crimson Giant' by triggering the initiation of new spines, as reported by Mauseth and Halperin (1975) and Mauseth (1977) for *O. polyacantha*. Cactus spines are presumed to be evolutionarily derived from leaves (Boke, 1944), and since directly applying GA₃ can stimulate leaf growth (Aloni and Pressman, 1980; Gray, 1957), it is feasible that GA₃ may also promote spine growth by increasing cell division or elongation or both within existing spine primordia. Additional research is needed to discern the mechanisms responsible for GA₃-induced spine growth in 'Crimson Giant'.

Intact plants and cultured phylloclades generally produced more new phylloclades as GA₃ concentration increased, i.e., the opposite trend observed for number of flower buds (Tables 3 and 5). This inverse relationship between vegetative growth and flowering is demonstrated by the highly significant negative correlation between the numbers of flower buds per plant and new apical phylloclades per plant ($r = -0.83$). Phylloclade growth of 'Crimson Giant' may have been directly and indirectly affected by GA₃. Direct stimulation of stem growth via increased cell division and cell elongation has been demonstrated in several species after GA₃ was applied (Sachs, 1961), and GA₃ would be expected to have similar effects on phylloclade growth. Also, GA₃ may have indirectly affected phylloclade growth by inhibiting flowering. Profuse flowering and a near absence of new phylloclades was the normal pattern of development, as exhibited by the controls (Tables 2 and 3). The flowering process may repress vegetative growth, and GA₃ may have reversed the repression of vegetative growth by inhibiting flowering, thus allowing phylloclades to form. GA₃ also promoted shoot growth in intact plants of *C. silvestri* (Sanderson et al., 1986), *M. elongata* (Sanderson et al., 1986), and *S. truncata* (Yonemura, 1979).

Horticultural significance. Applying GA₃ from 20 days before to 10 days after the start of LDs was either ineffective or inhibitory to flowering and, thus, would have no horticultural value for producing Easter cactus. However, a single application of GA₃ at 5 mg-liter⁻¹ accelerated flowering by 5 days when applied to plants with 1- to 2-mm-long flower buds. Accelerating flowering reduces the cropping period and may aid in Easter cactus crop scheduling. GA₃ at concentrations >5 mg-liter⁻¹ will also accelerate flowering, but may decrease plant quality and marketability due to deleterious effects on phylloclade and flower morphology.

Literature Cited

- Aloni, B. and E. Pressman. 1980. Interaction with salinity of GA₃ induced leaf elongation, petiole pithiness and bolting of celery. *Scientia Hort.* 13:135-142.
 Barthlott, W. 1987. New names in Rhipsalidinae (Cactaceae). *Bradleya* 5:97-100.

- Boke, N.H. 1944. Histogenesis of the leaf and areole in *Opuntia cylindrica*. *Amer. J. Bot.* 31:299-316.
 Boyle, T.H. 1991. Temperature and photoperiodic regulation of flowering in 'Crimson Giant' Easter cactus. *J. Amer. Soc. Hort. Sci.* 116:618-622.
 Boyle, T.H. 1992. Modification of plant architecture in 'Crimson Giant' Easter cactus with benzyladenine. *J. Amer. Soc. Hort. Sci.* 117:584-589.
 Boyle, T.H., D.J. Jacques, and D.P. Stimart. 1988. Influence of photoperiod and growth regulators on flowering of *Rhipsalidopsis gaertneri*. *J. Amer. Soc. Hort. Sci.* 113:75-78.
 Bünsow, R. and R. Harder. 1956. Blütenbildung von *Bryophyllum* durch Gibberellin. *Naturwissenschaften* 43:479-480.
 Chouard, P. 1960. Vernalization and its relations to dormancy. *Annu. Rev. Plant Physiol.* 11:191-238.
 Fujihara, K. 1959. Effect of gibberellin on the flowering of Christmas cactus. I. Effect of gibberellin given at various times prior to or after the start of short days. *J. Hort. Assn. Jpn.* 28:200-208.
 Gibson, A.C. and P.S. Nobel. 1986. The cactus primer. Harvard Univ., Cambridge, Mass.
 Gomez, K.A. and A.A. Gomez. 1984. Statistical procedures for agricultural research. 2nd ed. Wiley, New York. p. 187-240, 272-315.
 Gray, R.A. 1957. Alteration of leaf size and shape and other changes caused by gibberellins in plants. *Amer. J. Bot.* 44:674-682.
 Greulach, V.A. and J.G. Haesloop. 1958. Influence of gibberellin on *Xanthium* flowering as related to number of photoinductive cycles. *Science* 127:646-647.
 Halevy, A.H. and Y. Rudich. 1968. Effect of short-day and gibberellin on flowering in *Zygocactus*. *Symp. Plant Production Containers, Intl. Soc. Hort. Sci., Copenhagen, Denmark.*
 Ho, Y.S., K.C. Sanderson, and J.C. Williams. 1985. Effect of chemicals and photoperiod on the growth and flowering of Thanksgiving cactus. *J. Amer. Soc. Hort. Sci.* 110:658-662.
 Kaukovirta, E. 1979. The effect of ethephon and chlormequat on flowering in *Rhipsalidopsis gaertneri* and the *Zygocactus* hybrid, "Weihnachtsfreude". *Acta Hort.* 91:419-424.
 Ketellapper, H.J., and A. Barbaro. 1966. The role of photoperiod, vernalization and gibberellic acid in floral induction in *Coreopsis grandiflora* Nutt. *Phyton (Buenos Aires)* 23:33-41.
 Lang, A. and E. Reinhard. 1961. Gibberellins and flower formation. *Adv. Chem.* 28:71-79.
 Liberty Hyde Bailey Hortorium. 1976. *Hortus third: A concise dictionary of plants cultivated in the United States and Canada.* MacMillan, New York. p. 947-948.
 List, R.J. 1951. Smithsonian meteorological tables. 6th ed. Smithsonian Inst., Washington, D.C. p. 506-512.
 Mauseth, J.D. 1977. Cytokinin- and gibberellin acid-induced effects on the determination and morphogenesis of leaf primordia in *Opuntia polyacantha* (Cactaceae). *Amer. J. Bot.* 64:337-346.
 Mauseth, J.D. and W. Halperin. 1975. Hormonal control of organogenesis in *Opuntia polyacantha* (Cactaceae). *Amer. J. Bot.* 62:869-877.
 Murashige, T. and F. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plantarum* 15:473-497.
 Nanda, K.K., T.A. Anaradha, and K. Lal. 1967. Floral induction by gibberellic acid in *Impatiens balsamina* L., a qualitative short-day plant. *Planta* 76:367-370.
 Penner, J. 1960. Über den Einfluß von Gibberellin auf die photoperiodisch bedingten Blühvorgänge bei *Bryophyllum*. *Planta* 55:542-572.
 Rüniger, W. 1984. Einfluß von Wachstumsregulatoren im frühen und späten Stadium der Knospenentwicklung bei *Schlumbergera*. *Gartenbauwissenschaft* 49:100-103.
 Rüniger, W. and R.T. Poole. 1985. *Schlumbergera*, p. 277-282. In: A.H. Halevy (ed.). *Handbook of flowering*. vol. 4. CRC Press, Boca Raton, Fla.
 Sachs, R.M. 1961. Gibberellin, auxin, and growth retardant effects upon cell division and shoot histogenesis. *Adv. Chem. Ser.* 28:49-58.
 Salisbury, F.B. and C.W. Ross. 1978. *Plant physiology*. 2nd ed. Wadsworth Publishing, Belmont, Calif. p. 377-430.
 Sanderson, K.C., Y.S. Ho, W.C. Martin, and R.B. Reed. 1986. Effect of photoperiod and growth regulators on growth of three Cactaceae. *HortScience* 21:1381-1382.
 SAS Institute. 1985. SAS/STAT guide for personal computers. version 6.

- SAS Institute, Cary, N.C.
- Stuart, N.W. and H.M. Cathey. 1962. Control of growth and flowering of *Chrysanthemum morifolium* and *Hydrangea microphylla* by gibberellin, p. 391–399. In: J. Garnaud (ed.). Advances in horticultural science and their applications. vol. 2. Pergamon, New York.
- van Bragt, J. and R.L.M. Pierik. 1971. The effect of autoclaving on the gibberellin activity of aqueous solutions containing gibberellin A₃, p. 133–137. In: J. van Bragt, D.A.A. Mossel, R.L.M. Pierik, and H. Veldstra (eds.). Effects of sterilization on components in nutrient media. H. Veenman & Zonen N. V., Wageningen, The Netherlands.
- Vince-Prue, D. 1975. Photoperiodism in plants. McGraw-Hill, London. p. 217–224; 302–303.
- Wellensiek, S.J. 1960. Stem elongation and flower initiation. Proc. Akad. v. Wetenschappen C 63: 159–166.
- White, J. W., C. Allen, M. Easley, and P. Strock. 1978. Effects of growth regulants on *Opuntia basilaris*. HortScience 13:181–182.
- Wittwer, S.H. and M.J. Bukovac. 1958. The effects of gibberellin on economic crops. Econ. Bot. 12:213–255.
- Yonemura, K. 1979. Studies on the control of flowering in Christmas cactus. Spec. Bul. Aichi-Ken Agr. Res. Ctr., Nagakute, Aichi, Japan. p. 32–33.