Cloning and Characterization of a Sucrose Phosphate Synthase-encoding Gene from Muskmelon

Xiyan Yu, Xiufeng Wang, Jide Fan, and Hongmei Tian

College of Horticulture Science and Engineering, Shandong Agricultural University,
Tai'an 271018, China

Chengchao Zheng¹

State Key Laboratory of Crop Biology, College of Life Sciences, Shandong Agricultural University,
Tai'an, Shandong 271018, China

ADDITIONAL INDEX WORDS. *Cucumis melo*, *CmSPSI*, gene expression, sucrose metabolism

ABSTRACT. Sucrose phosphate synthase [SPS (EC 2.4.1.14)] is thought to play a critical role in sucrose accumulation in muskmelon (*Cucumis melo* L.) fruit. A full-length cDNA clone encoding sucrose phosphate synthase was isolated from muskmelon by reverse transcriptase–polymerase chain reaction and rapid amplification of cDNA ends. The clone, designated *CmSPSI*, contains 3377 nucleotides with an open reading frame of 3162 nucleotides. The deduced 1054 amino acids sequence showed high identities with other plant sucrose phosphate synthases. Northern blot analysis indicated that *CmSPSI* was expressed in leaves, stems, and mature fruit, but was not detected in roots or flowers. Moreover, the mRNA accumulation of *CmSPSI* started at 25 days after pollination (DAP) and reached highest level in mature fruit. Interestingly, both sucrose content and SPS activity increased dramatically between 20 and 30 DAP during fruit development, suggesting that sucrose accumulation may be linked to the *CmSPSI* transcript level in muskmelon fruit.

Sugars are the most important biochemical components for fruit quality. The kind and amount of sugars directly influence fruit flavor components such as sweetness. As the first step toward the genetic improvement of the quality of muskmelon fruit, it is necessary to determine the sugar components accumulated in fruit, elucidate enzymes involved in sugar metabolism, and clarify the relationship between the content of accumulated sugar and the activity of some related enzymes (McCollum et al., 1988; Moriguchi et al., 1992; Sakalo and Kurchii, 2004).

At the middle stage of fruit development, muskmelon fruit undergo a metabolic transition marked by both physical and compositional changes such as netting of the exocarp, mesocarp softening, and the onset of sucrose accumulation (Lester and Dunlap, 1985). Attempts to elucidate the changes in metabolism that lead to accumulation of sucrose have focused on sucrose-metabolizing enzymes during fruit growth and development (Geromel et al., 2006; Lingle and Dunlap, 1987; Schaffer et al., 1987; Winter and Huber, 2000). It was reported that both acid invertase (EC 3.2.1.26) and sucrose phosphate synthase (EC 2.4.1.14) are determinants of sucrose accumulation in melon fruit. However, the decline of the acid invertase appears to be a normal function of fruit maturation and is not the primary factor that determines sucrose accumulation. Rather, the capacity of sucrose synthesis, reflected in the sucrose phosphate synthase (SPS) activity, appears to determine

sucrose accumulation, which is a key component of fruit quality (Hubbard et al., 1989).

Recently, the SPS gene has been isolated from some higher plant species (Hesse et al., 1995; Hubbard et al., 1991; Huber and Huber, 1996; Li et al., 2003). In this report, we present the molecular cloning and characterization of *CmSPSI* from muskmelon. In addition, the differential expression of *CmSPSI* was determined in various tissues and fruit development stages of *Cucumis melo*.

Materials and Methods

PLANT MATERIAL AND TISSUE SAMPLING. Muskmelon inbred line M01-3 was grown in a greenhouse in an experimental farm of Shandong Agricultural University in Tai'an, China, from March through June 2006 with spacing 50 cm between plants and 120 cm between rows. Average day/night temperatures were $\approx 30/20$ °C. Average daylight was ≈ 12 h. Fertilizer was applied at two stages, a preplant broadcast application of 900 kg·ha⁻¹ of 14N–6.1P–29.9K followed by a sidedress application of 150 kg·ha⁻¹ N at flowering stage. Irrigation by furrows was applied as needed. Freshly opened female flowers were tagged on the day of hand-pollination to identify fruit of known age and one fruit per plant was allowed to develop. Different developing stage [5, 10, 15, 20, and 25 d after pollination (DAP)] and mature fruit were harvested. Thirty DAP is considered commercial maturity. Five fruit from each stage were pooled to isolate total RNA. A portion of each fruit was diced into small pieces and used for SPS activity assay and sucrose determination. The SPS activity assays and sucrose determination were carried out on duplicated samples. The experiments were repeated three times. For each replicate, five fruit were used. Vegetative tissues, including leaves, stems, and roots, were sampled along with

Received for publication 2 Dec. 2006. Accepted for publication 5 Apr. 2007.
This work was supported by the National Natural Sciences Foundation (No. 30471191), Provincial Breeding Project of Shandong and Post Doctor Program Foundation of Shandong Agricultural University in China.

¹Corresponding author. E-mail: cczheng@sdau.edu.cn.

flowers at bloom and harvested simultaneously from five 7- to 8-week-old plants. All tissues were quickly frozen in liquid nitrogen and stored at -80°C until late use.

EXTRACTION AND PURIFICATION OF RNA. Total RNA was isolated using the guanidine isothiocyanate–phenol–chloroform method as described by Sambrook et al. (1989) with modification. Frozen tissue samples (5 to 10 g) were ground to a powder in liquid N_2 using a mortar and pestle. Samples of the homogenized, powdered tissue were transferred to 50-mL screw-cap centrifuge tubes prechilled on ice followed by immediate addition of 15 mL solution D (4 M guanidine isothiocyanate, 0.025 M trisodium citrate dehydrate, 0.5% lauryl sarcosine) and of 1.5 mL of 2 M sodium acetate, pH 4.0, 15 mL of phenol equilibrated with distilled water, and 3 mL of 49 chloroform:1 isopropyl alcohol (v/v). After vortexing and incubation for 15 min on ice, the samples were processed according to Sambrook's method. The crude RNA preparations were then treated with DNase (TaKaRa Bio, Kyoto, Japan) to degrade genomic DNA, extracted with 1 phenol:1 chloroform (v/v), and total RNA was precipitated by addition of 1 volume isopropyl alcohol plus 0.1 volume of 3 M sodium acetate, pH 5.5. The precipitated RNA was pelleted by centrifugation, washed with cold 70% ethanol, and resuspended in diethyl pyrocarbonate-treated water. Quality of the extracted RNA was checked by agarose gel electrophoresis and RNA was quantified spectrophotomerically (ultraviolet 2450; Shimadzu Corp., Kyoto, Japan).

CLONING OF *CmSPSI* COMPLETE cDNA. Total RNA isolated from 25 DAP fruit of *C. melo* M01-3 plants was used for the initial cloning of SPS cDNA fragments, including products of 3' and 5' rapid amplification of cDNA ends (RACE). As the initial step in cloning a *C. melo* SPS gene, degenerate primers (P_1 : 5'-TCWAATCCAAGGAAGCCTATGA-3', P_2 : 5'-AACRGCCTGACAACGAAGT-3', W = T + A, R = A + G) were designed based on the conserved domain of SPS genes from other plants in GenBank. Reverse transcriptase–polymerase chain reaction (RT-PCR) was performed according to the protocol of the RNA PCR Kit (AMV) (version 3.0; TaKaRa Bio). The first cDNA was synthesized by reverse transcription using reverse transcriptase with 1 μg RNA as template and with Oligo (dT)-Adaptor primer as first strand primer. The RT reaction program is 42°C for 30 min, 99°C for 5 min, and 5°C for 5 min, 1 cycle. PCR was performed by running the following program: 94°C predenature for 5 min; 94°C for 1 min, 53°C for 1 min, and 72°C for 3 min, 30 cycles; and, 72°C for 10 min. The PCR product was separated by electrophoresis in 1.0% agarose gel. The isolation fragment was cloned by using PMD18-T vector (TaKaRa Bio) for sequencing.

A total of 1 μg RNA from muskmelon fruit was taken to convert mRNAs into cDNAs using a 3' RACE kit (TaKaRa Bio) provided with AMV RT and a universal oligo (dT) containing adapter primer (5'-CTGATCTAGAGGTACCG GATCCT₁₇-3'). The partial *CmSPSI* gene from the 3' end was then amplified by a pair of PCR primers: the gene specific forward primer 1 (5'-AAGACTTTGGTGCGGTGGGCT GATC-3') and reverse primer (5'-CTGATCTAGAGG TACCGGATCC-3') provided with the kit. According to the manufacturer's instructions, PCR was performed under the following conditions: cDNA was denatured at 94°C for 2 min followed by 35 cycles of amplification (94°C for 30 s, 55°C for 30 s, and 72°C for 1 min) and 7 min at 72°C . The PCR product was purified and cloned into pMD18-T vector for sequencing.

Based on the sequence of the initial *C. melo* SPS gene, the specific primers 2, 3, and 4 (primer 2 5'-CCAGCGTGGTTGC CTTGGTTTGC-3', primer 3 5'-CTGCCTCA ATCAAAGTC AGTCCA-3', primer 4 5'-TACCCATTATCAAGGTTAGAT TAGC-3') were designed to amplify the 5' end of *CmSPSI* gene using the 5' RACE system for rapid amplification of cDNA ends (version 2.0; Invitrogen Corp., Carlsbad, CA). The 5' RACE was performed according to the manufacturer's instructions. The PCR product was purified and cloned into pMD18-T vector for sequencing. DNAMAN (version 4.0; Lynnon Biosoft, Quebec), ANTHEPROT (version 5.0; Deleage et al., 2001), and NCBI web were used to analyze the DNA and protein sequences.

NORTHERN BLOT ANALYSIS. For RNA gel-blot analysis, 20 μg of total RNA was separated on 1.2% agarose gels containing formaldehyde. RNA gel blot analysis was performed according to the DIG Northern Starter Kit Manual (Roche, Penzberg, Germany). 3' UTR fragment of *CmSPSI* cDNA [≈ 300 base pairs (bp)] was used as a probe labeled by digoxigenin. RNA transfer and fixation were performed by capillary transfer with $20 \times \text{SSC}$ overnight and baking membrane at 80°C for 2 h. Prehybridization was done at 68°C in DIG Easy Hyb Buffer for 30 min with gentle agitation. Hybridization was carried out in the same buffer overnight at 68°C . The blot was washed twice with $2 \times \text{SSC}$ and 0.1% sodium dodecyl sulfate (SDS) at 15 to 25°C under constant agitation for 5 min. Then the blot was washed twice in $0.1 \times \text{SSC}$ and 0.1% SDS at 68°C under constant agitation for 15 min. The hybridized probes are immunodetected with antidigoxigenin-AP, Fab fragments, and then visualized with the chemiluminescence substrate CDP-*Star*. Enzymatic dephosphorylation of CDP-*Star* by alkaline phosphatase leads to a light emission at a maximum wavelength of 465 nm, which is recorded on x-ray films. Exposure time was 30 min.

SUCROSE CONTENT MEASUREMENT. Sucrose was extracted by grinding flesh tissues (10 g fresh weight) in 80% ethanol, adjusted to pH 7.0 with 0.1 N NaOH, and heated for 5 min at 80°C . The extraction was repeated three times and five fruit were used for each replicate. The extract corresponding to 0.5 g fresh weight was dried in vacuo and redissolved in water. An aliquot (10 μL) of the eluate was analyzed by high-performance liquid chromatography (Shimadzu Corp.), which was equipped with a RID-10A detector and a SCL-10AVP data acquisition system.

SUCROSE PHOSPHATE SYNTHASE EXTRACTION AND ASSAY. Sucrose phosphate synthase extraction and assay were performed in a manner similar to that previously described by Lowell et al. (1989) and Hubbard et al. (1989) with some minor modifications. SPS was extracted from ≈ 1 g of minced frozen melon tissue by homogenizing for 1 min on ice in 200 mM Hepes/NaOH (pH 7.5), 5 mM MgCl_2 , 1 mM EDTA, 0.5 $\text{mg}\cdot\text{mL}^{-1}$ of BSA, 0.005% Triton X-100 (Sigma-Aldrich Corp., St. Louis), 1 M NaCl, 1% insoluble PVP, and 5 mM DTT. Homogenates were quickly desalted at 4°C . Desalted buffer contained 20 mM Hepes/NaOH (pH 7.5), 0.25 mM MgCl_2 , 0.01% 2-mercaptoethanol, 1 mM EDTA, 0.05% BSA, 0.2% glycerine. Desalted extract was assayed immediately.

Sucrose phosphate synthase assay contained 50 mM Hepes/NaOH (pH 7.5), 15 mM MgCl_2 , 25 mM Fru-6-P, 25 mM Glu-6-P, and 25 mM UDP-Glu. Control was boiled for 10 min immediately after the addition of the desalted enzyme extract. The reactions were incubated for 30 min at 37°C and then stopped by transfer to a boiling water bath. To the 70- μL sample, 70 μL

C. melo	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRM	100
L. esculentum	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRM	100
S. tuberosum	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRM	100
A. chinensis	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRM	99
I. batatas	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRL	100
B. vulgaris	MAGNDWINSYLEAILDVGPGLDDAKSSLLLRERGRFSPTRYFVEVITGFDETDLERSWIRACATSPQERNTRLENMCWRIUNLARCKKQLEGCAQRK	100
C. melo	AKRRCEERERGRREAVADMSEDLSGEKGDIVTDMSSHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	200
L. esculentum	AKRRCEERERGRREAVADMSEDLSGEKGDIVTDMSSHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	200
S. tuberosum	AKRRCEERERGRREAVADMSEDLSGEKGDIVTDMSSHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	200
A. chinensis	AKRRCEERERGRREAVADMSEDLSGEKGDIVTDMSSHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	199
I. batatas	AKRRCEERERGRREAVADMSEDLSGEKGDIVTDMSSHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	200
B. vulgaris	AKRRCEERERGRREAVADMSEDLSGEKDI.....SAHGSETRGLPRISSEVTHEAVUSQCRCKLYIVLISLHGLIRGENMELGRSDTGGQVKYVVEL	195
C. melo	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...SEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	297
L. esculentum	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...SEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	297
S. tuberosum	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...TEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	296
A. chinensis	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...DEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	296
I. batatas	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...TEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	297
B. vulgaris	ARALGSMGVYRVLDLTVQVSSPVDVSYGEPTMLTPISTDGLN...DEMGESSGAYIRIPFGPREKYIPKEQLWPYIPEFVDGALNHIIQMSKVLGE	295
C. melo	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	397
L. esculentum	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	397
S. tuberosum	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	396
A. chinensis	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	396
I. batatas	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	397
B. vulgaris	QIGGCHPVWVPAIHGHYADAGDS AALLSGALNVPMLE TGHS LG RD KLEQLLRGRSKDEINSTYKIMRRIEAEELTLDASEIVITSTROEIDEQWRLYD	395
C. melo	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	496
L. esculentum	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	496
S. tuberosum	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	495
A. chinensis	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	495
I. batatas	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	496
B. vulgaris	GFDPFLERKLRLRIKRWVSCYGRFMFRMVIIPPGEHFEHIVPHEGDMGDTGESESGDKI..PDPPHIAEIMRFFSNPRKPMILALARPDPKKNLTTLVRAF	495
C. melo	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	596
L. esculentum	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	596
S. tuberosum	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	595
A. chinensis	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	596
I. batatas	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	596
B. vulgaris	GECPRLRLANLTLIMGNRDNIDEVSSSTNSALLSLILKMDIKYDLYGOVAYPKHHKQSDVDPDIYRLAAKTGVFINPAFIEPFGTLIEAAAYGLPMVAT	595
C. melo	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..PGDDDDENSETSPSPDSLRD	695
L. esculentum	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..PGDDDDENSETSPSPDSLRD	695
S. tuberosum	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..SIDDDDDENSETSPSPDSLRD	694
A. chinensis	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..S.DDGNENSETSPSPDSLRD	694
I. batatas	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..NA.DDGNENSETSPSPDSLRD	694
B. vulgaris	KNGGVPDIEHVLNGLLVDPHDQQAIDALLKLVDKQPMKCRANGLKNIHLFSUPEHCKTYLSRIASCKPQRPQWLR..DEGLDNEPSPSPDSLRD	694
C. melo	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	795
L. esculentum	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	795
S. tuberosum	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	794
A. chinensis	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	794
I. batatas	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	789
B. vulgaris	IEDISLNRFLSLDGEKNENKADSTLDPERKSKLENANVLSLSEKAPKSTKSWSSDRADQNPAGAKTFAIRRRRHIFVIAVDCDASSGLSGSVKKIFE	788
C. melo	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	895
L. esculentum	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	895
S. tuberosum	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	894
A. chinensis	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	894
I. batatas	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	889
B. vulgaris	AVERERSEGSIGFILASSFNISEVQSFLVSEGMSPDFAICNSGDLVYSSFHSEONPFVVDLYHSHIEYRUGGEGRLRTLVRWAASITDKNGENGE	888
C. melo	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	995
L. esculentum	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	995
S. tuberosum	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	994
A. chinensis	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	994
I. batatas	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	989
B. vulgaris	HIVVEDEDSADYCYTFKVKCPGKVPFAKELRRVIRIQALRCHAVYQNGSRINIPVLAASRSQALRYLYLRUGMLLSLKVVFVGESGDTDYEGLIGGLR	988
C. melo	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1034
L. esculentum	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1034
S. tuberosum	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1033
A. chinensis	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1033
I. batatas	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1048
B. vulgaris	RAVIMKGLCTNASSLIHGNRYPLSDVLPFDPSPNVIQADEECSSSTEIRSLLEKLAVALKG	1045

Fig. 1. Alignment of predicted amino acid sequences of sucrose phosphate synthase (SPS) genes from different plants. The GenBank accession numbers for the plant SPS sequences are: *Cucumis melo* ABF47344, *Lycopersicon esculentum* AAU29197, *Solanum tuberosum* Q43845, *Actinidia chinensis* AAL86360, *Ipomoea batatas* AAL34531, and *Beta vulgaris* CAA57500. This alignment was produced using DNAMAN (version 4.0; Lynnon Biosoft, Quebec). Conserved residues are shaded in black. Dark gray shading indicates similar residues in five of six of the sequences and clear gray shading indicates similar residues in three of six of the sequences.

of 30% NaOH was added, boiled for 10 min, and cooled. One milliliter anthrone reagent (76 mL of sulfuric acid, 30 mL H₂O, and 150 mg of anthrone) was then added, and the tubes were incubated for 20 min at 40 °C. Absorbance was measured with a microplate reader (model 3550-ultraviolet; Bio-Rad Laboratories, Hercules, CA) at 650 nm and compared with sucrose standards.

Results

CLONING AND CHARACTERIZATION OF *CmSPS1*. To isolate SPS cDNA from muskmelon, we first designed a pair of degenerated primers according to the conserved domain of SPS amino acid sequences from other plant species. A 1408-bp cDNA fragment was amplified by RT-PCR. A 1600-bp and a 750-bp cDNA fragments were produced by 5' RACE and 3' RACE, respectively. After sequencing confirmation, a 3377-bp cDNA clone including a full-length coding region was isolated from muskmelon and named *CmSPS1* (GenBank accession DQ521271). The deduced amino acid sequence of *CmSPS1* contains 1054 amino acid residues with a predicted signal sequence cleavage site between residue 179 and residue 180. The predicted mature protein might locate in the cytoplasm as a result of the mature protein without transmembrane regions. The alignment analysis showed that *CmSPS1* shared with overall amino acid identities of 98.96%, 96.77%, 84.72%, 75.65%, and 73.72% to tomato (*Lycopersicon esculentum* L.), potato (*Solanum tuberosum* L.), kiwifruit (*Actinidia chinensis* Planch.), sweetpotato (*Ipomoea batatas* L.), and sugar beet (*Beta vulgaris* L.) SPS, respectively (Fig. 1).

***CmSPS1* DIFFERENTIALLY EXPRESSED IN DIFFERENT MUSKMELON TISSUES.** To determine the pattern of *CmSPS1* expression in different muskmelon organs, a northern blot hybridization analysis was performed. The result showed that *CmSPS1* transcripts are easily detected in the leaves, stems, and mature fruit, but cannot be detected at all in roots and flowers (Fig. 2), suggesting that *CmSPS1* may play a role in leaf, stem, and fruit development in muskmelon.

REGULATION OF *CmSPS1* mRNA ABUNDANCE DURING FRUIT DEVELOPMENT AND RIPENING. To further investigate the devel-

Fig. 2. RNA blot analysis of *CmSPS1* spatial expression in muskmelon plants. Approximately 20 µg of total RNA from root, stem, leaf, flower, and mature fruit of muskmelon inbred line M01-3 were separated on 1.2% agarose gels containing formaldehyde. Root, stem, leaf, and flower tissues were harvested simultaneously from five 7- to 8-week-old plants and mature fruit were harvested as they mature from five 11- to 12-week-old plants. The blot was hybridized with a *CmSPS1*-specific probe. The ethidium bromide-stained rRNA band in the agarose gel is shown as a loading control.

Fig. 3. RNA blot analysis of *CmSPS1* expression in developmental muskmelon fruit. Approximately 20 µg of total RNA from muskmelon inbred line M01-3 fruit of 5, 10, 15, 20, and 25 d after pollination (DAP) and mature fruit, respectively, were separated on 1.2% agarose gels containing formaldehyde. Five fruit of each stage were pooled to isolate RNA. The blot was hybridized with a *CmSPS1*-specific probe. The ethidium bromide-stained rRNA band in the agarose gel is shown as a loading control.

opmental regulation of *CmSPS1* in fruit mesocarp tissues, a RNA blot analysis was carried out using different mesocarp tissues from 5 DAP to ripening. The result showed that *CmSPS1* mRNA was not detected in the mesocarp tissues before 20 DAP. However, the *CmSPS1* strongly expressed in the mesocarp tissues after 25 DAP, indicating that *CmSPS1* may be involved in sucrose accumulation in fruit ripening in muskmelon (Fig. 3).

SUCROSE CONTENT AND SUCROSE PHOSPHATE SYNTHASE ACTIVITY DURING FRUIT DEVELOPMENT AND RIPENING. To demonstrate the functions of *CmSPS1* in regulating fruit quality, sucrose content and SPS activity were analyzed. The results showed that very low concentration of sucrose was detected in young and unripe muskmelons. However, a rapid accumulation of sucrose in fruit was observed between 20 and 30 DAP (Fig. 4A). Similarly, SPS activity showed a significant increase during the later stage of fruit development as well (Fig. 4B), indicating an essential role of SPS in sucrose metabolism in muskmelon fruit.

Discussion

SPS catalyzes net sucrose synthesis in plants and its activity is generally high in source tissues and low in sink organs (Huber and Huber, 1992). It has been proven that SPS activity can be a controlling factor for sucrose synthesis and also for photosynthesis (Shinano et al., 2006; Stitt et al., 1988; Zuniga-Feest et al., 2005). To increase photosynthesis and sucrose synthesis, Worrell et al. (1991) genetically manipulated tomatoes by overexpressing a maize (*Zea mays* L.) SPS cDNA under the control of the promoter of the small subunit of Rubisco from tobacco (*Nicotiana tabacum* L.). Several transgenic lines showed a higher SPS activity and higher sucrose level in their leaves than untransformed controls. Previous studies demonstrated that the SPS activity in transformed tomato leaves was six times greater than that of the controls. At low irradiances, no increase in net photosynthesis was observed, but the light- and CO₂-saturated rates of photosynthesis in transformed leaves were increased by ≈20% (Galtier et al., 1993, 1995; Micallef et al., 1995). In this study, we demonstrated that *CmSPS1* was expressed in stem and leaf but not in root and flower tissues, implying that *CmSPS1* may be also involved in sucrose synthesis and photosynthesis in muskmelon. Langenkämper et al. (2002) reported that there were at least three families of

Fig. 4. Sucrose content and sucrose phosphate synthase (SPS) activity during M01-3 muskmelon fruit development. Five, 10, 15, 20, 25, and 30 d after pollination (DAP) fruit were harvested, respectively. Thirty DAP is considered commercial maturity. The SPS activity assays and sucrose determination were carried out on duplicated samples. The experiments were repeated three times and five fruit were used for each replicate. (A) Sucrose content during M01-3 muskmelon fruit development; (B) SPS activity during M01-3 muskmelon fruit development. Bars indicate SE.

SPS genes, designated A, B, and C based on the SPS genes isolated from *Arabidopsis* and other plant species. Recently, Castleden et al. (2004) revealed the finding of five families of SPS genes in wheat (*Triticum aestivum* L.) and other monocotyledonous plants from the family Poaceae (grasses). Each of the SPS gene families in wheat showed different, but overlapping, spatial and temporal expression patterns, and in most organs at least two different SPS genes are expressed. In this study, we cloned and characterized a SPS gene from muskmelon, *CmSPS1*. However, it is not clear how many SPS genes exist in muskmelon. We are constructing cDNA libraries from muskmelon leaf and fruit and try to isolate other SPS gene members involved in sucrose synthesis or photosynthesis in muskmelon by screening the cDNA libraries.

Sucrose phosphate synthase activity has been suggested to determine the rate of sucrose synthesis and the level of sucrose accumulation in late fruit development of various plant species such as tomato (Dali et al., 1992; Miron and Schaffer, 1991), muskmelon (Hubbard et al., 1989; Lingle and Dunlap, 1987), banana (*Musa acuminata* L.) (Hubbard et al., 1990), asian pear 'Chojuro' [*Pyrus pyrifolia* (Burm.f.) Nakai] and 'Yali' (*P. bretschneideri* Rehd.) (Moriguchi et al., 1992), peach [*Prunus persica* (L.) Batsch], strawberry (*Fragaria × ananassa* Duch.), kiwi (*Actinidia deliciosa* A. Chev.), and mango (*Mangifera indica* L.) (Hubbard et al., 1991). Recently, SPS-encoding genes have been isolated and characterized from some higher plant species (Hesse et al., 1995; Hubbard et al., 1991; Huber and Huber, 1996). To date, however, there have been only a few studies of SPS gene expression in relation to fruit tissues. Nascimanto et al. (1997) reported that the accumulation of sucrose occurred 4 d after SPS mRNA and activity had reached their maxima during banana fruit ripening. In citrus (*Citrus unshiu* Marc.), the levels of *CitSPS1* mRNA in immature fruit were low but very high in mature fruit (Komatsu et al., 1996). The SPS-antisense potato transformants showed a reduced expression of sucrose-phosphate synthase, which led to inhibition of sucrose synthesis in tuber discs (Geigenberger et al., 1999). Our present results showed that the *CmSPS1* transcripts were not detected in the young muskmelon fruit but rapid accumulation in the mature fruit (Fig. 3). This pattern of *CmSPS1* expression in mesocarp tissues is in accordance with the alteration of SPS activity and sucrose accumulation during fruit development of muskmelon (Fig. 4). Taken together, these data led us to conclude that the sucrose content in plant sink organs is regulated by the level of SPS expression. Therefore, increasing SPS expression might be an important regulatory event of sweetening during melon fruit ripening. Currently, we are overexpressing *CmSPS1* under the control of a fruit-specific promoter, which will help reveal the potential application of *CmSPS1* in controlling the quality of muskmelon fruit.

Literature Cited

- Castleden, C.K., N. Aoli, V.J. Gillespie, E.A. MacRae, W.P. Quick, P. Buchner, C.H. Foyer, R.T. Furbank, and J.E. Lunn. 2004. Evolution and function of the sucrose-phosphate synthase gene families in wheat and other grasses. *Plant Physiol.* 135:1753–1764.
- Dali, N., D. Michaud, and S. Yelle. 1992. Evidence for the involvement of sucrose phosphate synthase in the pathway of sugar accumulation in sucrose-accumulating tomato fruits. *Plant Physiol.* 99:434–438.
- Deleage, G., C. Combet, C. Blanchet, and C. Geourjon. 2001. ANTHEPROT: An integrated protein sequence analysis software with client/server capabilities. *Comput. Biol. Med.* 31:259–267.
- Galtier, N., C.H. Foyer, J. Huber, T.A. Voelker, and S.C. Huber. 1993. Effects of elevated sucrose-phosphate synthase activity on photosynthesis, assimilate partitioning, and growth in tomato (*Lycopersicon esculentum* var. UC82B). *Plant Physiol.* 101:535–543.
- Galtier, N., C.H. Foyer, E. Murchie, R. Alred, P. Ouick, T.A. Voelker, C. Thepenier, G. Lasceve, and T. Betsche. 1995. Effects of light and atmospheric carbon dioxide enrichment on photosynthesis and carbon partitioning in the leaves of tomato (*Lycopersicon esculentum* L.) plants overexpression sucrose phosphate synthase. *J. Expt. Bot.* 46:1335–1344.
- Geigenberger, P., R. Reimholz, U. Deiting, U. Sonnewald, and M. Stitt. 1999. Decrease expression of sucrose phosphate synthase strongly inhibits the water stress-induced synthesis of sucrose in growing potato tubers. *Plant J.* 19:119–129.

- Geromel, C., L.P. Ferreira, S.M.C. Guerreiro, A.A. Cavalari, D. Pot, L.F.P. Pereira, T. Leroy, L.G.E. Viera, P. Mazzafera, and P. Marraccini. 2006. Biochemical and genomic analysis of sucrose metabolism during coffee (*Coffea arabica*) fruit development. *J. Expt. Bot.* 57:3243–3258.
- Hesse, H., U. Sonnewald, and L. Willmitzer. 1995. Cloning and expression analysis of sucrose phosphate synthase from sugar beet. *Mol. Gen. Genet.* 247:515–520.
- Hubbard, N.L., S.C. Huber, and D.M. Pharr. 1989. Sucrose phosphate synthase and acid invertase as determinants of sucrose concentration in developing muskmelon (*Cucumis melo* L.) fruits. *Plant Physiol.* 91:1527–1534.
- Hubbard, N.L., D.M. Pharr, and S.C. Huber. 1990. Role of sucrose phosphate synthase in sucrose biosynthesis in ripening bananas and its relationship to the respiratory climacteric. *Plant Physiol.* 94: 201–208.
- Hubbard, N.L., D.M. Pharr, and S.C. Huber. 1991. Sucrose phosphate synthase and other sucrose metabolizing enzymes in fruits of various species. *Physiol. Plant.* 82:191–196.
- Huber, S.C. and J.L. Huber. 1992. Role of sucrose phosphate synthase in sucrose metabolism in leaves. *Plant Physiol.* 99:1275–1278.
- Huber, S.C. and J.L. Huber. 1996. Role and regulation sucrose phosphate synthase in higher plants. *Annu. Rev. Plant Mol. Biol.* 47:431–445.
- Komatsu, A., Y. Takanokura, T. Akihama, and M. Omura. 1996. Cloning and molecular analysis of cDNAs encoding three sucrose phosphate synthase isoforms from a citrus fruit (*Citrus unshiu* Marc.). *Mol. Gen. Genet.* 252:346–351.
- Langenkämper, G., R.W.M. Fung, R.D. Newcomb, R.G. Atkinson, R.C. Gardner, and E.A. MacRae. 2002. Sucrose phosphate synthase genes in plants belong to three different families. *J. Mol. Evol.* 54:322–332.
- Lester, G.E. and J.R. Dunlap. 1985. Physiological changes during development and ripening of ‘Perlita’ muskmelon fruits. *Scientia Hort.* 26:323–331.
- Li, C.R., X.B. Zhang, and C.S. Hew. 2003. Cloning of a sucrose phosphate synthase gene highly expressed in flowers from the tropical epiphytic orchid *oncidium goldiana*. *J. Expt. Bot.* 54: 2189–2191.
- Lingle, S.E. and J.R. Dunlap. 1987. Sucrose metabolism in netted muskmelon fruit during development. *Plant Physiol.* 84:386–389.
- Lowell, C.A., P.T. Tomlinson, and K.E. Koch. 1989. Sucrose metabolizing enzymes in transport tissues and adjacent sink structures in developing citrus fruit. *Plant Physiol.* 90:1394–1402.
- McCollum, T.G., D.J. Huber, and D.J. Cantliffe. 1988. Soluble sugar accumulation and activity of related enzymes during muskmelon fruit development. *J. Amer. Soc. Hort. Sci.* 113:399–403.
- Micallef, B.J., K.A. Haskins, P.J. Vanderveer, K.-S. Roh, C.K. Shewmaker, and T.D. Sharkey. 1995. Altered photosynthesis, flowering, and fruiting in transgenic tomato plants that have an increased capacity for sucrose synthesis. *Planta* 196:327–334.
- Miron, D. and A.A. Schaffer. 1991. Sucrose phosphate synthase, sucrose synthase, and invertase activities in developing fruit of *Lycopersicon esculentum* Mill. and sucrose accumulating *Lycopersicon hirsutum* Humb. and Bonpl. *Plant Physiol.* 95:623–627.
- Moriguchi, T., K. Abe, T. Sanada, and S. Yamaki. 1992. Levels and role of sucrose synthase, sucrose-phosphate synthase, and acid invertase in sucrose accumulation in fruit of asian pear. *J. Amer. Soc. Hort. Sci.* 117:274–278.
- Nascimento, J.R., B.R. Cordenunsi, F.M. Lajolo, and M.J. Alcocer. 1997. Banana sucrose-phosphate synthase gene expression during fruit ripening. *Planta* 203:283–288.
- Sakalo, V.D. and V.M. Kurchii. 2004. Hormonal control of sucrose phosphate synthase and sucrose synthase in sugar beet. *Russ. J. Plant Physiol.* 51:183–188.
- Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schaffer, A.A., B. Aloni, and E. Fogelman. 1987. Sucrose metabolism and accumulation in developing fruit of *Cucumis*. *Phytochemistry* 26:1883–1887.
- Shinano, T., K. Ando, K. Okazaki, and M. Osaki. 2006. Developmental changes of plant affecting primary photosynthate distribution in rice leaves. *Photosynthetica* 44:591–598.
- Stitt, M., I. Wilke, R. Feil, and H.W. Heldt. 1988. Coarse control of sucrose phosphate synthase in leaves: Alteration of the kinetic properties in the response to the rate of photosynthesis and the accumulation of sucrose. *Planta* 174:217–230.
- Winter, H. and S.C. Huber. 2000. Regulation of sucrose metabolism in higher plants: Localization and regulation of activity of key enzymes. *J. Crit. Rev. Biochem. Mol. Biol.* 35:253–289.
- Worrell, A.C., J.M. Bruneau, K. Summerfelt, M. Boersig, and T.A. Voelker. 1991. Expression of maize sucrose phosphate synthase in tomato alters leaf carbohydrate partitioning. *Plant Cell* 3:1121–1130.
- Zuniga-Feest, A., D.R. Ort, A. Gutierrez, M. Gidekel, L.A. Bravo, and L. Corcuera. 2005. Light regulation of sucrose-phosphate synthase activity in the freezing-tolerant grass *Deschampsia antarctica*. *Photosyn. Res.* 83:75–86.