Genetic Diversity and Population Structure Analysis of Chinese Wild Grape Using Simple Sequence Repeat Markers

Beibei Li

College of Enology, Northwest Agriculture & Forestry University, Yangling, 712100, China; and Zhengzhou Fruit Research Institute of the Chinese Academy of Agricultural Sciences, Zhengzhou, 450009, China

Xiucui Fan, Ying Zhang, Chonghuai Liu, and Jianfu Jiang

Zhengzhou Fruit Research Institute of the Chinese Academy of Agricultural Sciences, Zhengzhou, 450009, China

ADDITIONAL INDEX WORDS. genetic structure and differentiation, genetic variability, molecular markers, SSR, *Vitis*

ABSTRACT. Chinese wild *Vitis* is a useful gene source for resistance to biotic and abiotic stresses, although there is little research on its genetic diversity and structure. In this study, nine simple sequence repeat (SSR) markers were used to assess the genetic diversity and genetic structure among 100 *Vitis* materials. These materials included 77 indigenous accessions representing 23 of 38 wild *Vitis* species/cultivars in China, 18 *V. vinifera* cultivars, and the five North American species *V. aestivalis*, *V. girdiana*, *V. monticola*, *V. acerifolia*, and *V. riparia*. The SSR loci used in this study for establishing an international database (*Vitis* International Variety Catalogue) revealed a total of 186 alleles in 100 *Vitis* accessions. The mean values for the gene diversity (GD) and polymorphism information content (PIC) per locus were 0.91 and 0.90, respectively, which indicates that the discriminatory power of the markers is high. Based on the genetic distance data, the 100 *Vitis* accessions were divided into five primary clusters by cluster analysis, and five populations by structure analysis; these results indicate these Chinese wild grapes were more genetically close to European grapes than to North American species. In addition, the clustering patterns of most accessions correlated with the geographic distribution. An analysis of molecular variance (AMOVA) revealed that 3.28%, 3.27%, and 93.46% of the variance occurred between populations, between individuals within populations, and between individuals within the entire population, respectively. In addition, we identified three previously undescribed accessions (Wuzhi-1, MZL-5, and MZL-6) by cluster analysis. Our results reveal a high level of genetic diversity and variability in *Vitis* from China, which will be helpful in the use of genetic resources in future breeding programs. In addition, our study demonstrates that SSR markers are highly suitable for further genetic diversity analyses of Chinese wild grapes.

Grapes (*Vitis* sp.) are one of the most economically valuable horticultural crops in the world. They are widely used to produce wine, table grapes, raisins, juice, and health care products (Ren and Wen, 2007; Wan et al., 2013; Zhou et al., 2017). The genus, which includes more than 70 *Vitis* species, consists of two subgenera—*Vitis* and *Muscadinia*—in which more than 38 *Vitis* species originated from China (Liu et al., 2012; Wan et al., 2008). Currently, most cultivated grapes in China originated from Europe, and grape commercial breeding populations in China share a narrow genetic base as a result of their common origins from a number of popular cultivars, such as Muscat Hamburg and Kyoho. However, China, as one of the major gene centers of *Vitis* species, has abundant wild grape species that occur naturally in all the provinces except Xinjiang, and the number of wild grape species per province ranges from 1 to 29 (Jiang et al., 2015). Chinese wild grapes have strong abiotic stress tolerance and could be widely used for modern grape breeding (Liu et al., 2014; Wan et al., 2013). There is still

a great interest among grape breeders in broadening the genetic base of cultivated grapes and also in tapping into the gene pool of the wild relatives to enhance stress resistance.

Traditionally, the methods used for assessing the genetic diversity and relationships among wild grape species have relied mostly on morphology, palynology, and isoenzymes (Chao and Niu, 1981; He and Chao, 1982; Wan et al., 2008). However, these methods are easily affected by environmental conditions and developmental stages (Luo et al., 2001). Fortunately, DNA molecular marker techniques are able to overcome these limitations and act as powerful tools for evaluating genetic diversity and relationships (Guan et al., 2019; Queiroz et al., 2015; Riaz et al., 2018). Therefore, a variety of molecular markers, such as random amplified polymorphic DNA (Luo et al., 2001)], intersimple sequence repeat (Jing and Wang, 2013; Zhang et al., 2011)], sequence-related amplified polymorphism (SRAP) (Liu et al., 2013), and SSR (Jing et al., 2013; Liu et al., 2012)], have been used recently to examine the genetic diversity and relationships among wild Chinese grape species. Klein et al. (2018) also used high-throughput sequencing to clarify evolutionary relationships among North American *Vitis* species. Because SSR markers show high levels of polymorphisms, reproducibility, and codominant inheritance (Aradhya et al., 2013; Li et al., 2017), they have been widely used to study

Received for publication 5 Nov. 2020. Accepted for publication 22 Jan. 2021.
Published online 8 March 2021.

C.L. and J.J. are corresponding authors. E-mail: liuchonghuai@caas.cn or jiangjianfu@caas.cn.

This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

the genetic diversity of grapes (Cao et al., 2020; Guo et al., 2010), sugarcane [*Saccharum officinarum* (Ali et al., 2019)], wheat [*Triticum aestivum* (Abbasov et al., 2018)], cabbage [*Brassica oleracea* (El-Esawi et al., 2016)], and hazelnuts [*Corylus heterophylla* (Zhao et al., 2020)].

Early research focused on the classification of Chinese wild grapes, but the taxonomic status of many of the indigenous Chinese species and cultivars remains obscure, such as *V. amurensis* var. *yanshanensis* (Zhang et al., 2018). Later, as a result of the importance of wild grapes in breeding, a number of reports focused on the analysis of genetic diversity among wild grapes native to China and cultivated grapes originating from North America and Europe (Jing and Wang, 2013; Jing et al., 2013; Péros et al., 2011). However, there is still a need to investigate further the genetic diversity of wild grape resources to enable breeders to broaden the genetic base of cultivated grapevines through the incorporation of wild relatives (Fan et al., 2015; Jing et al., 2013; Liu et al., 2012; Zhang et al., 2018).

Recently, Liu et al. (2012) used SSR and SRAP to analyze the genetic diversity of 15 Chinese wild grape species (Liu et al., 2012). To understand more fully the genetic background of Chinese wild grapes, we extend that work in this study to include 23 species/cultivars, in which most wild species are included. This study aimed to characterize the genetic diversity and population structure of 100 accessions belonging to Chinese wild *Vitis* species/cultivars, *V. vinifera*, *V. aestivalis*, *V. girdiana*, *V. monticola*, *V. acerifolia*, and *V. riparia*. The results may provide invaluable information for the better use of Chinese wild grape germplasms in breeding. Moreover, to connect our findings with the *Vitis* International Variety Catalogue database (Maul and Töpfer, 2015), we assessed the genetic diversity and relationships of 100 *Vitis* materials standardized by using the nine SSR markers used in the database to obtain useful information on these materials. The genotypes obtained in this study can be connected to the *Vitis* International Variety Catalogue database after standardization, which will facilitate the sharing of information among different laboratories.

Materials and Methods

PLANT MATERIALS. Plant materials were collected from the China Grape Germplasm Repository in Zhengzhou Henan Province. A total of 100 *Vitis* accessions were used in this study. They consisted of 77 accessions representing 23 Chinese wild *Vitis* species/cultivars, 18 *V. vinifera* cultivars, and the North American *V. aestivalis*, *V. girdiana*, *V. monticola*, *V. acerifolia*, and *V. riparia*. Moreover, four cultivars of the 18 *V. vinifera* cultivars were used for SSR standardization for comparison within international databases (Table 1). The young leaves of each accession were harvested and stored at -80°C for later DNA extraction.

GENOMIC DNA EXTRACTION AND GENOTYPING. Genomic DNA was extracted from the young leaves using a Plant Genomic DNA Kit (Aidlab, Beijing, China) according to the manufacturer's instructions. The DNA quality was checked by 1% (w/v) agarose gel electrophoresis. The genomic DNA was diluted to 30 to 50 ng· μL^{-1} and was stored at -20°C .

Nine SSR loci distributed over six chromosomes in the grape were used, and they have shown a high discrimination capacity: VVMD5 and VVMD7 (Bowers et al., 1996); VVS2 (Thomas and Scott 1993); VVMD 32, VVMD25, VVMD27, and

VVMD28 (Bowers et al., 1999); and VrZAG62 and VrZAG79 (Sefc et al., 1999). All the forward primers were labeled with FAM (Carboxyfluorescein) or HEX (Hexachloro-Fluorescein) fluorescent dye at the 5'-end.

The polymerase chain reaction (PCR) was performed in a 20- μL volume containing 30 to 50 ng of template DNA, 1 $\times$ PCR buffer (Mg^{2+} plus) (TaKaRa, Beijing, China), 0.25 mM deoxynucleoside triphosphates, 0.5 μM of each primer set, and 1 U of Taq DNA polymerase (TaKaRa). The PCR amplification was conducted under the following thermal conditions: 4 min at 94°C for initial denaturation, followed by 35 cycles of denaturation at 94°C for 30 s, annealing at the optimum annealing temperature for 30 s, then at 72°C for 45 s, with a final extension at 72°C for 10 min. The final reaction was held at 4°C .

SSR DATA ACQUISITION. The PCR-amplified products were separated using capillary electrophoresis on an ABA-3730XL DNA Analyzer (Thermo Fisher Scientific, Waltham, MA). An internal size standard (GeneScan-500ROX, Thermo Fisher Scientific) was used to estimate the approximate molecular weights of the amplified products. Ultimately, the peaks were shown by size and height using generic description software (GeneMapper version 3.2, Thermo Fisher Scientific).

SSR DATA ANALYSIS. To assess the genetic diversity among the accessions, the following parameters were applied using the statistical software POPGEN version 1.32 (Krawczak et al., 2006): the number of alleles (N_a), number of effective alleles (N_e), observed heterozygosity (H_o), expected heterozygosity (H_e), and Shannon's information index (I) for each locus. The PIC and GD of each primer pair was estimated using PowerMarker version 3.25 (Kejun and Muse, 2005). The genetic diversity index among different populations was calculated using GenALEX version 6.41 (standard model) (Peakall and Smouse, 2006).

Genetic distances were calculated in accordance with Nei (1973) and were used to construct a dendrogram based on the unweighted pair-group method of mathematical averages (UPGMA) using MEGA5 software (Tamura et al., 2011), and the dendrogram was visualized using EvolView version 2 (He et al., 2016). A principal coordinate analysis (PCA) and an AMOVA were also conducted with GenALEX version 6.41. The population structure was investigated with STRUCTURE software (Li et al., 2019), which is based on an admixture model with correlated allelic frequencies. The number of populations (K) was set from 2 to 10, and 10 independent runs for each K were conducted using a burn-in period of 10,000 steps, followed by 100,000 iterations. The best K was estimated based on the delta K method using STRUCTURE HARVESTER (Earl and Vonholdt, 2012).

Results

ALLELIC VARIANCE AND GENETIC DIVERSITY. During this study, nine SSR primers were used to detect the molecular variance among the 100 grape samples; 100 unique genotypes were detected (Supplemental Table 1). Each of the nine loci analyzed was polymorphic, and resulted in 186 alleles, with an average of 20.667 alleles per locus (Table 2). N_a varied from 16 (VVMD7) to 29 (VVMD28). N_e ranged from 8.306 to 17.549, with a mean value of 11.472. GD was between 0.883 and 0.943, with an average of 0.910. The average value for I was 1.245, with a range from 0.966 to 1.582. PIC values varied from 0.873 for VVMD5 to 0.940 for VVMD28, with an average of 0.903

Table 1. Geographic characteristics and the group information of 100 *Vitis* materials examined in this study.

Species	Code no.	Clone	Origin	Cluster no.
<i>V. riparia</i>	83	DVIT1884 b43-15M44	United States	I
<i>V. aestivalis</i>	84	DVIT2382Xiaputao	United States	I
<i>V. monticola</i>	85	DVIT1847Tianshanputao	United States	I
<i>V. acerifolia</i>	87	DVIT1296Qiyeputao	United States	I
<i>V. girdiana</i>	86	DVIT1387Shanguputao	United States	I
<i>V. davidii</i>	88	ZLL	Hunan Province, R6 ^y	II
	90	HHBG	Huaihua, Hunan Province, R6 ^y	II
	89	BPT-1	Hunan Province, R6 ^y	II
	92	HT-1	Huaihua, Hunan Province, R6 ^y	II
	91	WH-ci	Wuhan Province, R6 ^y	II
	98	Gaoshan-1	Jiangxi Province, R2 ^y	II
	99	Ziqiu	Hunan Province, R2 ^y	II
	100	Gaoshan-2	Jiangxi Province, R2 ^y	II
	94	Black pearl	Hunan Province, R6 ^y	II
	93	ZF-1	Huaihua, Hunan Province, R6 ^y	II
	95	Xiangzhenzhu (Red)	Hunan Province, R6 ^y	II
	96	Xiangzhenzhu (Green)	Hunan Province, R6 ^y	II
	97	ES1424	Hubei Province, R6 ^y	II
<i>V. amurensis</i>	78	SPT-1	Northeastern China, R4 ^y	III
	79	SPT-2	Northeastern China, R4 ^y	III
	80	SPT0933	Northeastern China, R4 ^y	III
	81	DLS	Beijing Province, R3 ^y	III
<i>V. amurensis</i> var. <i>yanshanensis</i>	82	Beizhi1702	Beijing Province, R3 ^y	III
<i>V. vinifera</i>	72	Sauvignon Blanc ^z	France	III
	74	Carignan ^z	Spain	III
	75	Gamay ^z	Spain	III
	76	Syrah ^z	Iran	III
	67	Cardinal	United States	III
	62	Muscat Hamburg	England	III
	59	6-28	China	III
	60	Jinanzaozhong	China	III
	65	Jingfeng	China	III
	64	Centennial Seedless	United States	III
	73	Jingkejing	China	III
	70	Jingyu	China	III
	66	Queen of Vineyard	Hungary	III
	63	Shenyangmeigui	China	III
	71	Augusta	Romania	III
	68	Miskat Plevenski	Bulgaria	III
	61	Pobeda	Uzbekistan	III
	69	Flaming Muscat	South Africa	III
Unknown	56	MZL-5	Songxian, Henan Province, R6 ^y	III
Unknown	55	MZL-6	Songxian, Henan Province, R6 ^y	III
<i>V. betulifolia</i>	58	BYS	Songxian, Henan Province, R6 ^y	III
	57	SX	Songxian, Henan Province, R6 ^y	III
<i>V. ficifolia</i>	45	QTH-6	Jiaozuo, Henan Province, R6 ^y	IV
	44	NWH1403	Xinyang, Henan Province, R6 ^y	IV
	41	QYS-8	Luoyang, Henan Province, R6 ^y	IV
	43	JLG-4	Luoyang, Henan Province, R6 ^y	IV
	42	JLG-1	Luoyang, Henan Province, R6 ^y	IV
	51	BTM-1	Nanyang, Henan Province, R6 ^y	IV
	53	FJG-3	Luoyang, Henan Province, R6 ^y	IV
	52	XY	Xinyang, Henan Province, R6 ^y	IV
	47	QL-3	Shaanxi Province, R5 ^y	IV
	46	HX	Xian, Shaanxi Province, R5 ^y	IV
	48	WDS-1	Henan Province, R6 ^y	IV
	50	SBY-5	Linzhou, Henan Province, R6 ^y	IV
	49	JLG1401	Luoyang, Henan Province, R6 ^y	IV

Continued next page

Table 1. Continued.

Species	Code no.	Clone	Origin	Cluster no.
<i>V. romaneti</i>	26	BTM	Nanyang, Henan Province, R6 ^y	IV
	27	LB-qiu	Henan Province, R6 ^y	IV
	38	ZX1734	Hubei Province, R6 ^y	IV
<i>V. bellula</i>	3	MLPT1104	Hubei Province, R6 ^y	IV
<i>V. pseudoreticulata</i>	36	GX	Guangxi Province, R2 ^y	V
	37	LY	Jiangsu Province, R2 ^y	V
	4	WX1135	Chongqing, R6 ^y	IV
	5	HDPT1057	Hubei Province, R6 ^y	IV
	7	LYW-1	Luoyang, Henan Province, R6 ^y	IV
<i>V. piasezkii</i>	10	LJS-1	Luoyang, Henan Province, R6 ^y	IV
	77	SNJ	Hubei Province, R6 ^y	III
	40	CHS1401	Xian, Shaanxi Province, R5 ^y	IV
<i>V. shenxiensis</i>	15	LB-shanxi	Henan Province, R6 ^y	IV
	12	DW1707	Hubei Province, R6 ^y	V
<i>V. wuhanensis</i>	9	XN1710	Hubei Province, R6 ^y	V
	39	WH-3	Wuhan Province, R6 ^y	V
<i>V. bryoniaefolia</i>	6	WD1418	Wuhan Province, R6 ^y	V
	21	QYS-1	Luoyang, Henan Province, R6 ^y	V
	1	XGPT-2	Guangxi Province, R2 ^y	V
<i>V. balanseana</i>	2	NN1606	Guangxi Province, R2 ^y	V
	13	YC-1	Chongqing, R1 ^y	V
<i>V. heyneana</i>	14	YMN-1	Yuanmou, Yunnan Province, R1 ^y	V
	18	YMCB	Yuanmou, Yunnan Province, R1 ^y	V
	54	NWH1402	Xinyang, Henan Province, R1 ^y	IV
	25	YM-2	Yunnan Province, R1 ^y	V
<i>V. yunnanensis</i>	28	YM-3	Yunnan Province, R1 ^y	V
	29	YN-hn	Hunan Province, R2 ^y	V
	30	YN-2	Yunnan Province, R1 ^y	V
	24	XS1614	Jiangxi Province, R2 ^y	V
<i>V. chungii</i>	23	SQS	Jiangxi Province, R2 ^y	V
<i>V. chunganensis</i>	19	WYS1709	Fujian Province, R2 ^y	V
<i>V. hancockii</i>	20	LYPT0945	Jiangxi Province, R2 ^y	V
<i>V. sinocinerea</i>	8	CS	Changsha, Hunan Province, R6 ^y	V
<i>V. wilsonae</i>	34	BTM-2	Nanyang, Henan Province, R6 ^y	V
	35	BTM-1	Nanyang, Henan Province, R6 ^y	V
	31	LS-2	Luoyang, Henan Province, R6 ^y	V
	33	LS-1	Luoyang, Henan Province, R6 ^y	V
<i>V. adenoclada</i>	16	SX-1	Hongjiang, Hunan Province, R6 ^y	V
	17	GX4020101	Guangxi Province, R2 ^y	V
<i>V. hekouensis</i>	22	HKPT	Yunnan Province, R1 ^y	V
<i>V. ruyuanensis</i>	11	HZ199	Wuhan Province, R6 ^y	V
Unknown	32	Wuzhi-1	Wuhan Province, R6 ^y	V

^zSauvignon Blanc, Carignan, Gamay, and Syrah are used for simple sequence repeat standardization for comparison with international databases.

^yR1 = southwestern China (i.e., Yunnan and Chongqing); R2 = southeastern China (i.e., Guangxi, Guangdong, Jiangxi, Jiangsu, and Fujian); R3 = north China (i.e., Beijing); R4 = northeastern China (i.e., Heilongjiang); R5 = northwestern China (i.e., Shanxi and Qinling); R6 = central China (i.e., Hubei, Hunan, Henan).

(Table 2). H_o ranged from 0.476 to 0.859, with a mean value of 0.644; H_e ranged from 0.503 to 0.726, with a mean value of 0.608 (Table 2). The fixation index (F) values were close to zero for most of the loci, indicating that the locus fit Hardy-Weinberg equilibrium. The gene flow values of all the SSR loci were less than one, revealing that the allelic variation at different SSR loci showed genetic differentiation among the different materials.

According to the cluster analysis, 100 grape samples could be divided into five groups. The genetic diversity of the

different groups (I, II, III, IV, and V) evaluated using the SSR markers is shown in Table 3. According to the values for N_e and I, group V had the highest diversity, followed by groups IV, III, I, and II, whereas the number of accessions in groups III and II are more than group IV and I, respectively, which indicates a little relationship with the number of accessions in each group.

To understand the genetic variation in grape accessions from different origins, genetic diversity was also evaluated with the SSR data (Table 4). Central China (R6) had the highest values

Table 2. Genetic diversity analysis of the 100 *Vitis* accessions by the nine simple sequence repeat markers.

Locus	Parameter ^z								
	N _a	N _e	H _o	H _e	GD	PIC	I	F	Nm
VrZAG62	20	10.959	0.637	0.646	0.909	0.902	1.319	-0.023	0.608
VrZAG79	18	11.138	0.691	0.610	0.911	0.904	1.274	-0.142	0.489
VVS2	23	13.441	0.859	0.726	0.927	0.923	1.582	-0.216	0.937
VVMD5	19	8.306	0.582	0.544	0.883	0.873	1.102	-0.091	0.413
VVMD7	16	9.268	0.515	0.503	0.894	0.885	0.966	-0.057	0.309
VVMD28	29	17.549	0.622	0.626	0.943	0.940	1.358	0.003	0.480
VVMD32	22	12.107	0.476	0.543	0.918	0.912	1.051	0.099	0.369
VVMD25	21	10.455	0.751	0.667	0.906	0.899	1.358	-0.171	0.675
VVMD27	18	10.030	0.662	0.605	0.899	0.891	1.197	-0.101	0.539
Mean	20.667	11.472	0.644	0.608	0.910	0.903	1.245	-0.078	0.535

^zN_a = observed number of alleles; N_e = effective number of alleles; H_o = observed heterozygosity; H_e = expected heterozygosity; GD = gene diversity; PIC = polymorphism information content; I = Shannon's information index; F = average fixation index; Nm = gene flow estimate according to Wright's equation.

Table 3. The parameters of genetic diversity among five *Vitis* groups estimated by nine simple sequence repeat markers.

Group no.	Parameter ^z					Total genotypes (no.)
	N _a	N _e	I	H _o	H _e	
I	5.556	4.433	1.473	0.733	0.700	5
II	5.667	3.455	1.345	0.744	0.650	13
III	11.556	5.764	2.000	0.798	0.816	28
IV	12.556	7.034	2.141	0.755	0.838	24
V	14.667	8.503	2.362	0.566	0.878	30

^zN_a = observed number of alleles; N_e = effective number of alleles; I = Shannon's information index; H_o = observed heterozygosity; H_e = expected heterozygosity.

Table 4. The parameters of genetic diversity of *Vitis* from different regions estimated by nine simple sequence repeat markers.

Origin ^z	Parameter ^y						Total genotypes (no.)
	N _a	N _e	I	H _o	H _e	F	
R1	7.778	5.819	1.783	0.611	0.775	0.028**	8
R2	9.444	6.861	2.052	0.640	0.850	0.248**	13
R3	2.667	2.415	0.866	0.722	0.528	-0.378	2
R4	3.111	2.690	0.973	0.667	0.556	-0.187	3
R5	3.667	3.003	1.126	0.778	0.611	-0.272**	3
R6	16.778	10.002	2.469	0.692	0.891	0.225***	48

^zR1 = southwestern China (i.e., Yunnan and Chongqing); R2 = southeastern China (i.e., Guangxi, Guangdong, Jiangxi, Jiangsu, and Fujian); R3 = North China (i.e., Beijing); R4 = northeastern China (i.e., Heilongjiang); R5 = northwestern China (i.e., Shanxi and Qinling); R6 = Central China (i.e., Hubei, Hunan, Henan).

^yN_a = observed number of alleles; N_e = effective number of alleles; I = Shannon's information index; H_o = observed heterozygosity; H_e = expected heterozygosity; F = average fixation index.

, *Significant at $P \leq 0.10$ or 0.05 , respectively, calculated over 1000 permutations.

of the major parameters N_e and I, followed by southeastern China (R2), southwestern China (R1), northwestern China (R5), northeastern China (R4), and north China (R3), and this trend showed that the genetic diversity of the six regions varied, with the accessions derived from central China exhibiting the greatest genetic diversity. This result may be related to the sample size of each region.

POPULATION DIFFERENTIATION ANALYSIS. AMOVA was performed among the 100 accessions by considering the five populations generated by the structure analysis (Table 5). The AMOVA results in Chinese wild grape accessions indicated that 3.28% of the total variance was among populations, 3.27% was among individuals within populations, and 93.46% was within individuals. The inbreeding coefficients within subpopulations relative to the total, within individuals relative to the subpopulation, and within individuals relative to the total parameters were 0.033, 0.034, and 0.065, respectively (Table 5).

PRINCIPAL COORDINATE ANALYSIS. PCA was chosen to complement the cluster analysis information and genetic relationships (Zhu et al., 2018). The PCA results for 100 accessions based on the genetic distance matrix data from the nine SSR markers is shown in Fig. 1. Generally speaking, 100 accessions could be divided into three primary groups (i.e., North American species, European species, and East Asian species), although there was little overlap between the European species and East Asian species, which is consistent with the geography [i.e., North America, East Asia, and Europe (Fig. 1)].

CLUSTER ANALYSIS. To assess the genetic diversity and relationships among 100 *Vitis* materials, a cluster analysis was conducted based on the genetic distance matrix data estimated by using SSR markers. The dendrogram constructed by UPGMA is shown in Fig. 2. In addition, the North American accessions separated from the other cultivars, forming one individual group—namely, group I—which showed there was a marked divergence between the North American accessions and the Eurasian species of the subgenus *Vitis*, whereas the cultivated grapes (European species) and Chinese wild grape species (East Asian species) were grouped together in group III. These results provide additional support for the sister relationship between the East Asian species and the European species.

All 100 grape accessions examined were divided into five groups based on the cluster analysis. Group I contained exclusively North American accessions; group II contained only *V. davidii*. Group III was subdivided into three subgroups. Group III-1 contained *V. amurensis* and *V. amurensis* var. *yan-sha-nensis*, and one accession belonged to *V. piasezkii*. Group III-2 contained European cultivars. Group III-3 included *V. betulifolia*, and two accessions shared similar morphologies with *V. betulifolia*. Group IV consisted of two subgroups. Group IV-1 included *V. ficifolia*. Group IV-2 contained *V. heyneana*, *V.*

Table 5. Results of the analysis of molecular variance carried out among 100 *Vitis* accessions considering the five populations generated by the structure analysis.

Source of variation	df	Sum of squares	Variance components	Proportion of variation (%)	Coefficient ^z		
					F _{st}	F _{is}	F _{it}
Among populations	4	36.102	0.135	3.28			
Among individuals within populations	95	390.298	0.134	3.27			
Within individuals	100	384.000	3.840	93.46			
Total	199	810.400	4.109	100	0.033	0.034	0.065

^zF_{st} = the inbreeding coefficient within subpopulations relative to the total; F_{is} = the inbreeding coefficient within individuals relative to the subpopulation; F_{it} = the inbreeding coefficient within individuals relative to the total.

Fig. 1. Associations among the 100 *Vitis* accessions revealed by principal coordinate (PC) analysis as performed on Nei's genetic distance matrix data calculated from nine simple sequence repeat markers. The red circle represents the North American species. The green circle represents the East Asian species. The purple circle represents the European species.

shenxiensis, *V. piasezkii*, and *V. romaneti*. Group V was the largest, with 30 accessions, and included 15 wild species native to China. Group V was subdivided into 12 subgroups. Group V-1 included the single species *V. wuhanensis*. Group V-2 included *V. bryoniaefolia*. Group V-3 contained *V. balanseana*, *V. pseudoreticulata*, and *V. bellula*. Group V-4 included a single species, *V. yunnanensis*. Group V-5 contained *V. chungii*. Group V-6 consisted of *V. chunganensis*. Group V-7 contained a single species, *V. hancockii*. Group V-8 contained only *V. sinocinerea*. Group V-9 included two accessions of *V. pseudoreticulata*, and four accessions of *V. pseudoreticulata* did not group closely within one branch. Group V-10 contained *V. wilsonae* and one accession, Wuzhi-1. Group V-11 consisted of *V. adenoclada* and *V. hekouensis*. Group V-12 contained only *V. ruyuanensis*.

In addition, some important exceptions were discovered by cluster analysis. Three previously undescribed accessions [Wuzhi-1 (ID 32), MZL-5 (ID 56), and MZL-6 (ID 55)] were assigned to the *V. wilsonae* and *V. betulifolia* cluster, with mean genetic distance values of 0.39, 0.31, and 0.38, respectively.

POPULATION STRUCTURE ANALYSIS. The statistic pK calculated with STRUCTURE was used to determine a suitable K value (population number). The maximum likelihood value and a sharp peak were revealed at K = 5. Thus, the 100 grape accessions were divided into five populations: POP I, POP II,

POP III, POP IV, and POP V (Fig. 3). On the whole, the results of the genetic structure analysis were consistent with those revealed by the cluster analysis. POP I contained five accessions that originated only from group I, and the accessions of POP I were from the US. POP II included 13 accessions, all of which were from group II, and the accessions of this population were from Hunan, Hubei, Jiangxi, and Wuhan. The 24 accessions that originated from group III were assigned to POP III, which were mostly from Europe and Heilongjiang, among others. POP IV consisted of 18 accessions, which were mostly from Henan, including 14 accessions from group IV, and the other four accessions that originated from group III. POP V included 40 accessions containing all 30 accessions from POP V along with 10 accessions from group IV, which were from Yunnan, Hunan, Jiangxi, Guangxi, and Fujian, among others.

Discussion

SSR MARKER. The SSR marker is a powerful, reliable, and useful tool for molecular breeding and for assessing the genetic diversity of plants because of its codominance and ability to reveal a high number of alleles per locus (Bassil et al., 2018; El-Esawi et al., 2016; Liang et al., 2015; Liu et al., 2015; Rodolfi et al., 2018). This is the first study to assess the genetic diversity and relationships among 23 wild Chinese *Vitis* species/cultivars using this powerful microsatellite technique. The nine SSR markers selected in this study, which are high in polymorphisms, have been used by the European GrapeGen06 consortium (Maul et al., 2012) and have been used frequently to identify grape cultivars (Li et al., 2018; Mihaljevic et al., 2013; This et al., 2004). Internationally, France, Germany, Italy, and other countries have used the nine SSR markers to establish a molecular database of grape cultivars (Maul and Töpfer, 2015), providing an inquiry service for grape research workers. Furthermore, Li et al. (2017, 2018) showed that the nine SSR markers can be used to identify the Chinese grape landraces and Euro-American hybrids.

Fig. 2. Phylogenetic trees of 100 *Vitis* accessions based on simple sequence repeat data. A distance tree was constructed in MEGA5 (Tamura et al., 2011) using the unweighted pair-group method of mathematical averages method. Code number is the same reported in Table 1. POP = population.

The average allele per marker (20.667) is greater than that noted in the earlier report by Jing et al. (2013), who found an average of 10.5 alleles per marker in 62 Chinese wild grapes using 10 SSR markers. The PIC value (0.903) is greater than the values reported by Karataş et al. (2014) (SSR, 0.63) and Ramezani et al. (2009) (SSR, 0.89) in grapes. In particular, the PIC value in our study (0.903) is greater than the earlier report by Li et al. (2018), who found an average PIC value of 0.815 in 94 cultivated accessions using the nine SSR markers. The greater PIC values evidence a greater genetic diversity in the currently studied Chinese wild grape species than in the

cultivated accessions that were studied before. In addition, we found greater H_o values than expected (0.644 and 0.608, respectively), which is in accordance with Riaz et al. (2018), who reported greater H_o values than expected among 1378 grape accessions (0.742 and 0.678, respectively). The high heterozygosity values found our study suggest high levels of genetic variation among the analyzed Chinese wild grape species. Currently, many wild grapes have been used in breeding programs, and some even were applied in production directly (Liu et al., 2014). Therefore, this set of SSR primers can be used for the identification of wild grape cultivars in the grape industry.

Fig. 3. Genetic structure of the 100 *Vitis* accessions inferred by a Bayesian mixed-model-based cluster analysis assuming correlations among allele frequencies across clusters. Each individual is represented by a thin line partitioned into K colored segments representing the membership fraction in K clusters. Each color represents a population. POP = population; K = number of populations.

SSR markers are also a powerful tool for characterizing the cultivars. In this study, cluster, principal coordinate, and structure analyses allowed us to identify three previously undescribed accessions (Wuzhi-1, MZL-5, and MZL-6). For instance, Wuzhi-1, MZL-5, and MZL-6 not only have a closer relationship with *V. betulifolia* at the molecular level, but also have similar morphological characteristics. The branchlets have arachnoid tomentum and fall off later; the shape of the mature leaf is oval and the number of mature leaf lobes is three; the lower side prostrate hairs between the veins of young leaves are sparse; and there are no glandular hairs on shoots. Bassil et al. (2018) also used SSRs to identify four accessions of previously undescribed hybrid origin during an analysis of the genetic diversity of wild *Vaccinium* species. Cao et al. (2020) reported that five previously unknown grape accessions were matched by SSRs.

PATTERN OF GENETIC DIVERSITY DISTRIBUTION AMONG DIFFERENT POPULATIONS. The genetic diversity within and among the different geographic groups in Chinese wild grape species, as demonstrated by I and N_e , indicates there is significant diversity (Table 4). The *Vitis* accessions from R6 (i.e., Hubei, Hunan, and Henan) had the highest number of effective alleles, and the I value suggests this region may be the center of diversity for wild grape species. These results may be a result of the large sample size for this region. The H_o value of the accessions from R1, R2, and R6 was slightly less than the H_e values, with a contrasting trend for the accessions from R3, R4, and R5. These differences are consistent with the positive F values, particularly in the populations from R2 and R6, which indicates a close genetic relationship among individuals from the same populations (Table 4). The negative F values for accessions from R2, R3, and R6 indicate an excess of heterozygotes, but only the last one (R6), was statistically significant (-0.272) (Table 4).

The AMOVA results showed that similar levels of genetic variation among and within populations were observed, and low levels of genetic diversity were present among populations, whereas high levels of genetic diversity were found within individuals (Table 5). These results are in accordance with the findings from Riaz et al. (2018), who analyzed the genetic diversity of cultivated and wild grapevine accessions around the Mediterranean Basin and Central Asia. Riaz et al. (2018) also reported that the level of genetic diversity within individuals was greater than that among populations.

GENETIC STRUCTURE AND DIFFERENTIATION. A significant differentiation between wild North American *Vitis* species and Eurasian *Vitis* species was detected by cluster analysis and PCA (Figs. 1 and 2). This result is consistent with previous studies that support the division of the subgenus *Vitis* into two clades that mirror their geographic distribution—North America and Eurasia—based on resequencing technology, plastid and nuclear markers, and so on (Klein et al., 2018; Liu et al., 2016; Ma et al., 2018; Péros et al., 2011; Wan et al., 2013). Nevertheless, the relationship between European and East Asian species remains controversial.

In our study, the European species and the cultivated accessions native to China were grouped with two Chinese wild grape species, indicating these Chinese wild grape species are more genetically close to European grapes than North American species. On the other hand, these results were not consistent with the findings of Liang et al. (2019), who indicated that wild East Asian and wild North American accessions shared more similarity in genetic background than Eurasian accessions based on the whole-genome resequencing method. This finding is consistent with those of Liu et al. (2016) and Zhang et al. (2018), all of whom affirmed that the European species (*V. vinifera*) was grouped within the East Asian clade. Many explanations have been made regarding this phenomenon. For example, Péros et al. (2011) held the opinion that there might be an Asian origin for the subgenus *Vitis*, and their study suggested that the subgenus *Vitis* first diverged in Asia and later colonized Europe. Moreover, one hypothesis was suggested that hybridization could be conducive to the integration of the European grape genome into the Chinese germplasm (Zhang et al., 2018). This finding is based on the observation that viticulture has been common since ancient times in regions where wild species grew as well. This finding will increase the gene flow possibility between the cultivated grapes (*V. vinifera*) and the wild grape species.

Furthermore, 100 *Vitis* accessions were divided into five major groups by their SSRs (Fig. 2), and the PCA also showed similar groupings. The reason that group V exhibited greater genetic diversity might be related to the sample size. In addition, the American *Vitis* species from the US grouped together and formed group I. Group II was distributed primarily in the south-central and southeastern regions of China. Groups III and IV were distributed primarily throughout the central region, with some extending to the northern, northeastern, and

northwestern regions of China, whereas the accessions that originated from Europe grouped together and were also assigned to group III. Group V was distributed primarily throughout the southwestern and southeastern regions of China. Thus, the clustering patterns of most accessions correlated with geographic distribution (Fig. 2).

In addition, a population structure analysis showed that five populations were obtained from the 100 *Vitis* accessions. However, there were a few discrepancies between the cluster analysis and the structure analysis, which might be because each accession was assigned a fixed branch by cluster analysis, whereas the structural analysis assigned individuals to groups based on the percentage of population members (Zhu et al., 2018). The admixture of all the populations might indicate gene flow in these accessions (Aradhya et al., 2013).

In addition, the results of the cluster analysis revealed a close relationship among *V. piasezkii*, *V. romaneti*, *V. shenxiensis*, and *V. heyneana*. This finding is similar to the results of Ma et al. (2018). Our result supports the idea that some types of *V. piasezkii* have similar morphological characteristics relative to *V. shenxiensis* (Ren and Wen, 2007), which also indicates that a high level of genetic diversity is present within *V. piasezkii*. In addition, according to the review by Wan et al. (2008), *V. romaneti* and *V. shenxiensis* belong to section *Romanetianae*. *V. heyneana* did not cluster with *V. ficifolia*, unlike the results reported by Zhang et al. (2018) and Wang et al. (2008), but consistent with those of Liu et al. (2016). In addition, the results also revealed a closer relationship between *V. wilsonae* and two accessions of *V. pseudoreticulata*, and this finding is similar to the results of the phylogenetic studies on the grape genus using five plastid and two nuclear markers (Liu et al., 2016). We also found that some accessions belonging to the same species are in different clades (Fig. 2), such as *V. pseudoreticulata* and *V. piasezkii*. This result implies high genetic variation and genetic diversity within species, which was further supported by AMOVA.

Conclusion

Today, China holds more than 3000 accessions of *Vitis*, of which some are wild types. These accessions are either native to China or through foreign introduction. With the increase of grape germplasms, genetic information among accessions is becoming more critical for maintenance and utilization strategies in China's breeding programs. We conclude that the estimation of genetic diversity and the population structure of 100 accessions of *Vitis* using SSR markers may provide more accurate information to grape breeders than the classical pedigree method. In addition, the nine SSR markers used in our study are significantly polymorphic and can be used efficiently to distinguish wild Chinese grape species. Moreover, three previously undescribed accessions were identified to species (Wuzhi-1, MZL-5, and MZL-6). The nine SSR primer pairs in our study may also be of potential value for further research on genetic mapping, marker-assisted selections, and so on, in grapes.

Literature Cited

- Ali, A., Y.B. Pan, Q.N. Wang, J.D. Wang, J.L. Chen, and S.J. Gao. 2019. Genetic diversity and population structure analysis of *Saccharum* and *Erianthus* genera using microsatellite (SSR) markers. *Sci. Rep.* 9:395, doi: 10.1038/s41598-018-36630-7.
- Abbasov, M., Z. Akparov, T. Gross, S. Babayeva, V. Izzatullayeva, E. Hajiyev, K. Rustamov, P. Cross, M. Tekin, T. Akar, S.M. Chao, and R. Brueggeman. 2018. Genetic relationship of diploid wheat (*Triticum* sp.) species assessed by SSR markers. *Genet. Resources Crop Evol.* 65:1441–1453, doi: 10.1007/s10722-018-0629-2.
- Aradhya, M., Y. Wang, M.A. Walker, B.H. Prins, A.M. Koehmstedt, D. Velasco, and J.E. Preece. 2013. Genetic diversity, structure, and patterns of differentiation in the genus *Vitis*. *Plant Syst. Evol.* 299:317–330, doi: 10.1007/s00606-012-0723-4.
- Bassil, N., A. Bidani, K. Hummer, L.J. Rowland, J. Olmstead, P. Lyrene, and C. Richards. 2018. Assessing genetic diversity of wild southeastern North American *Vaccinium* species using microsatellite markers. *Genet. Resources Crop Evol.* 65:939–950, doi: 10.1007/s10722-017-0585-2.
- Bowers, J., J.M. Boursiquot, P. This, K. Chu, H. Johansson, and C. Meredith. 1999. Historical genetics: The parentage of chardonnay, gamay, and other wine grapes of northeastern France. *Science* 285:1562–1565, doi: 10.1126/science.285.5433.1562.
- Bowers, J.E., G.S. Dangl, R. Vignani, and C.P. Meredith. 1996. Isolation and characterization of new polymorphic simple sequence repeat loci in grape (*Vitis vinifera* L.). *Genome* 39:628–633, doi: 10.1139/g96-080.
- Cao, S., S. Stringer, G. Gunawan, C. McGregor, and P.J. Conner. 2020. Genetic diversity and pedigree analysis of muscadine grape using SSR markers. *J. Amer. Soc. Hort. Sci.* 145:143–151, doi: 10.21273/JASHS04856-20.
- Chao, W.J. and S.Z. Niu. 1981. Preliminary study on isoenzyme of Chinese wild *Vitis* in China. *China Fruits* 4:41–44, doi: 10.16626/j.cnki.issn1000-8047.1981.04.022. (in Chinese).
- Earl, D.A. and B.M. Vonholdt. 2012. Structure Harvester: A website and program for visualizing structure output and implementing the evanno method. *Conserv. Genet. Resources* 4:359–361, doi: 10.1007/s12686-011-9548-7.
- El-Esawi, M.A., K. Germaine, P. Bourke, and R. Malone. 2016. Genetic diversity and population structure of *Brassica oleracea* germplasm in Ireland using SSR markers. *C. R. Biol.* 339:133–140, doi: 10.1016/j.crv.2016.02.002.
- Fan, X.C., J.F. Jiang, Y. Zhang, H.S. Sun, J. Jiao, and C.H. Liu. 2015. Genetic diversity assessment of *Vitis ficifolia* Bge. populations from Henan Province of China by SRAP markers. *Biotechnol. Biotech. Equipment* 29:15–20, doi: 10.1080/13102818.2014.984414.
- Guan, C., P. Zhang, C. Hu, S. Chachar, A. Riaz, R. Wang, and Y. Yang. 2019. Genetic diversity, germplasm identification and population structure of *Diospyros kaki* Thunb. from different geographic regions in China using SSR markers. *Scientia Hort.* 251:233–240, doi: 10.1016/j.scientia.2019.02.062.
- Guo, D.L., J.Y. Zhang, C.H. Liu, G.H. Zhang, and M. Li. 2010. Genetic relationships of Chinese grape accessions to European and American cultivars assessed by microsatellite markers. *Biotechnol. Biotech. Equipment* 24:2054–2059, doi: 10.2478/v10133-010-0073-8.
- He, P.C. and W.J. Chao. 1982. Study on pollen electron microscopy of Chinese wild *Vitis*. *China Fruits* 3:43–45, doi: 10.7606/j.1004-1389.2014.04.013. (in Chinese).
- He, Z., H. Zhang, S. Gao, M.J. Lercher, W.H. Chen, and S. Hu. 2016. Evolvview v2: An online visualization and management tool for customized and annotated phylogenetic trees. *Nucl. Acids Res.* 44:W236–W241, doi: 10.1093/nar/gkw370.
- Jiang, J.F., S. Kell, X.C. Fan, Y. Zhang, W. Wei, D.M. Kang, N. Maxted, B.F. Lloyd, and C.H. Liu. 2015. The wild relatives of grape in China: Diversity, conservation gaps and impact of climate change. *Agr. Ecosyst. Environ.* 209:155–163, doi: 10.1016/j.agee.2015.03.021.
- Jing, Z.B. and X.P. Wang. 2013. Genetic relationship between Chinese wild *Vitis* species and American and European cultivars based on ISSR markers. *Biochem. Syst. Ecol.* 46:120–126, doi: 10.1016/j.bse.2012.08.004.
- Jing, Z.B., X.P. Wang, and J.M. Cheng. 2013. Analysis of genetic diversity among Chinese wild *Vitis* species revealed with SSR and SRAP markers. *Genet. Mol. Res.* 12:1962–1973, doi: 10.4238/2013.June.13.5.

- Karataş, D.D., H. Karataş, V. Laucou, G. Sarikamiş, L. Riahi, R. Bacilieri, and P. This. 2014. Genetic diversity of wild and cultivated grapevine accessions from southeast Turkey. *Hereditas* 151:73–80, doi: 10.1111/hrd2.00039.
- Kejun, L. and S.V. Muse. 2005. Powermarker: An integrated analysis environment for genetic marker analysis. *Bioinformatics* 9:2128–2129, doi: 10.1093/bioinformatics/bti282.
- Klein, L.L., A.J. Miller, C. Ciotir, K. Hyma, S. Uribe-Convers, and J. Londo. 2018. High-throughput sequencing data clarify evolutionary relationships among North American *Vitis* species and improve identification in USDA *Vitis* germplasm collections. *Amer. J. Bot.* 105:215–226, doi: 10.1002/ajb2.1033.
- Krawczak, M., S. Nikolaus, H. Von Eberstein, P.J.P. Croucher, N.E. El Mokhtari, and S. Schreiber. 2006. Popgen: Population-based recruitment of patients and controls for the analysis of complex genotype–phenotype relationships. *Community Genet.* 9:55–61, doi: 10.1159/000090694.
- Li, B.B., J.F. Jiang, X.C. Fan, Y. Zhang, H.S. Sun, G.H. Zhang, and C.H. Liu. 2017. Molecular characterization of Chinese grape landraces (*Vitis* L.) using microsatellite DNA markers. *HortScience* 52:533–540, doi: 10.21273/HORTSCI11802-17.
- Li, B.B., J.F. Jiang, Y. Zhang, X.C. Fan, H.S. Sun, G.H. Zhang, and C.H. Liu. 2018. DNA fingerprinting and genetic diversity analysis of grape cultivars based on SSR markers. *J. Plant Genet. Resources* 19:301–313, doi: 10.13430/j.cnki.Jpgr.2018.02.017. (in Chinese).
- Li, Y., K. Cao, G.R. Zhu, W.C. Fang, C.W. Chen, X.W. Wang, P. Zhao, J. Guo, T.Y. Ding, L.P. Guan, Q. Zhang, W.W. Guo, Z.J. Fei, and L.R. Wang. 2019. Genomic analyses of an extensive collection of wild and cultivated accessions provide new insights into peach breeding history. *Genome Biol.* 20:36, doi: 10.1186/s13059-019-1648-9.
- Liang, W., L. Dondini, P.D. Franceschi, R. Paris, S. Sansavini, and S. Tartarini. 2015. Genetic diversity, population structure and construction of a core collection of apple cultivars from Italian germplasm. *Plant Mol. Biol. Rpt.* 33:458–473, doi: 10.1186/s13059-019-1648-9.
- Liang, Z., S. Duan, J. Sheng, S. Zhu, X. Ni, J. Shao, and R. Mao. 2019. Whole-genome resequencing of 472 *Vitis* accessions for grapevine diversity and demographic history analyses. *Nat. Commun.* 10:1190, doi: 10.1038/s41467-019-09135-8.
- Liu, C.H., X.C. Fan, J.F. Jiang, D.L. Guo, H.S. Sun, Y. Zhang, and J.G. Feng. 2012. Genetic diversity of Chinese wild grape species by SSR and SRAP markers. *Biotechnol. Biotech. Equipment* 26:2899–2903, doi: 10.5504/bbeq.2011.0146.
- Liu, X.Q., S.M. Ickert-Bond, Z.L. Nie, Z. Zhou, L.Q. Chen, and J. Wen. 2016. Phylogeny of the *Ampelocissus*–*Vitis* clade in Vitaceae supports the New World origin of the grape genus. *Mol. Phylogenet. Evol.* 95:217–228, doi: 10.1016/j.ympev.2015.10.013.
- Liu, C.H., J.F. Jiang, X.C. Fan, and Y. Zhang. 2014. The utilization of Chinese wild grape species in production and breeding. *J. Plant Genet. Resources* 15:720–727, doi: 10.13430/j.cnki.Jpgr.2014.04.006. (in Chinese).
- Liu, Q., Y. Song, L. Liu, M. Zhang, J. Sun, S. Zhang, and J. Wu. 2015. Genetic diversity and population structure of pear (*Pyrus* spp.) collections revealed by a set of core genome-wide SSR markers. *Tree Genet. Genomes* 11:1–22, doi: 10.1007/s11295-015-0953-z.
- Liu, K.Y., F. Xu, X.H. Shi, X.H. Zhong, G.F. Yang, and J.J. Ni. 2013. Relationship analysis for *Vitis davidii* with SRAP molecular markers. *J. Hunan Agr. Univ.* 38:607–611, doi: 10.3724/SP.J.1238.2012.00607. (in Chinese).
- Luo, S.L., P.C. He, X.Q. Zheng, and P. Zhou. 2001. Genetic diversity in wild grapes native to China based on randomly amplified polymorphic DNA (RAPD) analysis. *Acta Bot. Sin.* 43:158–163, doi: 10.1016/S0168-9452(00)00427-1.
- Ma, Z.Y., J. Wen, S.M. Ickert-Bond, Z.L. Nie, L.Q. Chen, and X.Q. Liu. 2018. Phylogenomics, biogeography, and adaptive radiation of grapes. *Mol. Phylogenet. Evol.* 129:258–267, doi: 10.1016/j.ympev.2018.08.021.
- Maul, E., K.N. Sudharma, S. Kecke, G. Marx, and P. This. 2012. The European *Vitis* database (www.eu-vitis.de): A technical innovation through an online uploading and interactive modification system-*Vitis27985*, doi: 10.5073/VITIS.2012.51.79-85.
- Maul, E. and R. Töpfer. 2015. *Vitis International Variety Catalogue (VIVC): A cultivar database referenced by genetic profiles and morphology.* Bio. Web Conf. 5:01009, doi: 10.1051/bioconf/20150501009.
- Mihaljevic, M.Z., S. Simon, I. Pejic, F. Carka, R. Sevo, A. Kojic, and D. Preiner. 2013. Molecular characterization of old local grapevine varieties from southeast European countries. *Vitis* 52:69–76, doi: 10.5073/vitis.2013.52.69-76.
- Nei, M. 1973. Analysis of gene diversity in subdivided populations. *Proc. Nat. Acad. Sci. USA* 70:3321–3323, doi: 10.1073/pnas.70.12.3321.
- Peakall, R. and P.E. Smouse. 2006. GENALEX 6: Genetic analysis in Excel: Population genetic software for teaching and research. *Mol. Ecol. Notes* 6:288–295, doi: 10.1111/j.1471-8286.2005.01155.x.
- Péros, J.P., G. Berger, A. Portemont, J.M. Boursiquot, and T. Lacombe. 2011. Genetic variation and biogeography of the disjunct *Vitis* subg. *Vitis* (Vitaceae). *J. Biogeogr.* 38:471–486, doi: 10.1111/j.1365-2699.2010.02410.x.
- Queiroz, A., A. Assunção, I. Ramadas, W. Viegas, and M.M. Veloso. 2015. Molecular characterization of Portuguese pear landraces (*Pyrus communis* L.) using SSR markers. *Scientia Hort.* 183:72–76, doi: 10.1016/j.scienta.2014.11.016.
- Ramezani, A., R. Haddad, M. Dorostkar, M. Mardi, and M. Naghavi. 2009. Evaluation of genetic diversity of Iranian grapevine accessions using microsatellite markers. *Vitis* 48:151–152, doi: 10.1007/s00122-011-1585-9.
- Ren, H. and J. Wen. 2007. *VITIS Linnaeus*, p. 210–222. In: H. Peng (ed.), *Flora of China* (English edition). Science Press, Beijing, China.
- Riaz, S., G. De Lorenzis, D. Velasco, A. Koehmstedt, D. Maghradze, Z. Bobokashvili, and O. Failla. 2018. Genetic diversity analysis of cultivated and wild grapevine (*Vitis vinifera* L.) accessions around the Mediterranean Basin and Central Asia. *BMC Plant Biol.* 18:137, doi: 10.1186/s12870-018-1351-0.
- Rodolfi, M., A. Silvanini, B. Chiancone, M. Marieschi, A. Fabbri, R. Bruni, and T. Ganino. 2018. Identification and genetic structure of wild Italian *Humulus lupulus* L. and comparison with European and American hop cultivars using nuclear microsatellite markers. *Genet. Resources Crop Evol.* 65:1405–1422, doi: 10.1007/s10722-018-0622-9.
- Sefc, K.M., F. Regner, E. Turetschek, J. Glossl, and H. Steinkellner. 1999. Identification of microsatellite sequences in *Vitis* riparia and their applicability for genotyping of different *Vitis* species. *Genome* 42:367–373, doi: 10.1139/g98-168.
- Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28:2731–2739, doi: 10.1093/molbev/msr121.
- This, P., A. Jung, P. Boccacci, J. Borrego, R. Botta, L. Costantini, M. Crespan, G.S. Dengl, C. Eisenheld, F. Ferreira-Monteiro, J. Grando, S. Ibanez, T. Lacombe, V. Laucou, R. Magalhaes, C.P. Meredith, N. Milani, E. Peterlunger, F. Regner, L. Zulini, and E. Maul. 2004. Development of a standard set of microsatellite reference alleles for identification of grape cultivars. *Theor. Appl. Genet.* 109:1448–1458, doi: 10.1007/s00122-004-1760-3.
- Thomas, M.R. and N.S. Scott. 1993. Microsatellite repeats in grapevine reveal DNA polymorphisms when analysed as sequence-tagged sites (STSs). *Theor. Appl. Genet.* 86:985–990, doi: 10.1007/BF00211051.
- Wan, Y., H.R. Schwaninger, A.M. Baldo, J.A. Labate, G.Y. Zhong, and C.J. Simon. 2013. A phylogenetic analysis of the grape genus (*Vitis* L.) reveals broad reticulation and concurrent diversification during neogene and quaternary climate change. *BMC Evol. Biol.* 13:141, doi: 10.1186/1471-2148-13-141.
- Wan, Y., H. Schwaninger, D. Li, C.J. Simon, Y. Wang, and C. Zhang. 2008. A review of taxonomic research in Chinese wild grapes. *Vitis* 47:81–88, doi: 10.5073/vitis.2008.47.81-88.

- Wang, J., C.H. Liu, X.C. Fan, H.S. Sun, and D. Dong. 2008. Study on the biodiversity of the morphology and polymorphism based on SSR markers for *Vitis ficifolia* native to Henan Province in China. *J. Fruit Sci.* 4:496–500, doi: 10.3724/SP.J.1148.2008.00259. (in Chinese).
- Zhao, T.T., W.X. Ma, Q.H. Ma, Z. Yang, and L.J. Wang. 2020. Genetic diversity and population structure of Chinese *Corylus heterophylla* and *Corylus kweichowensis* using simple sequence repeat markers. *J. Amer. Soc. Hort. Sci.* 145:1–10, doi: 10.21273/JASHS04887-19.
- Zhang, X., P. Gong, Y. Shi, Y. Wang, and C. Zhang. 2018. Genetic inter-relationships among Chinese wild grapes based on SRAP marker analyses. *Vitis* 57:151–157, doi: 10.5073/vitis.2018.57.
- Zhang, Y.H., C.H. Liu, X.C. Fan, Y. Zhang, H.S. Sun, B. Peng, and J.F. Jiang. 2011. Application of ISSR markers in classification of wild grapes in China. *J. Fruit Sci.* 28:406–412, doi: 10.13925/j.cnki.gsxb.2011.03.012. (in Chinese).
- Zhou, Y., M. Massonnet, J.S. Sanjak, D. Cantu, and B.S. Gaut. 2017. Evolutionary genomics of grape (*Vitis vinifera* ssp. *vinifera*) domestication. *Proc. Natl. Acad. Sci. USA* 114:11715–11720, doi: 10.1073/pnas.1709257114.
- Zhu, S.Y., X.L. Zhang, Q. Liu, T.K. Luo, and Z. Tang. 2018. The genetic diversity and relationships of cauliflower (*Brassica oleracea* var. *botrytis*) inbred lines assessed by using SSR markers. *PLoS One* 13:e0208551, doi: 10.1371/journal.pone.0208551.

Supplemental Table 1. Genetic profiles of 100 *Vitis* accessions at nine microsatellite loci.

Code no.	Microsatellite loci								
	VrZAG62	VrZAG79	VVS2	VVMD5	VVMD7	VVMD28	VVMD32	VVMD25	VVMD27
1	190/196	248/250	135/139	245/245	259/263	243/249	180/246	236/238	175/179
2	190/196	248/250	135/139	245/245	259/263	249/249	246/246	236/238	175/179
3	188/194	246/260	123/133	243/245	243/251	243/259	236/238	248/250	187/209
4	170/198	256/264	121/131	249/255	243/251	223/235	236/238	260/260	185/197
5	190/194	246/260	123/133	243/245	243/251	245/261	236/238	250/252	187/209
6	182/182	242/252	129/129	245/245	231/247	227/243	220/220	238/238	183/201
7	206/206	242/242	129/129	229/243	237/241	249/255	234/246	234/250	175/179
8	186/188	240/240	133/133	235/235	241/243	255/255	228/228	252/268	181/191
9	182/182	242/254	137/153	239/241	241/247	227/227	240/246	230/242	183/187
10	206/206	242/254	127/127	243/245	237/241	227/249	234/244	234/250	193/193
11	188/190	??	139/143	243/243	??	??	238/238	246/252	181/189
12	182/192	242/242	139/145	233/235	231/231	241/251	234/258	242/242	183/187
13	172/172	234/250	127/155	245/245	231/239	247/255	246/246	238/272	179/181
14	174/174	250/254	129/133	245/245	231/237	241/249	246/246	240/248	179/181
15	194/196	242/244	143/161	239/243	235/237	219/243	234/244	234/258	175/181
16	176/190	252/260	123/131	239/239	245/251	253/253	236/236	232/232	183/183
17	186/194	254/254	125/133	239/239	245/245	247/253	236/236	234/248	187/187
18	174/174	234/250	129/155	245/245	231/237	249/257	246/246	240/274	179/181
19	182/186	236/248	131/135	229/251	233/233	229/229	246/248	236/256	179/199
20	172/186	240/240	137/147	247/247	243/243	229/239	220/220	246/248	201/201
21	194/194	244/250	123/129	245/245	237/237	225/241	218/218	236/236	177/183
22	188/188	236/236	123/123	243/245	245/245	219/219	242/246	234/246	179/179
23	186/186	250/258	121/133	231/233	223/233	229/229	234/234	260/262	175/179
24	174/200	250/256	143/149	243/243	243/243	229/229	250/250	236/236	179/207
25	186/186	240/248	139/139	243/243	241/243	229/245	216/250	244/252	179/181
26	196/206	246/254	131/135	239/243	241/243	221/243	244/244	244/258	179/179
27	198/200	246/248	135/139	243/243	239/243	239/259	244/246	248/254	179/181
28	186/186	242/242	133/133	243/243	243/243	229/245	240/246	252/252	179/179
29	186/186	240/248	131/139	243/245	243/243	229/229	216/250	242/252	179/179
30	176/186	240/248	129/133	243/245	235/245	215/229	216/216	254/256	179/179
31	184/184	250/258	135/137	229/229	229/241	231/231	252/252	248/258	183/183
32	170/170	258/258	129/135	249/249	229/241	233/233	246/246	248/248	183/183
33	170/170	240/256	135/139	229/231	229/229	215/243	246/246	248/248	181/183
34	170/170	250/260	137/139	229/229	229/241	233/235	246/252	248/248	177/183
35	170/186	250/260	135/139	231/233	229/241	231/233	252/252	248/248	183/193
36	192/192	256/258	125/139	247/251	229/229	249/251	232/232	250/260	177/177
37	190/190	256/258	125/139	247/251	229/229	249/251	232/232	250/260	177/177
38	212/212	246/246	131/137	239/243	239/241	245/245	228/244	236/254	179/189
39	194/194	252/256	129/135	245/245	237/237	227/243	220/220	238/238	189/199
40	186/192	240/256	125/129	243/243	237/237	239/243	234/236	234/258	179/187
41	194/194	256/266	129/163	233/235	237/237	217/245	238/240	256/256	189/193
42	194/194	246/256	129/139	235/235	237/245	239/241	238/238	248/254	195/195
43	194/194	246/256	129/129	235/241	235/237	239/239	238/238	240/246	189/195
44	174/174	246/252	129/159	231/235	237/237	235/239	236/238	252/256	185/189
45	184/194	256/256	131/131	235/241	237/237	235/239	236/238	250/250	185/189
46	192/192	252/258	135/167	243/245	233/237	235/239	234/240	246/250	189/189
47	192/192	246/258	135/165	231/243	231/237	235/239	240/240	240/246	189/193
48	184/192	246/252	127/167	235/243	235/235	235/245	238/238	240/246	197/197
49	172/192	252/256	127/155	243/243	237/237	235/239	238/238	240/260	185/189
50	192/192	252/282	133/151	239/243	237/237	217/235	238/238	246/262	193/197
51	172/184	252/256	155/165	235/259	235/237	239/247	240/240	240/248	185/193
52	174/194	256/258	131/163	231/243	235/237	241/243	238/242	246/252	185/185
53	192/192	246/258	129/139	235/243	235/235	217/217	238/240	246/252	185/185
54	174/190	250/264	127/135	239/243	237/243	235/235	236/238	238/262	193/193
55	184/192	240/254	125/131	231/245	235/237	223/245	236/244	232/246	185/187
56	184/204	248/254	125/135	231/243	237/241	223/243	236/244	246/248	181/181
57	184/210	254/258	133/135	233/251	237/237	241/249	238/244	238/246	181/187

Continued next page

Supplemental Table 1. Continued.

Code no.	Microsatellite loci								
	VrZAG62	VrZAG79	VVS2	VVMD5	VVMD7	VVMD28	VVMD32	VVMD25	VVMD27
58	198/204	250/254	133/137	231/241	237/241	243/243	246/246	246/246	179/181
59	190/202	236/248	131/147	233/235	245/245	233/233	268/268	238/246	181/181
60	190/202	236/248	133/147	233/235	245/245	233/233	268/268	238/246	181/181
61	186/190	236/254	133/149	229/233	245/247	233/255	268/268	246/252	181/191
62	184/190	236/252	133/147	229/235	245/247	233/241	268/268	246/252	175/181
63	184/190	236/252	131/133	229/235	245/247	233/241	268/268	246/252	175/181
64	184/186	240/248	131/133	223/229	237/247	231/255	254/268	246/252	181/181
65	184/186	248/254	131/149	223/237	247/251	231/241	252/268	238/252	181/191
66	184/202	248/252	131/149	223/233	245/247	231/265	268/268	238/252	175/181
67	184/184	248/252	123/133	223/233	247/247	241/265	248/268	238/252	175/181
68	184/184	236/252	131/133	233/235	247/247	233/265	268/268	238/252	175/181
69	184/202	244/252	131/147	225/229	247/249	241/265	260/268	238/246	175/191
70	184/202	248/254	131/131	229/233	241/247	231/265	248/268	236/238	181/191
71	184/202	248/254	131/131	233/235	241/247	241/265	268/268	236/238	181/191
72 ^z	198/202	248/248	135/141	235/237	237/261	233/255	246/248	238/252	175/181
73	184/186	244/252	133/139	231/233	247/251	215/265	246/268	238/252	175/177
74 ^z	184/186	248/256	141/143	223/225	237/237	245/255	246/248	238/252	177/181
75 ^z	184/192	242/248	131/141	225/229	237/237	233/255	246/246	240/252	181/197
76 ^z	186/192	242/248	131/131	223/229	237/237	217/227	236/268	238/238	185/187
77	184/192	248/252	127/133	233/233	241/249	239/239	226/226	230/254	177/177
78	178/186	250/262	127/137	231/231	237/239	245/245	242/244	238/264	177/177
79	178/186	250/252	125/135	231/231	237/239	225/243	234/242	238/262	177/177
80	184/186	252/264	125/141	223/233	239/239	225/225	244/244	244/262	177/177
81	184/184	252/252	133/141	233/233	239/239	225/243	234/238	236/236	177/179
82	184/188	238/252	131/139	233/251	229/239	225/243	228/234	244/252	181/193
83	172/178	242/256	139/141	253/261	231/249	225/239	248/248	244/256	185/217
84	192/194	252/254	131/137	249/259	249/249	231/233	234/234	234/234	183/209
85	188/190	252/252	143/159	261/261	249/249	231/235	234/234	236/236	187/203
86	188/196	252/256	139/143	263/263	249/263	213/241	232/234	234/236	203/207
87	186/192	252/256	123/139	261/263	249/249	211/241	232/234	236/236	181/185
88	190/192	240/248	123/145	231/243	239/249	227/269	230/230	248/252	177/187
89	188/188	240/248	121/143	243/243	239/239	227/269	230/230	248/252	177/187
90	188/188	240/248	121/143	243/243	229/229	227/269	230/240	248/252	177/187
91	192/192	236/248	121/143	243/245	241/249	227/251	230/252	244/252	183/187
92	190/192	236/236	121/143	243/243	241/249	237/267	230/230	246/252	177/187
93	194/196	238/240	123/125	231/243	241/249	229/229	230/230	238/250	181/187
94	190/194	238/240	125/145	231/243	239/249	229/243	230/230	238/252	181/187
95	194/196	240/258	125/145	243/243	249/249	229/233	230/230	238/252	177/187
96	192/194	240/258	121/143	243/243	249/249	227/231	230/230	238/254	177/187
97	192/194	240/258	121/143	243/243	249/249	227/231	230/232	238/252	177/187
98	192/194	250/262	125/145	231/251	245/249	227/227	230/230	238/252	177/183
99	194/194	248/248	125/145	243/245	241/249	229/253	230/230	244/252	183/187
100	194/194	236/248	125/145	243/245	241/249	227/251	230/230	244/252	183/187

^z72, 74, 75, and 76 are used for simple sequence repeat standardization for comparison with international databases. Code number is the same reported in Table 1.