

Fig. 1 Relation of root and leaf quality ratings to pH for hydroponically grown iris; 1 (poor) to 5 (excellent).

Root and leaf quality were judged on a scale of 0 (lowest) to 5 (highest). Overall quality for roots was based on root size (root length and total volume of roots) and presence of necrosis. Leaf quality was based on height and presence of chlorosis or necrosis.

Best foliar and root quality was attained at a pH 8.0 and 9.0 for hydroponically-grown plants (Fig. 1). Data from the tissue analysis suggested that higher Ca levels in root tissue tended to reduce necrosis in iris roots; and the subsequent decline in root tissue could directly lower foliage quality (tipburn). The poor plant performance at low pH was not readily explained by leaf or root tissue analyses since no nutrient level standards exist for this crop.

There were no differences between hydroponic treatments for plant height. Onset of foliar chlorosis was delayed significantly and root necrosis was minimized at pH 8.0 and 9.0.

A second greenhouse study was established to more closely duplicate field growing conditions. Iris plants were grown in clay pots containing a Pembroke silt loam (pH-5.1). This study was conducted at a mean temperature of 26°C under natural daylength. The pH levels were decreased by 1 unit using elemental sulfur at a rate of 681g/10 m² of soil surface area and raised with hydrated lime according to Woodruff (8). Treatment levels were 4.5, 5.5, 7.0, 8.5 and 9.5.

Daughter rhizomes of 'Captain Gallant' iris were obtained as in Experiment 1 and similarly surface sterilized. Rhizomes were planted (2/pot) in 15 cm clay containers in their preadjusted soil. Treatments were ar-




Fig. 2. Leaf rating by pH values for irises grown in soils; 1 (poor) to 5 (excellent).

anged in a randomized complete block design with 15 replications. All treatments received 14 N-6-OP-11.6K slow release fertilizer (Osmocote) at a rate of 5 g/pot applied to the media surface following planting.

Container studies using field soil also showed a relation between low pH and foliar tipburn. Although plants at lower pH initially grew more rapidly final plant height was the same. Leaves of most soil grown plants at pH 4.5 and 5.5 had leaf tipburn while those from alkaline soil were free of foliar damage. Leaf quality ratings are shown in Fig. 2.

Information derived from both hydroponic culture and soil studies indicates that the bearded iris grows better in alkaline soils (pH 8-9) than in acid soils. Plant quality and early foliage growth is enhanced by the higher pH.

HortScience 16(5):673-674. 1981.

Effects of Growth Regulators on Diseases of Begonia, Chrysanthemum, Poinsettia, and Rose¹

Bernard Sammons² and Jane F. Rissler

Department of Botany, University of Maryland, College Park, MD 20742

James B. Shanks

Department of Horticulture, University of Maryland, College Park, MD 20742

Additional index words. *Botrytis cinerea*, *Sphaerotheca pannosa*, *Ascochyta chrysanthemi*, *Erysiphe cichoracearum*, *Begonia* × *hiemalis*, *Rosa hybrida*, *Euphorbia pulcherrima*, *Chrysanthemum morifolium*

Abstract. Seven growth regulators representing ethylene-generating materials, auxins, gibberellins, cytokinins and inhibitors were applied prior to inoculation with selected pathogens of chrysanthemum (*Chrysanthemum morifolium* Ramat.), begonia (*Begonia* × *hiemalis* Fotech.), rose (*Rosa hybrida* L.), and poinsettia (*Euphorbia pulcherrima* Willd.). (2-Chloroethyl)phosphonic acid (ethephon) and 6-benzylamino purine (BA) increased the incidence of *Ascochyta* ray blight (incited by *Ascochyta chrysanthemi* Stevens) on 'Ritz' chrysanthemum but the other materials had no effect. None of the growth regulators influenced powdery mildew (*Sphaerotheca pannosa* (Wallr.) Lévl.) of rose, powdery mildew (*Erysiphe cichoracearum* DC.) of begonia, or grey mold (*Botrytis cinerea* Pers. ex Fr.) of poinsettia.

There are various examples of growth regulators increasing (1, 4, 8, 9) or decreasing (2, 4, 5) disease development. The objective of this work was to determine the effects of certain growth-regulators with potential use in greenhouse production on the incidence

and severity of *Ascochyta* ray blight of chrysanthemum, grey mold of poinsettia, and powdery mildew of rose and begonia.

Plants of each species were grown in plastic pots in a root medium of 2 sphagnum peat moss: 1 perlite : 1 vermiculite : 1 sandy loam (by volume) amended with 2.6 g/m³ fritted trace elements, and limed to pH 6.5. Experiments were conducted in the greenhouse at a night temperature of 17°C. Other cultural procedures of fertilization, watering, and insect control were consistent with those in use for the commercial production of the crop (6). The growth regulator treatments made 1 week before inoculation with pathogens were:

1. control — water with 1% polyoxyethylene sorbitan monolaurate (Tween 20) used as a wetting agent in all treatments,

2. 1:100 dilution of a seaweed extract having cytokinin activity (Cytex, Atlantic and Pacific Research, Inc., North Palm Beach, Florida),

¹Received for publication September 20, 1980. Scientific Article No. A2862, Contribution No. 5914, Maryland Agricultural Experiment Station, Departments of Botany and Horticulture, College Park, MD 20742. From a thesis prepared by the senior author in partial fulfillment of the requirements for the MS degree, University of Maryland, 1980. Computer time was supported in full through the Computer Science Center of the University of Maryland. Plants were provided by De Vor Nurseries, Inc., Pleasanton, California; Paul Ecke Poinsettias, Encinitas, California; and Mikkelsen, Inc., Ashtabula, Ohio.

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked advertisement solely to indicate this fact.

²Current address: Department of Plant Pathology and Physiology, Clemson University, Clemson, SC 29631.

Literature Cited

- Burch, J. 1980. Sound advice? Amer. Iris Soc. Bul. 237.
- Gaskill, F. 1964. Soil fertility for irises. Amer. Iris Soc. Bul. 174.
- Lindner, R.C. 1944. Rapid analytical methods for some of the more common inorganic constituents of plant tissues. Plant Physiol. 19:76-89.
- Price, M. 1966. The iris book. D. Van Nostrand, Princeton, N.J.
- Ramsbottom, J.K. 1915. Iris leaf-blotch disease (*Heterosporium gracile* Sacc.). J. Royal Hort. Soc. 40:481-492.
- Shelton, W.R. and A.J. Harper. 1945. Total P determination. Iowa State Col. J. Sci. 15:403.
- Williamson, C. 1980. Soil conditioning for irises. Amer. Iris Soc. Bul. 238.
- Woodruff, C.M. 1946. Determination of exchangeable hydrogen and lime requirements by means of a glass electrode and buffered solution. Soil. Sci. Soc. Amer. Proc. 12:141-142.

3. 50 and 150 ppm BA (Abbott Laboratories, North Chicago, Illinois),

4. 25 ppm BA + 25 ppm GA_{4,7} (Promalin, Abbott Laboratories, North Chicago, Illinois),

5. 150 ppm 6-(benzylamino)-9-(2-tetrahydropyran-9H-purine) (PBA, Shell Development Co., Modesto, California),

6. 25, 50, and 100 ppm α-cyclopropyl-α-(p-methoxyphenyl)-5-pyrimidine methanol (ancymidol, Elanco Products Co., Indianapolis, Indiana),

7. 25 and 50 ppm GA₃ (Abbott Laboratories, North Chicago, Illinois),

8. 5 and 20 ppm naphthaleneacetic acid (NAA, Union Carbide Co., Ambler, Pennsylvania),

9. 500 and 1,000 ppm ethephon (Union Carbide Co., Ambler, Pennsylvania).

Solutions were applied to runoff with a Sure-Shot Sprayer (Milwaukee Sprayer, Milwaukee, Wisconsin).

The design was randomized block with 4 single-plant replicates of each cultivar in each treatment. Each test was repeated at least once.

Plants of 'Fiesta' and 'Ritz' chrysanthemum, propagated in the University of Maryland greenhouses and growing in 12.5 cm pots, were pinched to induce branching and exposed to 8 weeks of 14-hr dark periods to induce flowering. Growth regulators were applied as flowers were opening. Flowers were inoculated by spraying with a conidial suspension of *A. chrysanthemi* (5.0–7.0 × 10⁵ conidia/ml in 0.04% Tween 20) prepared from infected dried plant material. Plants were then placed under intermittent, low pressure mist at 21°C for 48 hr. Ray blight incidence was based upon the estimated percentage of flower area blighted: 1 = 0–5%, 2 = 6–10%, 3 = 11–25%, 4 = 26–50%, and 5 = 51–100% in 2 tests.

Plants of 'Annette Hegg Top Star' poinsettia in 12.5 cm pots were treated with growth regulators after 8 weeks of 14-hr nights. The leaves and developing bracts were inoculated by spraying with a conidial suspension of *B. cinerea* (1.5–1.7 × 10⁶ conidia/ml in 0.04% Tween 20) prepared from potato dextrose agar cultures. Plants were then covered with plastic bags for 48 hr. Grey mold incidence was determined as the percentage of leaves and bracts showing disease symptoms in 2 tests.

Young plants of 'Nixe', 'Improved Schwabenland Red', and 'Improved Schwabenland Pink' begonia were transplanted from 6 cm to 12.5 cm pots. Growth regulator treatments were applied 2 weeks later. Plants were inoculated by spraying with a conidial suspension of *E. cichoracearum*

Table 1. Effect of growth regulators on incidence and severity of Ascochyta ray blight on chrysanthemum 'Ritz' 21 and 30 days after inoculation.

Treatment & rate	Incidence (%) ^w		Severity rating ^x	
	Days after inoculation		Days after inoculation	
	21	30	21	30
Cytex (1:100)	82 ab ^y	78 a ^y	2.2 a ^y	4.3 a ^y
BA (50 ppm)	81 ab	91 abc	2.4 a	4.3 a
(150 ppm)	86 abc	99 b	2.6 a	4.5 a
BA (25 ppm) + GA _{4,7} (25 ppm) ^z	83 ab	88 ab	2.1 a	4.4 a
PBA (150 ppm)	82 ab	88 ab	2.2 a	4.6 a
GA ₃ (25 ppm)	85 abc	82 ab	2.6 a	4.6 a
(50 ppm)	84 abc	82 ab	2.8 a	4.2 a
Ancymidol (25 ppm)	85 abc	80 a	2.2 a	4.2 a
(50 ppm)	73 a	91 ab	2.1 a	4.1 a
(100 ppm)	73 a	93 ab	2.3 a	4.1 a
NAA (5 ppm)	76 a	83 ab	2.3 a	4.1 a
(20 ppm)	80 ab	90 ab	2.9 a	4.1 a
Ethephon (500 ppm)	99 c	94 ab	4.5 b	4.9 a
(1,000 ppm)	97 bc	98 b	4.8 b	5.0 a
Control H ₂ O + Tween 20	84 abc	80 a	2.7 a	4.3 a

^wIncidence = percentage of flowers showing symptoms.

^xSeverity scale: 1 = 0–5%, 2 = 6–10%, 3 = 11–25%, 4 = 26–50%, 5 = 51–100% flower area blighted.

^yMean separation within columns by Student-Newman-Keuls test, 5% level. Each value is the mean of 4 replicates.

^zPromalin.

(0.8–7.0 × 10⁵ conidia/ml in 0.04% Tween 20) prepared from infected begonias and were then placed under low pressure, intermittent mist at 21°C for 24 hr. Mildew incidence was determined as the percentage of leaves showing symptoms in 2 tests. Severity was rated visually at the same time periods as the percentage of leaf area covered by conidia and conidiophores: 1 = 0–2%, 2 = 3–5%, 3 = 6–10%, 4 = 11–25%, 5 = 26–50%, and 6 = 51–100%.

Dormant, bare root, xx grade plants 'Forever Yours' rose were received from the nursery and grown in 30 cm pots for 4 weeks before growth regulator treatments were applied. Plants were inoculated by spraying with a conidial suspension of *S. pannosa* (5.0–7.0 × 10⁴ conidia/ml in Tween 20 at 0.04%) prepared from infected rose leaves and then subjected to low pressure, intermittent mist at 21°C for 24 hr. Mildew incidence and severity were estimated as for begonia in 2 tests.

Incidence of Ascochyta ray blight on 'Ritz' chrysanthemums was significantly different from the control with ethephon at 1000 ppm but not with 500 ppm 30 days after inoculation. At both rates, severity was significantly greater at 21 days after inoculation. Incidence was significantly greater at 30 days with BA at 150 ppm (Table 1). Except for ethephon and BA, the regulators did not significantly change the incidence or severity of the diseases studied. Ray blight was significantly influenced by ethephon and BA in both tests.

The results with ethephon are consistent with the concept that ethylene contributes to symptom development in many plant diseases

(7). The lack of observable change in disease development in most of the plant-pathogen-chemical interactions could indicate the failure of the chemicals to affect the processes determining disease development or could be due to ineffectual concentrations of chemicals, inappropriate environmental conditions, or non-optimal host developmental stages.

Literature Cited

1. Buchenauer, H. and D. C. Erwin. 1976. Effect of the plant growth retardant Pydanon on Verticillium wilt of cotton and tomato. *Phytopathology* 66:1140–1143.
2. Crossan, D. F. and D. J. Fieldhouse. 1964. A comparison of dwarfing and other compounds with and without fixed copper fungicide, for control of bacterial spot of pepper. *Plant Dis. Rptr.* 48:549–550.
3. Davis, D. and A. E. Dimond. 1953. Inducing disease resistance with plant growth-regulators. *Phytopathology* 43:137–140.
4. Dropkin, V. H., J. P. Helgeson, and C. D. Upper. 1968. The hypersensitivity reaction of tomatoes resistant to *Meloidogyne incognita*: reversal by cytokinins. *J. Nematol.* 1:55–61.
5. Fliegel, P., K. G. Parker, and L. J. Edgerton. 1966. Gibberellic acid treatment of sour cherry infected by sour cherry yellows virus: response to sprays applied throughout the growing season and the influence of environmental conditions. *Plant Dis. Rptr.* 50:240–242.
6. Larson, R. A. (ed.). 1980. *Introduction to floriculture*. Academic Press, New York.
7. Primrose, S. B. 1977. Ethylene and agriculture: the role of the microbe. *J. Appl. Bacteriol.* 46:1–25.
8. Sinha, A. K. and R. K. S. Wood. 1967. The effect of growth substances on Verticillium wilt of tomato plants. *Ann. Appl. Biol.* 59:117–128.
9. Tahori, A. S., G. Zeidler, and A. H. Halevy. 1965. Effect of some plant growth-retarding compounds on three fungal diseases and one viral disease. *Plant Dis. Rptr.* 49:775–777.