

CORRIGENDA

• In the article “‘Aylés’, ‘Guara’, and ‘Moncayo’ Almonds”, by A. Felipe and R. Socias i Company (*HortScience* 22(5):961–962, October 1987), the percentage bar did not appear on Fig. 1. The correct figure appears on the right.

• In the article “Efficacy of Ancymidol, Daminozide, Flurprimidol, Paclobutrazol, and XE-1019 when Followed by Irrigation”, by James E. Barrett, Carolyn A. Bartuska, and Terril A. Nell (*HortScience* 22(6):1287–1289, December 1987), Table 1 was printed incorrectly. The corrected table appears below.




Table 1. Stem elongation for chrysanthemums given overhead irrigation at intervals after growth retardant spray.^z

Irrigation time ^y (hr)	Stem elongation (cm)				
	Paclobutrazol (100 mg·liter ⁻¹)	XE-1019 (50 mg·liter ⁻¹)	Chemical Ancymidol (100 mg·liter ⁻¹)	Daminozide (5000 mg·liter ⁻¹)	Flurprimidol (75 mg·liter ⁻¹)
0.5	11	9	4	17	16
1	12	9	4	10	15
2	11	8	4	7	15
4	14	8	4	6	13
8	14	8	3	4	13
24	12	8	3	5	15
None	10	8	4	5	14
Significance:					
HSD (5%)	5		2		NS
Chemical (C)	**		**		---
Irrigation (I)	NS		**		---
C × I	NS		**		---
Nontreated plants	38		21		42

^zData are from three separate experiments. Plants were single-stem ‘Nob Hill’ in 12.5-cm pots.

^yOverhead irrigation (1.5 cm) was applied at indicated time after spray treatment or no irrigation was applied.

***Significant at $P = 0.05$ and 0.01 , respectively.

• In the article “Two Methods of Studying Rest: Temperature Alteration and Genetic Analysis”, by Frank G. Dennis, Jr. (*HortScience* 22(5):820–823, October 1987), Table 1 was printed incorrectly. The corrected table appears below.

Table 1. Effects of alternating^x vs. constant temperatures (°C) on breaking of rest.

Species and cultivar	Effect	Ref.
Peach leaf buds		
Redhaven	6°/15° > > 4°	10
Redskin	6°/15° > 4°	10
Redhaven	4°/15° > > 4°	5
	4°/15° = 4°	5, 9
	6°/15° = 6°	9
	6°/9°, 6°/11°, 6°/13° and 6°/15° > 6°	9
Cornet	0°/15°, 4°/15°, and 6°/15° > > 0°, 4°, or 8°	9
Peach flower buds		
Redhaven	4°/15° = 4°	5
	4°/15° > 4°	9
	6°/9°, 6°/11°, 6°/13° and 6°/15° ≥ 6°	9
Sungold ^x	7°/15° > 7° ^y	15
Sour cherry flower buds		
Montmorency	5°/15° > 5° ^y	13
Peach seeds		
Halford	5°/10° > 5°	1
	5°/15° < 5°	1
Siberian C.	5°/10° > 5° or 10°	(A. Mahhou, unpublished data)
	5°/15° < 5°	

^xAll 18 hr at low and 6 hr at high temperature except for ‘Sungold’ nectarine (14/10 hr).

^yEffective only during latter part of chilling period.

^zNectarine.