

Calcium, Iron, Potassium, Phosphorus, and Vitamin C Content of Organic and Hydroponic Tomatoes

Zdenka Premuzic, Martha Bargiela, Ana Garcia, Alicia Rendina, and Alicia Iorio

Department of Chemistry, Faculty of Agriculture, University of Buenos Aires, Argentina

Additional index words. *Lycopersicon esculentum*, vermicompost, fruit quality, nutrition

Abstract. Two tomato (*Lycopersicon esculentum* Mill.) cultivars were grown in two organic and two inorganic media to evaluate the effects upon the levels of Ca, Fe, K, P, and vitamin C in the fruit. 'Platense' tomato was grown in a glasshouse, on sand or peat-perlite (hydroponic substrates) irrigated with a complete solution of macro and microelements, or on 100% vermicompost or 50% vermicompost-50% soil (organic substrates) irrigated with water. Fruit were harvested at physiological maturity, and levels of P, K, Ca, Fe, and vitamin C were determined. Fruit grown on organic substrates contained significantly more Ca and vitamin C and less Fe than did fruit grown on hydroponic media. Phosphorus and K content did not differ between fruit from organic and hydroponic substrates.

The intensive nature of vegetable production necessitates fertilizer management aimed at maximizing yield and quality (Huett and Rose, 1988). There is great interest in organic crop production (Montagu and Goh, 1990), particularly in adding organic matter to the soil, which liberates nutrients in a gradual and controlled way, allowing a greater production of vegetables with a minor environmental impact (Premuzic and Iorio, 1996; Premuzic and Nicolossi, 1995). Organic wastes should be applied to soil after being processed to obtain a mature and stabilized compost (Senesi, 1989). Earthworms act on organic wastes to form vermicompost, which releases nutrients slowly (Brandjes, 1984; Ferruzi, 1994).

To compare substrate effect on the levels of P, K, Ca, Fe, and vitamin C in the fruit, tomatoes were grown on two hydroponic substrates irrigated with a nutrient solution (Premuzic et al., 1996) and on two organic (vermicompost) substrates irrigated with water.

Materials and Methods

'Platense' tomatoes were grown in a naturally lighted glasshouse with average maximum-minimum temperatures of 24-16 °C, using 3-L black opaque polyethylene pots. Seed was directly sown in pots filled with one of four substrates. Two substrates were hydroponic: purified sand (S) (Hewitt, 1966) and peat-perlite (P-P), both irrigated with a solution complete in macro and microelements (Table 1) (Premuzic et al., 1996), taking into account the nutrient requirements of tomato in an open hydroponic system (Cheng and Dube,

1976; Cooper, 1976; Hoagland and Arnon, 1950; Martinez-Caldevilla and Garcia-Lozano, 1993; Pardossi et al., 1987; Sonneveld, 1985). The other substrates were organic: 100% vermicompost (V100) and 50% vermicompost : 50% soil (V50) (Table 2), irrigated with water. All four substrates were irrigated daily, early in the morning and late in the afternoon. The experimental design was completely randomized with three replications and four treatments (substrates). Fruit were harvested at physiological maturity and dried for 48 h at 70 °C for determination of P by molecular absorption spectrophotometry (Iorio et al., 1992), K by flame photometry, and Ca and Fe by atomic absorption. Vitamin C (Official Methods of Analysis of the Association of Analytical Chemists, 1980) content was measured on fruits frozen at -18 °C for 60 d. The data were analyzed by analysis of variance, and when F values were significant, by comparison of means with Tukey's test (Tukey, 1950).

Results and Discussion

Phosphorus and K levels of fruit did not differ significantly among the four substrates. The mean levels of P (0.083%) were below, and the mean levels of K (3.31%) were similar

to those previously reported (Auclair et al., 1995; Magalhaes and Wilcox, 1983a, 1983b; Sobulo et al., 1975).

Calcium concentration (Fig. 1A) was significantly higher for fruit grown in the organic than in the hydroponic substrates, and the levels were higher than previously reported (Agui et al., 1979; Maher, 1976; McMurtry et al., 1993; Muller-Haslach et al., 1986). There were no significant differences in Ca content between fruit from V50 and V100 or between fruit from S and P-P. Iron concentration (Fig. 1B) was significantly higher in the fruit grown hydroponically, but did not differ between V50 and V100 or between S and P-P.

Vitamin C concentration (Fig. 1C) was higher in fruit grown on vermicompost than in fruit grown on the hydroponic substrates, but did not vary significantly between fruit from V50 and V100 or between fruit from S and P-P. Fruit with greater vitamin C concentration have greater nutritional value (Bangerth, 1976; Gull et al., 1978; Lopez-Andreu et al., 1988; Silva and Scotti, 1973). Vitamin C levels were similar to those previously reported (Adams et al., 1978; Ketsa and Wongveerakhan, 1987; Klein and Perry, 1982; Muller-Haslach et al., 1986), although a little lower, as they were measured in frozen fruit (Barowski et al., 1981).

The higher concentration of Ca and vitamin C in the organic tomatoes suggests a positive correlation of Ca and vitamin C levels, in agreement with Bangerth (1976). According to the values presented in the USDA Food Agricultural Handbook (1963), the fruits grown organically had an adequate level of nutrients.

The quality of food and its composition are being scrutinized more critically than ever before by the consumer (Goh and Vitykon, 1986). The nutritional value of tomato consists essentially of carotene (provitamin A), lycopene, and vitamin C (Lopez-Andreu et al., 1988; Muller-Haslach et al., 1986). Most research indicates that fruit quality is affected not only by adequate amounts of N, P, and K, but also by the balance between the levels of these nutrients (Varis and George, 1985). The relationship between fertilizer application and vitamin C content in fruit has been discussed by various authors. Sharma and Mann (1971) noted a positive correlation between N and P fertilization and level of vitamin C. Mengel (1972) reported that higher rates of N increase

Table 1. Composition of the nutrient solution employed for irrigation of the hydroponic substrates. CaCO₃ (0.42g·L⁻¹) and MgCO₃ (0.17g·L⁻¹) were added directly to the hydroponic solution.

Nutrient	Stock solution (g·L ⁻¹)	Stock solution used in hydroponic solution (mL·L ⁻¹)
NH ₄ NO ₃	21.50	21
NaH ₂ PO ₄	63.00	4
KCl	19.00	33
Na ₂ SO ₄	50.00	13
FeSO ₄	26.20	5
MnSO ₄	0.80	12
H ₃ BO ₃	0.34	2
CuSO ₄	0.63	1
Na ₂ MoO ₄	0.15	1
ZnSO ₄	0.40	1

Received for publication 27 May 1997. Accepted for publication 10 Dec. 1997. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

levels of amino acids and vitamin C at the expense of carbohydrate. A negative correlation between fertility and vitamin C was cited by both Montagu and Goh (1990) and Muller-Haslach et al. (1986). High rates of fertilizers promote foliar growth and shading of the developing fruit, especially in glasshouse-grown tomatoes. This would affect vitamin C, as its synthesis depends on exposure to light (Venter, 1977).

In our experiment, tomatoes grown with vermicompost as the nutrient supply had a higher concentration of vitamin C, which could be related to the shading effect, as foliage development was greater on the hydroponic substrates (Premuzic and Iorio, 1996). Higher Ca levels in organically grown tomatoes were associated with reduced levels of other cations such as Na, Mg, and K, which are usually more abundant in substrates irrigated with nutrient solutions (Matev and Stanchev, 1979). An-

other important factor is that the fruit grown on organic substrate received nutrients in a slow-release fashion, optimizing the nutrient availability during fruit growth, when the uptake of nutrients is more important (Huett and Dettman, 1988).

Composting is an environmentally safe means of converting a waste product into an acceptable source of organic matter (Sterrett et al., 1983). Compost prepared from biogenic waste is an excellent material for use in intensive vegetable production, as it improves yield and nutritional and storage quality of the product (Vogtmann et al., 1993).

The results of our experiment and of others may lead to increased organic production of high-quality, contaminant-free food and contribute to the use of biogenic wastes in crop production, while reducing environmental impact.

Literature Cited

Adams, P., J.N. Davies, and G.W. Winsor. 1978. Tomato yield in relation to the N, K and Mg status of the plants and of the peat substrate. *Plant and Soil* 49:137-148.
 Agui, Y., M. Gomez, and L. Recalde. 1979. Interactions between K, Ca and Mg on the uptake level and distribution in tomato plants. *Anales Edafologia y Agrobiologia* 38:267-281.
 Association of Analytical Chemists. 1980. Official methods of analysis of the Association of Analytical Chemists. Washington, D.C.
 Auclair, L., J.A. Zee, and A. Karam. 1995. Nutritive value, organoleptic quality and productivity of greenhouse tomatoes in relation to production

method: Organic-conventional-hydroponic. *Sciences des Aliments* 15:511-528.
 Bangerth, F. 1976. Relationship between the Ca content and Ca treatment and the ascorbic acid content of apple, pear and tomato fruits. *Qualitas Plantarum* 26:341-348.
 Barowski, J.K. Elner, and H. Michalik. 1981. Freezing of tomatoes in liquid N. *Biuletyn Warzywnicy* 25:297-310.
 Brandjes, P. 1984. Worm compost, a good organic manure. *Vakblad voor de Bloemisterij* 39:3-37.
 Cheng, B.T and A. Dube. 1976. A comparison of various artificial substrates for hydroponic culture in relation to nutrient solutions. *Agriculture-Quebec* 33:29-31.
 Cooper, A.J. 1976. Crop production with NFT. *Inter. Congres Soilless Culture*, Las Palmas, Canarias, Spain.
 Ferruzi, C. 1994. *Manual de lombricultura*. Mundiprensa. Madrid, Spain.
 Goh, K.M. and P. Vytakoni. 1986. Effects of fertilizers on vegetable production. II. Effects of nitrogen fertilizers on nitrogen content and nitrate accumulation of spinach and beetroot. *N.Z. J. Agr. Res.* 29:485-494.
 Gull, D.D, S.J. Locascio, and S.R. Kostewicz. 1978. Composition of greenhouse tomatoes as affected by cultivar, production media and fertilizer. *Proc. Fla. State Hort. Soc.* 90:395-397.
 Hewitt, E.J. 1966. Sand and water culture methods used in study of plant nutrition. *Commonwealth Agr. Bureaux*.
 Hoagland D.R. and D.I. Arnon. 1950. The water culture method for growing plants without soil. *California Agr. Expt. Sta. Circ.* 347.
 Huett, D.O. and B.E. Dettmann. 1988. Effect of nitrogen on growth, fruit quality and nutrient uptake of tomatoes grown in sand culture. *Austral. J. Expt. Agr.* 28:391-399.
 Huett, D.O. and G. Rose. 1988. Diagnostic nitrogen correlations for tomatoes grown in sand culture. *Austral. J. Expt. Agr.* 28:401-409.
 Iorio, A., A. Rendina, A. Garcia, and A. Schor. 1992. Efecto de la temperatura, matriz vegetal y tiempo sobre la reduccion del acido fosfomolibdico. *Rev. Fac. Agron.* 13:261-265.
 Ketsa, S. and Wongveerakhan. 1987. The ascorbic content at maturity stages in tomato (*Lycopersicon esculentum*) cultivars. *Thai J. Agr. Sci.* 20:257-261.
 Klein, B.P. and A.K. Perry. 1982. Ascorbic acid and vitamin A activity in selected vegetables from different geographical areas of the United States. *J. Food Sci.* 47:941-945.
 Lopez-Andreu, F.J., R.M. Esteban, E. Molla, and O. Carpena. 1988. Incidence of nutrition system on the quality of tomato fruits. I. Yields, physic parameters, nitrogen fractions, acidity and sugars. *Anal. Edafologia y Agrobiologia*. XVII:1181-1190.
 Magalhaes, J.R. and G.E. Wilcox. 1983a. Tomato growth and nutrient uptake patterns as influenced by nitrogen and light intensity. *J. Plant Nutr.* 6:941-956.
 Magalhaes, J.R. and G.E. Wilcox. 1983b. Tomato growth and mineral composition as influenced by nitrogen form and light intensity. *J. Plant Nutr.* 6:847-862.
 Maher, M.J. 1976. Growth and nutrient content of glasshouse tomato crop grown in peat. *Scientia Hort.* 4:23-26.
 Martinez-Caldevilla, E. and M. Garcia-Lozano. 1993. *Cultivos de Hortalizas sin suelo en clima Mediterraneo*. p. 41-60.
 Matev, Y. and L. Stanchev. 1979. Effect of Na⁺, K⁺, Ca²⁺ and Mg²⁺ imbalance on the development and biological value of glasshouse tomatoes. *Gradinska Lozarska Nauka* 16:76-82.

Table 2. Chemical properties of the soil and the organic substrate

Properties	Soil	Vermicompost
Total N (%)	0.12	1.30
pH	5.80	7.00
OM (%)	2.80	13.20
Ca (g·kg ⁻¹)	0.36	0.46
Mg (g·kg ⁻¹)	0.34	0.60
K (g·kg ⁻¹)	0.54	1.75
Na (g·kg ⁻¹)	0.13	0.11
Fe (mg·kg ⁻¹)	----	43.00
P (mg·kg ⁻¹)	5.00	350.00

Method used for exchangeable cations: NH₄CH₃COO; pH = 7.




Fig. 1. Effects of media on concentrations of Ca, Fe, and Vitamin C in tomato fruit. Media used were 100% vermicompost (V100), 50% vermicompost-50% soil (V50), sand (S), and peat-perlite (P-P). Three replications per treatment with mean separation by Tukey's test (P ≤ 0.05).

- McMurtry, M.R., D.C. Sanders, P.V. Nelson, and A. Nash. 1993. Mineral nutrient concentration and uptake by tomato irrigated with recirculating aquaculture water as influenced by quantity of fish waste products supplied. *J. Plant Nutr.* 16:407-419.
- Mengel, K. 1972. Ernährung und Stoffwechsel der Pflanze. Fischer, Stuttgart.
- Montagu, K.D. and K.M. Goh. 1990. Effects of forms and rates of organic and inorganic nitrogen fertilizers on the yield and some quality indices of tomatoes (*Lycopersicon esculentum* Mill.). *N.Z. J. Crop Hort. Sci.* 18:31-37.
- Muller-Haslach, V.W., G. Arold, and V. Kimmel. 1986. Effect of nutrient intensity on the quality of tomatoes. *Bayerisches Landwirtschaftliches Jahrb. Sonderheft* 63:81-104.
- Pardossi, A., F. Tognoni, and G. Bertero. 1987. The influence of nutrient standard concentration on growth, mineral uptake and yield of tomato plants grown in NFT. *Adv. Hort. Sci.* 1:55-60.
- Premuzic, Z., G. Bilotti, A. de los Rios, and A. Iorio. 1996. El aprendizaje del tema de soluciones en química: Una alternativa hidropónica. *CIT: Rev. Inform. Tecnológica, Serena, Chile* 7:91-94.
- Premuzic, Z. and A. Iorio. 1995. Comportamiento de una variedad de tomate frente a dos sistemas de fertilización: Orgánico e inorgánico. *Revista Frutihortícola.* Nov. p. 10.
- Premuzic, Z. and A. Iorio. 1996. La producción y el contenido de nitrógeno en tomates orgánicos respecto de hidropónicos. *Actas XII Congreso Latinoamericano de Química, Concepcion, Chile.* p. 451.
- Premuzic, Z. and R. Nicolossi. 1995. Contenidos de N, P, K en plántulas de tomate con un sustrato orgánico. *Actas XXXV Congr. Brasileiro de Química, Salvador, Bahia, Brazil.*
- Pulupol, L.U., M.H. Behboudian, and K.J. Fisher. 1996. Growth, yield, and postharvest attributes of glasshouse tomatoes produced under deficit irrigation. *HortScience* 31:926-929.
- Senesi, N. 1989. Composted materials as organic fertilizers. *Sci. Total Environ.* 2:521-542.
- Sharma, C.B. and H.S. Mann. 1971. Effect of phosphatic fertilizers at varying levels of nitrogen and phosphate on the quality of tomato fruits. *Indian J. Hort.* 28:228-233.
- Silva, S. and Y. Scotti. 1973. Influenza della concimazione oligominerale su alcune caratteristiche nutritive del pomodoro. *Annu. Fac. Agraria XIII*:240-248.
- Sobulo, R.A., A. Fayemi, and A. Agboola. 1975. Nutrient requirements of tomatoes (*Lycopersicon esculentum*) in southwestern Nigeria. II. Foliar analysis for assessing N, P and K requirements. *Expt. Agr.* 11:137-143.
- Sonnenveld, C. 1985. A method for calculating the composition of nutrient solutions for soilless cultures. Second translated edition N 10 Glasshouse crops Res. and Expt. Sta., Noaldwijk, Netherlands.
- Sterrett, S.B., C.W. Reynolds, F.D. Schales, R.L. Chaney, and L.W. Douglass. 1983. Transplant quality, yield and heavy metal accumulation of tomato, muskmelon and cabbage grown in media containing sewage sludge compost. *J. Amer. Soc. Hort. Sci.* 108:36-41.
- Tukey, J.W. and G. Snedecor. 1950. Statistical methods. Iowa State College Press, Ames.
- U.S. Department of Agriculture. Food Agriculture Handbook. 1963. N 8.
- Varis, S. and R.A.T. George. 1985. The influence of mineral nutrition on fruit yield and quality in tomato. *J. Hort Sci.* 60:373-376.
- Venter, F. 1977. Solar radiation and vitamin C content of tomato fruit. *Acta Hort.* 58:121-127.
- Vogtmann, H., K. Matthies, B. Kehres, and A. Meier-Ploeger. 1993. Enhanced food quality: Effects of composts on the quality of foods. *Compost Sci. and Util.* 1:82-100.