

Response of Landscape Groundcovers to Deficit Irrigation: An Assessment Based on Normalized Difference Vegetation Index and Visual Quality Rating

Anish Sapkota and Amir Haghverdi

Department of Environmental Sciences, University of California–Riverside, Riverside, CA 92521, USA

Donald Merhaut

Department of Botany and Plant Sciences, University of California–Riverside, Riverside, CA 92521, USA

Amninder Singh and Jean Claude Iradukunda

Department of Environmental Sciences, University of California–Riverside, Riverside, CA 92521, USA

Keywords. evapotranspiration, groundcover water response function, ornamental plants, urban irrigation, water conservation

Abstract. Developing water conservation strategies for urban landscape groundcovers grown in hot and dry summers like inland Southern California, USA, is crucial because they are one of the largest residential water users. A 2-year (2020–21) study was conducted in Riverside, CA, to assess the effect of irrigation rates on the growth of landscape groundcovers as evaluated by visual quality ratings (VR) and normalized difference vegetation index (NDVI). Relationships between VR and NDVI were also established to obtain the minimum threshold values of NDVI for each groundcover. Lastly, the groundcover water response function was developed to estimate groundcover response to irrigation rates over time. Four reference evapotranspiration (ET_0)-based irrigation treatments ranging from 24% to 99% ET_0 and 10 landscape groundcovers were laid in a randomized complete block design and replicated three times. Data were collected from May to October in 2020 and 2021. The irrigation controller overirrigated the plots on average by 7.7% and 4.7% in 2020 and 2021, respectively. A significant relationship ($P < 0.05$, $0.35 \leq R^2 \leq 0.82$) between NDVI and VR for each landscape groundcovers was found. On the basis of the NDVI values and VR, it was found that three landscape groundcovers, including *Rhadowia spinescens*, *Baccharis* × ‘Starn’ Thompson, *Eriogonum fasciculatum* ‘Warriner Lytle’ can withstand water stress and can maintain their growth and visual quality at 24% ET_0 irrigation. Groundcovers *Ruschia lineolata* nana, *Rosmarinus officinalis* ‘Roman Beauty’, and *Eremphila glabra* showed the potential to perform well with as low as 49% ET_0 irrigation, whereas *Lantana montevidensis*, *Oenothera stubbei*, and *Lonicera japonica* required 75% ET_0 or more.

Water-efficient horticultural alternatives to turfgrass are recommended in many urban areas where water is scarce. Predominant

among these recommended alternatives are perennial groundcovers, which are low-growing plants that form a continuous soil covering (Davison 1999). They vary significantly in shape, size, texture, and color, with heights typically ranging from 7.5 cm to 1 m (Davison 1999; Pittenger et al. 2001). However, the assumption that groundcovers consume less water (Pittenger et al. 2001) is mainly anecdotal because few studies have assessed the response of groundcovers to deficit irrigation (Costello and Jones 2014; Garcia-Navarro et al. 2004; Nazemi Rafi et al. 2019; Pittenger et al. 2001). Moreover, the quality assessment of groundcovers is mostly based on the visual ratings (Nazemi Rafi et al. 2019; Pittenger et al. 2001), and the potential use of quantitative quality predictor, including normalized difference vegetation index (NDVI) has not been studied for landscape groundcover.

Visual quality rating (VR) is a dominant and traditional method to assign a numerical value to plant appearance. Nevertheless, estimating groundcover quality based on visual methods can be time-consuming and requires a skilled and trained evaluator (Wang et al. 2022). Even a well-trained person may introduce bias because the VR process is subjective and prone to rater’s fatigue (Horst et al. 1984; Lusier et al. 2006; Wang et al. 2022). The relationship between VR and the NDVI, widely used indicator of vegetative health (Easterday et al. 2019; Haghverdi et al. 2021c), has been studied in turfgrass plots (Fitz-Rodríguez and Choi 2002; Haghverdi et al. 2021b, 2021c; Leinauer et al. 2014). In turfgrass research, some studies suggested the use of NDVI as an alternative to visual rating because it provides consistent and reliable evaluation of turfgrass quality in less time compared with visual quality (Bell et al. 2009; Fitz-Rodríguez and Choi 2002; Haghverdi et al. 2021c). In contrast, some studies in turfgrass highlighted the practical limitations of using NDVI for the quality assessment (Bremer et al. 2011; Leinauer et al. 2014). However, studies evaluating the potential of NDVI values to assess the quality of landscape groundcovers are yet to be done.

Haghverdi et al. (2021c) introduced the turfgrass water response function as an empirical regression-based model to estimate the response of turfgrass to extreme drought and limited irrigation scenarios. These models were developed using data from turfgrass fields in southern and central California along with long-term weather data obtained from nearby weather stations (Haghverdi et al. 2021b, 2021c). Development of these models can be helpful in irrigation management as they estimate the quality (based on NDVI values) of plants for different rates of irrigation. Because the water requirements of landscape groundcovers can significantly differ among species, development of a species-specific water response function model [hereafter called groundcover water response function (GCWRF)] can help predict the quality of groundcovers based on NDVI values at varying rates of irrigation. This in turn can be helpful in optimizing irrigation rates while maintaining the quality of groundcovers.

Evapotranspiration (ET)-based smart irrigation controllers with on-site weather measurements can be used for autonomous landscape irrigation management. These controllers have been reported to reduce irrigation water by 40% to 61% in plot studies and 28% to 32% in residential studies (Dukes 2020). However, few studies evaluated their performance for landscape groundcovers (Shober et al. 2009; US Bureau of Reclamation 2008), particularly in arid and semiarid regions such as inland Southern California, where keeping plants alive with minimum water application is often required (Haghverdi et al. 2021c; Serena et al. 2020). Efficient irrigation scheduling using ET-based controllers depends on the

Received for publication 28 Sep 2022. Accepted for publication 27 Dec 2022.

Published online 27 Jan 2023.

This research was funded in part by CANERS Foundation and the Metropolitan Water District of Southern California (#180897). We acknowledge the help of Janet Hartin, Margaret O’Neil, and Agricultural Operations at the University of California Riverside for their help with the project. We also acknowledge the detailed and constructive comments from three anonymous reviewers. A.S. and A.H. are the corresponding authors. E-mail: asapk001@ucr.edu (A.S.) or amirh@ucr.edu (A.H.).

This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

availability of science-based plant factor information for each landscape species and the accuracy of the ET_o estimations by the irrigation controller (Haghighi et al. 2021a). Hence, the objectives of this study were to 1) evaluate the effect of irrigation rates on VR and NDVI of 10 landscape groundcovers; 2) examine the strength of linear relationships between VR and NDVI for each groundcover species; 3) estimate the response of groundcover species to multiple irrigation regimes under extremely low, high, and mean atmospheric evaporative demand using GCWRFs; and 4) determine the reliability of the Weathermatic smart ET-based controller for autonomous irrigation scheduling.

Materials and Methods

Study area. A 2-year (2020–21) study was conducted in a year-old established field at the Agricultural Experimental Station, Riverside (lat. 33°58' N, long. 117°19' W, 307 m. elevation) at the University of California–Riverside, Riverside, CA. The experimental field had a soil classified as Hanford coarse sandy loam (websoilsurvey.sc.egov.usda.gov). The climate in Riverside is semiarid. The ET_o demand during the experimental season (May–October) was higher than the long-term average in both years, while precipitation was negligible (Table 1). Fertilizer 15–5–8 microgreen (Simplot Turf & Horticulture, San Diego, CA) was top-dressed at 49 kg-ha⁻¹ nitrogen (N) and the soil was treated with preemergent herbicide in 2019 to control weeds. The experimental plots were hand-weeded during the study, and alleys were sprayed with herbicides. Fast-growing groundcovers were pruned to maintain 38 cm height. Similarly, the lateral growth of the groundcover was always confined within the designated plot size (3.05 m × 3.05 m) by trimming the excess growth.

Experimental design, groundcover species selection, and irrigation application. Ten woody, herbaceous, and succulent landscape groundcovers, including some native and widely grown species in California with different growth habits and water requirements, were planted in 2019 (Fig. 1). Two other plant species (*Delosperma cooperi* ‘John Proffitt’ and *Frankenia thymifolia*) did not grow well and were not included in this study. *Eriogonum* did not become fully established in the first year after planting, so results for it are presented only for 2021. For six species (i.e., *Rhagodia*, *Eriogonum*, *Baccharis*, *Eremphila*, *Ruschia*, and

Oenothera), 12 to 16 plants per plot were acquired in 2.8-L (1 gallon) containers. For the remaining species, 10-cm pot plants in full trays were obtained and planted at a higher density to ensure proper plot coverage and plant establishment. Four irrigation treatments (80%, 60%, 40%, and 20% ET_o) replicated three times were laid in two adjacent randomized complete block designs totaling 144 individual experimental plots. Each plot was ~3 m × 3 m, with a 1.2-m alley between the neighboring plots. Four 300-mm tall quarter-circle pop-up heads (Toro 570Z series; The Toro Company, Bloomington, MN, USA) with pressure-compensating precision series spray nozzles (Model 0-10-Q, The TORO Company) were used to irrigate each plot. Each plot was independently controlled using a Hunter PGV-101G solenoid valve (Hunter Industries, Inc., San Marcos, CA). In addition, a pressure regulator was installed in the field to maintain steady water pressure.

The automatic irrigation scheduling was done by a Weathermatic SmartLine SL4800 smart irrigation controller (Weathermatic, Garland, TX, USA). The controller works in the principle of the Hargreaves and Samani (1985) equation, which uses on-site temperature data and latitude-based solar radiation to estimate ET_o . The SmartLine irrigation controller was connected to an SLW1 weather sensor and a Badger Meter Recordall Turbo flowmeter (Badger Meter, Inc., Milwaukee, WI, USA). Flowmeter was calibrated using field-based flow test data. The low half distribution uniformity of the system (86%) was determined at the beginning of the experiment using a catch-cans test. The irrigation controller was programmed to apply the desired ET_o level divided by the irrigation frequency for each treatment. Therefore, programmed irrigation rates were 93%, 70%, 47%, and 23% ET_o (Table 2). The controller initiates the irrigation whenever the minimum deficit irrigation threshold is reached, can irrigate multiple times a day until the desired level is reached; however, it does not irrigate outside of the pre-defined irrigation window. Plots were irrigated between midnight to 8 AM to avoid evaporative water loss. The smart controller automatically performed run-soak cycles to eliminate runoff. The maximum runtime and minimum soak time between the irrigation event were set to be 10 and 30 min, respectively. The irrigation trial ran from early May to late October in 2020 and 2021, and uniform nonlimiting (80% ET_o) irrigation was applied from November to April.

The performance of the ET-based irrigation controller was evaluated using CIMIS- ET_o rates obtained from the nearby California Irrigation Management Information System (CIMIS #44). The irrigation runtime data for each treatment were retrieved from the controller, converted to CIMIS- ET_o , and compared with the programmed ET_o values at the beginning of the trial.

Data collection. The effect of irrigation rates on the landscape groundcovers was evaluated by measuring the NDVI and VR. The NDVI is a widely used index for vegetation assessment (Huang et al. 2021) as it correlates strongly to green coverage, above-ground biomass, and plant vigor (Easterday et al. 2019; Garg et al. 2022).

The NDVI data were collected using handheld GreenSeeker (Trimble Inc., Sunnyvale, CA) close to solar noon on cloud-free days. The GreenSeeker was held at waist height and hovered over the plot (~3 m²) in an inverse Z-shape keeping the trigger engaged to get a representative and average NDVI value from each experimental plot. Data were collected during solar noon in a cloud free day. In both years, the NDVI data were collected on 12 dates during the experimental season (May to October).

Canopy pictures from each experimental plot were captured on the same day of NDVI data collection using a 12-megapixel Olympus digital camera (TG-5; Olympus Korea Co., Ltd., Seoul, Korea). The canopy pictures were obtained for VR. A scale of 1 to 9 was used to rate the visual appearance of the groundcovers, where 1 = dead or dying plants, 6 = minimally acceptable, and 9 = ideal or optimum quality (Pittenger et al. 2001). Ground coverage, plant vigor, and color were taken into consideration during the rating process (Pittenger et al. 2001). In 2020, images from six data collection dates and in 2021, canopy pictures from 12 different data collection dates were used for the visual quality assessment. To maintain the consistency of rating, one person rated all the pictures using the same screen for all the images obtained in both years. Simple linear regression models were developed for all the species to evaluate the relationships between NDVI and VR. The models were subsequently used to identify minimum NDVI thresholds for each species equivalent to the VR value of six.

Statistical analysis. Data were analyzed using PROC GLIMMIX in SAS ver 9.4 (SAS Institute Inc., Cary, NC, USA). When data

Table 1. Growing season monthly, seasonal, and 30-yr average reference evapotranspiration (ET_o), precipitation, and air temperature obtained from the nearby California Irrigation Management Information System (CIMIS) weather station (CIMIS #44).

Month	ET_o (mm)			Precipitation (mm)			Air temp (°C)		
	2020	2021	1992–2021	2020	2021	1992–2021	2020	2021	1992–2021
May	184	164	157	0	0	5	20	18	18
Jun	163	188	174	1	4	2	21	22	21
Jul	208	206	189	0	3	2	24	25	24
Aug	197	181	183	0	0	2	27	25	25
Sep	161	149	142	0	0	3	25	23	23
Oct	122	102	103	0	11	8	22	18	19
Season	1,035	990	948	1	18	22	23	22	22




Fig. 1. Canopy pictures, the scientific name (italic and bold) and the common name of landscape groundcovers selected in this study.

from 2020 and 2021 were combined, there was a significant year effect. Also, the meteorological information, including precipitation, ET_o , and air temperature (Table 1), differed considerably between the 2020 and 2021 experimental seasons. Therefore, data were analyzed separately for each year. Also, because there were significant differences between species, each landscape groundcover was individually analyzed (Pittenger et al. 2001). The landscape groundcover *Baccharis* × ‘Starn’ Thompson was pruned just before the data collection on 22 May, 14 Oct, and 27 Oct in 2021; data from these dates for this specific plant were not included in the analysis because it would have skewed the results. Also, the groundcover *Eriogonum fasciculatum* ‘Warriner Lytle’ did not grow well in 2020, so data from 2021 were only included for this groundcover in the analysis. Irrigation treatments, data collection date, and interaction were used as fixed effects

for the response variables. Block and its interaction with irrigation treatment were random effects. The LSMEANS option LINES statement was used for pairwise least square mean comparisons, and treatment effects were considered significant at $\alpha = 0.05$.

Data in 2020 and 2021 for each groundcover (only from 2021 for *Eriogonum fasciculatum* ‘Warriner Lytle’) were pulled together to determine the relationship between NDVI and VR. The mean value of NDVI and VR for each species were obtained for all four irrigation treatments and each day of data collection. The regression option from the data analysis tool in Microsoft Excel 2016 was used to compute the regression statistics, identify the relationship’s significance, and get the coefficients of slope and intercepts for the linear regression equation. Graphs were made using GraphPad Prism version 9.3 (GraphPad Software, LLC, San Diego, CA, USA).

A simple linear regression-based GCWRF model for each groundcover was developed using 2 years of experimental data. Applied irrigation amount (percentage of ET_o), atmospheric evaporative demand (cumulative ET_o), and their interaction were used as predictor variables, and the NDVI was the response variable. The significant difference between the models was determined using the analysis of variance function in SAS. Long-term ET_o data (30 years) from CIMIS station 44 was used to identify minimum, mean and maximum daily ET_o values for six months (1 May–31 Oct). Then GCWRFs were used to estimate the response of groundcovers to four irrigation levels (80%, 60%, 40%, and 20% ET_o) under extremely low (minimum daily ET_o), high

(maximum daily ET_o), and mean (mean daily ET_o) atmospheric evaporative demands. The performance of models was evaluated using the coefficient of determination (R^2 ; Eq. [1]), mean absolute error (MAE; Eq. [2]), mean biased error (MBE; Eq. [3]), and the root mean square error (RMSE; Eq. [4]).

$$R^2 = 1 - \frac{\sum_{i=1}^N (M_i - E_i)^2}{\sum_{i=1}^N (M_i - \bar{M})^2} \quad [1]$$

$$MAE = \frac{1}{N} \sum_{i=1}^N |E_i - M_i| \quad [2]$$

$$MBE = \frac{1}{N} \sum_{i=1}^N (E_i - M_i) \quad [3]$$

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (E_i - M_i)^2} \quad [4]$$

where N is the total number of observations, M_i is the measured and E_i is the predicted value of i^{th} observation, and \bar{M} is the mean of the measured values.

Results

Performance of the ET-based smart irrigation controller. Table 2 summarizes the irrigation treatment values, programmed irrigation rates, and the actual irrigation applied as percentages of CIMIS- ET_o . The irrigation controller overirrigated the landscape groundcovers by an average of 7.7% (range: 7.5% to 8.7%) in 2020 and 4.7% (range: 3.2% to 7.1%) in 2021. The controller closely followed the programmed watering days (irrigation frequency).

Impact of irrigation on the VR of landscape groundcovers. In 2020, the data collection date (i.e., time of the season) significantly affected ($P < 0.01$) the VR (Table 3, Fig. 2) of

Table 2. Irrigation treatments implemented in the study in 2020 and 2021.

Irrigation		Percentages of ET_o			
2020	Treatment	20	40	60	80
	Programmed ⁱ	23	47	70	93
	Applied ⁱⁱ	25	51	75	99
2021	Treatment	20	40	60	80
	Programmed	23	47	70	93
	Applied	24	49	75	96

ⁱ Programmed irrigation is equal to treatment levels divided by irrigation efficiency of 86%.

ⁱⁱ Applied irrigation is equal to actual irrigation applications based on the precipitation rates of the irrigation system and flowmeter data. ET_o = reference evapotranspiration.

Table 3. Analysis of variance table presenting the effect of irrigation treatments, date of data collection, and their interaction on the normalized difference vegetation index (NDVI) and visual rating (VR) of different groundcover species in the years 2020 and 2021.

ET ₀ (%)	<i>Rhagodia spinescens</i>		<i>Baccharis</i> × 'Starn'		<i>Eremphila glabra</i> 'Mingew Gold'		<i>Lonicera japonica</i>		<i>Ruschia lineolata nana</i>		<i>Trachelospermum jasminoides</i>		<i>Lantana montevidensis</i>		<i>Rosmarinus officinalis</i> 'Roman Beauty'		<i>Oenothera stubbei</i>		<i>Eriogonum fasciculatum</i> 'Warriner Lytle'	
	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR	NDVI	VR
2020																				
25	0.48 c ¹	6.94 b	0.50 a	6.56 a	0.51 c	5.5 c	0.51 c	5.28 b	0.55 b	6.56 b	0.34 c	3.56 b	0.36 c	4.17 c	0.50 b	5.67 b	0.43 c	5.72 ab	-	-
51	0.49 bc	7.61 ab	0.50 a	6.33 a	0.57 b	6.28 bc	0.59 b	5.94 b	0.66 a	8.39 a	0.49 b	5.44 ab	0.58 b	5.78 b	0.63 a	7.89 a	0.43 c	5.06 b	-	-
75	0.54 ab	8.61 a	0.52 a	6.89 a	0.61 b	6.94 ab	0.69 a	7.61 a	0.68 a	8.72 a	0.60 ab	5.67 a	0.64 ab	6.72 b	0.66 a	8.39 a	0.54 b	7.17 ab	-	-
99	0.56 a	7.89 ab	0.55 a	6.39 a	0.66 a	7.78 a	0.71 a	7.56 a	0.71 a	8.72 a	0.69 a	6.78 a	0.68 a	8.06 a	0.66 a	8.22 a	0.64 a	8.00 a	-	-
SE	0.01	0.34	0.01	0.23	0.01	0.22	0.04	0.42	0.02	0.35	0.05	0.53	0.04	0.47	0.02	0.41	0.02	0.74	-	-
<i>P</i> values																				
Irrigation (I)	0.031	0.059	0.178	0.411	0.001	0.003	0.0018	0.003	0.005	0.009	0.001	0.034	0.0006	0.0008	0.001	0.02	0.002	0.144	-	-
Date (D)	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.003	<0.0001	<0.0001	<0.0001	<0.0001	-	-
I × D	<0.0001	0.257	<0.0001	0.709	0.0001	0.003	0.0004	0.041	<0.0001	<0.0001	<0.0001	0.022	<0.0001	0.002	<0.0001	0.0004	<0.0001	0.055	-	-
2021																				
24	0.51 a	8.64 a	0.53 b	7.74 a	0.46 b	7.19 b	0.51 b	6.33 b	0.46 b	6.33 b	0.22 b	2.28 b	0.34 d	4.14 c	0.48 b	6.33 b	0.26 b	3.86 b	0.47 a	7.72 a
49	0.50 a	8.53 a	0.58 ab	7.89 a	0.65 a	8.86 a	0.52 b	6.42 b	0.62 a	8.61 a	0.51 a	6.36 a	0.46 c	6.03 b	0.68 a	8.86 a	0.34 b	5.19 b	0.49 a	8.06 a
75	0.54 a	8.86 a	0.58 ab	7.93 a	0.65 a	8.89 a	0.79 a	8.69 a	0.66 a	8.78 a	0.56 a	7.11 a	0.60 b	7.83 a	0.68 a	8.83 a	0.55 a	7.53 a	0.53 a	7.69 a
96	0.53 a	8.69 a	0.63 a	8.22 a	0.70 a	8.97 a	0.72 a	8.53 a	0.67 a	8.50 a	0.66 a	7.83 a	0.66 a	8.67 a	0.67 a	8.89 a	0.60 a	8.28 a	0.53 a	7.25 a
SE	0.02	0.12	0.01	0.21	0.03	0.26	0.04	0.19	0.03	0.42	0.06	0.66	0.03	0.35	0.01	0.17	0.03	0.55	0.02	0.51
<i>P</i> values																				
Irrigation (I)	0.564	0.353	0.106	0.635	0.001	0.008	0.008	0.0003	0.006	0.010	0.003	0.001	<0.0001	0.0005	<0.0001	<0.0001	0.0008	0.003	0.151	0.570
Date (D)	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.001	<0.0001	<0.0001	<0.0001	0.0002	0.013	0.358	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.198
I × D	0.999	0.538	0.0002	0.966	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.058	0.133	0.0002	<0.0001	<0.0001	<0.0001	0.151	<0.0001	0.195	0.502

¹ For each year, different letter assignment within the column represents statistical significance at $\alpha = 0.05$. SE = standard error of the mean.

all nine groundcovers (*Eriogonum* was not included in 2020). Irrigation rates and their interaction with data collection dates also had significant ($P \leq 0.05$; Table 3) impacts on the VR of six groundcovers except for *Rhagodia*, *Baccharis*, and *Oenothera*. In 2021, the effect of data collection dates on the VR was significant ($P \leq 0.001$; Table 3) for eight landscape groundcovers during the experimental season except for *Trachelospermum* ($P = 0.358$) and *Eriogonum* ($P = 0.198$). In addition, the varying irrigation rates also significantly ($P < 0.01$) affected the VR of seven groundcovers other than *Rhagodia*, *Baccharis*, and *Eriogonum* ($P > 0.05$). The interaction effect of irrigation and data collection date on VR was significant ($P < 0.001$; Table 3) for only six groundcovers excluding *Rhagodia*, *Baccharis*, *Trachelospermum*, and *Eriogonum*.

The mean VR values of *Rhagodia* for 75% and 99% ET₀ irrigation treatments were above the minimum VR threshold (VR = 6) for the whole experimental period in 2020 (Fig. 2). The VR values dropped below 6 in mid-August for 51% and 25% ET₀ irrigation treatments. In 2021, the lowest mean VR value was 7.33, and irrigation rates did not affect the VR values of this groundcover (Fig. 3). *Eriogonum*, which was evaluated only in 2021, was also not affected by irrigation rates (Table 3), and the VR values were above the minimum threshold of 6 (Fig. 3).

Groundcovers *Ruschia* and *Rosmarinus* had a similar trend in 2020 and 2021. Irrigation treatments $\geq 49\%$ ET₀ had significantly the same VR values and were well above the minimum threshold (Figs. 2 and 3). Only the plots with irrigation treatments $< 25\%$ ET₀ showed signs of water stress and VR values < 6 , mainly from August. In 2020, *Eremphila* had VR values < 6 for irrigation treatments 51% and 25% ET₀ (Fig. 2). In 2021, three irrigation treatments ($\geq 49\%$ ET₀) did not significantly affect the VR values (Fig. 3) and had VR values consistently above 6. *Eremphila* treated with 24% ET₀ irrigation treatment had significantly different VR values than the other three irrigation treatments; however, the lowest mean VR values were 6, suggesting the plants still maintained acceptable visual quality.

Groundcovers such as *Lantana*, *Oenothera*, and *Lonicera* mostly had VR values above the minimum acceptable threshold (i.e., VR = 6; Figs. 2 and 3) for the irrigation rates $\geq 75\%$ ET₀ in both years. *Lonicera* once had VR < 6 in September 2020 (Fig. 2). The VR values for the other two irrigation rates diminished as the experimental season progressed and dropped below the minimum acceptable threshold of 6.

Baccharis was not affected by any of the four irrigation treatments in 2020 and 2021, nor by the interaction effect of irrigation rates and data collection dates (Table 3, Figs. 2 and 3). In 2020, the VR values for all irrigation treatments started falling after the start of the experimental season. Beginning in August, VR values were mainly below the minimum threshold of 6 to be visually acceptable (Fig. 2). However, in 2021, for all irrigation rates and the whole experimental season,




Fig. 2. Visual rating (VR) of multiple groundcover species over the growing season in 2020 as affected by varying irrigation rates (25%, 51%, 75%, and 99% ET_0). For each groundcover species, VR values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance (i.e., VR < 6 on a scale of 1 to 9). Error bar represents the standard error of the means for each groundcover species during the growing season. ET_0 = reference evapotranspiration.

the VR values for *Baccharis* remained at or above the acceptable threshold of six (Fig. 3). Unlike *Baccharis*, *Trachelospermum* was greatly affected by irrigation treatments. The mean VR values were above the minimum threshold of 6 only for irrigation levels $\geq 96\%$ ET_0 for the whole experimental period (Figs. 2 and 3). The VR values were as low as two for the irrigation rates $\leq 25\%$ ET_0 .

Relationship between NDVI and VR. Figure 4 shows the relationship between the VR and NDVI of multiple groundcovers in this study. Minimum NDVI values identified for each groundcover are presented in Table 4. A statistically significant ($P < 0.001$) linear relationship was established for nine landscape groundcovers. *Eriogonum* did not yield a significant relationship, so a minimum NDVI threshold value for this groundcover was not developed. However, based on the VR (Fig. 3), it had maintained the minimum acceptable VR ratings throughout the experimental period in 2021; therefore, all the NDVI values (0.40–0.67) were within the acceptable quality range for this groundcover. The relationship between NDVI and VR for three groundcover species, including *Trachelospermum*, *Rosmarinus*, and *Oenothera* showed a strong correlation

with $R^2 \geq 0.80$. Landscape groundcovers, including *Baccharis*, *Lonicera*, *Ruschia*, and *Lantana*, showed a significant and robust correlation between NDVI and VR with $70 < R^2 < 80$ (Fig. 4). *Rhagodia* ($R^2 = 0.35$), and *Eremphila* ($R^2 = 0.49$) also showed a significant relationship between NDVI and VR (Fig. 4).

Impact of irrigation on NDVI of landscape groundcovers. In 2020, the data collection date (i.e., time of the season) significantly affected ($P < 0.001$) the NDVI readings (Table 3, Fig. 5) of all nine groundcovers (*Eriogonum* was not included in 2020). The effect of irrigation rates also was significant ($P \leq 0.03$; Table 3) on the NDVI values of all the groundcovers except *Baccharis* ($P = 0.178$; Table 3). The interaction between irrigation and the data collection date had significant ($P < 0.001$) effects on NDVI readings of all groundcovers in 2020. Like in 2020, the effect of data collection dates on the NDVI was significant ($P \leq 0.013$; Table 3) for all 10 landscape groundcovers in 2021 during the experimental season (Fig. 6). In addition, the varying irrigation rates also significantly ($P < 0.01$) affected the NDVI values of seven groundcovers except for *Rhagodia*, *Baccharis*, and *Eriogonum* ($P > 0.05$). The interaction

effect of irrigation rates and data collection date was significant for six groundcover species and groundcover, but *Rhagodia*, *Trachelospermum*, *Oenothera*, and *Eriogonum* were not significantly ($P > 0.05$) influenced by the interaction of irrigation rates and data collection dates.

Among all groundcovers, *Rhagodia* had slight variation of the NDVI values during the experimental season and between the irrigation treatments. *Rhagodia* with silvery green leaves had mean NDVI values ranging between 0.41 and 0.62 in 2020. On the basis of the fitted linear regression between NDVI and VR, the minimum NDVI threshold for *Rhagodia* was 0.46 (Table 4). For irrigation treatments $\geq 75\%$ ET_0 , this groundcover had NDVI values above the minimum threshold (0.46) throughout the experimental season in 2020. Additionally, for the irrigation rates of 25% and 51% ET_0 , the NDVI values were on the threshold borderline from late July to October, as shown in the light orange shaded region in Fig. 5. However, in 2021, the mean NDVI values (ranged between 0.46 and 0.58) for all irrigation rates and data collection dates were at or well above the acceptable minimum threshold. Figure 6 shows the performance of *Rhagodia* in 2021 for four irrigation rates from May to October. *Baccharis*




Fig. 3. Visual rating (VR) of multiple groundcover species over the growing season in 2021 as affected by varying irrigation rates (24%, 49%, 75%, and 96% ET_0). For each groundcover species, VR values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance (i.e., VR < 6 on a scale of 1 to 9). Error bar represents the standard error of the means for each groundcover species during the growing season. ET_0 = reference evapotranspiration.

mostly maintained its steady growth and health under all the irrigation rates in 2020 and 2021. However, the interaction effect of irrigation rates and data collection date was

significant. In 2020, its mean NDVI values ranged from 0.27 to 0.78, with the lowest recorded in early to mid-August. Similarly, the mean NDVI ranged from 0.43 to 0.69 in

2021. Given the NDVI threshold of 0.41 (Table 4), *Baccharis* maintained its acceptable quality for all irrigation rates in 2021 (Fig. 6). However, in 2020, it fell below that threshold




Fig. 4. Relationships between normalized difference vegetation index (NDVI) and visual rating (VR) of multiple groundcover species used in the study. Data in 2020 and 2021 for all groundcover (only from 2021 for *Eriogonum*) were combined to determine the relationship between NDVI and VR.

Table 4. Landscape groundcover water response functions and the minimum NDVI threshold developed for each groundcover to be minimally acceptable.

Scientific name	Minimum NDVI ⁱ	Water response function to estimate NDVI
<i>Rhagodia spinescens</i>	0.46	$0.55 + (2.73 \times 10^{-4} I^{ii}) - (1.47 \times 10^{-4} \text{CET}_0^{iii}) + (8.47 \times 10^{-7} I \text{CET}_0)$
<i>Baccharis</i> × ‘Starn’ Thompson	0.41	$0.73 - (9.30 \times 10^{-4} I) - (4.26 \times 10^{-4} \text{CET}_0) + (3.10 \times 10^{-6} I \text{CET}_0)$
<i>Eremphila glabra</i> ‘Mingenew Gold’	0.50	$0.66 + (2.21 \times 10^{-4} I) - (3.78 \times 10^{-4} \text{CET}_0) + (3.95 \times 10^{-6} I \text{CET}_0)$
<i>Lonicera japonica</i>	0.47	$0.77 + (3.70 \times 10^{-4} I) - (6.01 \times 10^{-4} \text{CET}_0) + (5.13 \times 10^{-6} I \text{CET}_0)$
<i>Ruschia lineolate nana</i>	0.48	$0.71 + (2.58 \times 10^{-4} I) - (4.11 \times 10^{-4} \text{CET}_0) + (3.72 \times 10^{-6} I \text{CET}_0)$
<i>Trachelospermum jasminoides</i>	0.53	$0.42 + (2.31 \times 10^{-3} I) - (4.07 \times 10^{-4} \text{CET}_0) + (5.10 \times 10^{-6} I \text{CET}_0)$
<i>Lantana montevidensis</i>	0.50	$0.53 + (2.53 \times 10^{-3} I) - (4.59 \times 10^{-4} \text{CET}_0) + (3.30 \times 10^{-6} I \text{CET}_0)$
<i>Rosmarinus officinalis</i> ‘Roman Beauty’	0.50	$0.68 + (3.96 \times 10^{-4} I) - (3.43 \times 10^{-4} \text{CET}_0) + (3.14 \times 10^{-6} I \text{CET}_0)$
<i>Oenothera stubbei</i>	0.44	$0.45 + (4.08 \times 10^{-3} I) - (3.95 \times 10^{-4} \text{CET}_0)$
<i>Eriogonum fasciculatum</i> ‘Warriner Lytle’	—	$0.57 + (8.48 \times 10^{-4} I) - (2.08 \times 10^{-4} \text{CET}_0)$

ⁱ NDVI = normalized difference vegetation index.

ⁱⁱ I = irrigation percentage expressed in terms of reference evapotranspiration.

ⁱⁱⁱ CET₀ = cumulative evapotranspiration (millimeters).

in August, and irrigation treatments 51% and 25% ET₀ struggled to rise above the threshold of 0.41 (Fig. 5).

Groundcovers *Ruschia* and *Rosmarinus* showed a similar trend of NDVI values in 2020 and 2021. Three irrigation rates ($\geq 49\%$ ET₀) had significantly the same NDVI values, and they were all above the minimum acceptable NDVI values of 0.48 and 0.50, respectively. Only the plots with $\leq 25\%$ ET₀ irrigation treatments showed signs of water stress, as reflected by the NDVI values. The NDVI values of both species deteriorated starting from late July and

dropped below the minimum acceptable NDVI threshold. Groundcover *Eremphila* showed a similar trend in 2021. For three irrigation rates ($\geq 49\%$ ET₀), the NDVI values were significantly the same and were above the established minimum acceptable NDVI threshold of 0.50. For 24% ET₀ irrigation treatments, the NDVI values dropped significantly below 0.5 starting in July (Fig. 6). In 2020, the NDVI values for *Eremphila* were above 0.50 minimum acceptable threshold only for two irrigation rates (75% and 99% ET₀). The NDVI values dropped significantly below the acceptable

threshold starting in July for 25% ET₀, and it remained at the borderline of the NDVI = 0.50 for the 75% ET₀ irrigation treatment for most of the experimental season in 2020 (Fig. 5).

Until mid-August 2020 (Fig. 5) and early August 2021 (Fig. 6), *Lonicera* grew well for all four irrigations without showing signs of drought injury. After that, the NDVI values for deficit irrigation treatments $\leq 51\%$ ET₀ fell below the desired threshold value of 0.47 and showed signs of water stress. In 2020, for September (Fig. 5), NDVI values for 60% ET₀ treatment were significantly dropped and came




Fig. 5. The normalized difference vegetation index (NDVI) of multiple groundcover species over the growing season in 2020 under varying irrigation rates (25%, 51%, 75%, and 99% ET₀). For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptable appearance. Error bar represents the standard error of the means for each groundcover species during the growing season. ET₀ = reference evapotranspiration.




Fig. 6. The normalized difference vegetation index (NDVI) of multiple groundcover species over the growing season in 2021 under varying irrigation rates (24%, 49%, 75%, and 96% ET_0). For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance. Error bar represents the standard error of the means for each groundcover species during the growing season. ET_0 = reference evapotranspiration.

close to the minimum threshold value of 0.47; however, groundcover *Lonicera* grew sufficiently for the 75% ET_0 treatment, and NDVI values started getting better. In both years (except for September 2020), the NDVI values for $\geq 75\%$ ET_0 irrigation treatments were significantly the same, with mean NDVI values for 75% ET_0 treatment being slightly more than that of $\geq 96\%$ ET_0 treatments (Figs. 5 and 6). For *Lantana*, in both years, irrigation treatments $\geq 75\%$ ET_0 had acceptable mean NDVI values above its NDVI threshold of 0.5 (Table 4). However, as the summer progressed, the NDVI values for all the irrigation rates decreased such that the NDVI values of the irrigation treatments $\leq 51\%$ ET_0 fell below the acceptable threshold showing visible water-stress symptoms. As a result, the minimum NDVI readings were only 0.22 and 0.21 for the lowest irrigation treatments in 2020 (Fig. 5) and 2021 (Fig. 6), respectively.

Oenothera maintained acceptable NDVI values (Figs. 5 and 6) for irrigation treatment $\geq 96\%$ ET_0 in both years. However, plants treated with irrigation treatments $\leq 51\%$ ET_0 showed signs of water stress and had mean NDVI values less than the NDVI threshold of 0.44 (Table 4). For 75% ET_0 irrigation treatment, the groundcover maintained the NDVI values above the minimum threshold of 0.44 for the experimental season in 2021, whereas it was right below that threshold in 2020 starting from August.

The mean NDVI values of *Trachelospermum* were significantly affected by irrigation rates and data collection dates in both years (Table 3). However, the interaction effect of

irrigation rates and data collection dates was significant ($P < 0.001$) only in 2020. In both years, the irrigation treatment $\geq 96\%$ ET_0 only had NDVI values well above the minimum required threshold (i.e., NDVI = 0.53; Table 4) for the whole experimental season (Figs. 5 and 6). The 75% ET_0 irrigation treatment had NDVI values above the threshold in 2020, but it was not the case in 2021. Plants at $< 25\%$ ET_0 treatment always had the lowest NDVI values. Switching back to nonlimiting irrigation between Nov 2020 and Apr 2021 did not help this species regenerate from the water stress. Hence, NDVI values were relatively lower in 2021 than in 2020.

Eriogonum did not grow well in 2020, so only 2021 data were processed and presented for the results. The NDVI was not significantly affected by different irrigation rates and their interaction with the data collection date (Table 3). For all four irrigation treatments, the NDVI values followed the same trend (Fig. 6). The maximum mean NDVI value recorded was 0.67, whereas the minimum was 0.40.

Groundcover water response function. Table 4 shows the GCWRFs developed using 2-year data for all 10 groundcover species. The relationships between the measured and the GCWRFs-estimated NDVI are presented in Fig. 7. The strength and accuracy of the models developed were presented in terms of the coefficient of correlation (R^2), MAE, MBE, and RMSE.

The cumulative ET_0 for scenarios minimum, mean, and maximum atmospheric evaporative demand based on long-term data were 403, 949,

and 1302 mm, respectively, for the experimental period from May to October. All groundcovers maintained their growth and aesthetic values at 80% ET_0 irrigation (Fig. 8) for all three scenarios except *Oenothera*, which only performed well under minimum atmospheric evaporative demand was minimum. A significant relationship between NDVI and VR was not obtained for *Eriogonum*. However, NDVI > 0.4 were considered acceptable for this species because all their corresponding VR values were above the minimum threshold. The NDVI of *Baccharis* and *Lantana* under maximum evaporative demand dropped to values close to their thresholds toward the end of October.

Under 60% ET_0 irrigation application, three landscape groundcover, including *Eremphila*, *Ruschia*, and *Rosmarinus* maintained their acceptable NDVI values for all weather scenarios (Fig. 9). Groundcovers *Rhagodia*, *Baccharis*, and *Lonicera* maintained their acceptable quality at 60% ET_0 irrigation only under minimum and mean atmospheric evaporative demands. Groundcovers *Trachelospermum*, *Lantana*, and *Oenothera* had NDVI values below the minimum threshold at 60% ET_0 irrigation rate for the scenario when the daily ET_0 equals the mean or maximum of the long-term average (Fig. 9).

Rhagodia maintained its quality for all three atmospheric evaporation demand at 40% ET_0 until early September, after which the quality of *Rhagodia* dropped below the acceptable threshold except for minimum atmospheric evaporative demand (Fig. 10). Groundcovers *Rosmarinus*, *Ruschia*, and *Baccharis* maintained NDVI above their thresholds at 40% ET_0 except under the maximum atmospheric




Fig. 7. Relationships between measured and estimated normalized difference vegetation index (NDVI) for multiple groundcovers obtained using landscape groundcover water response functions.

evaporative demand scenario. The NDVI for *Eremphila* and *Lonicera* fell below the acceptable threshold toward the end of the season under the mean atmospheric evaporative

demand scenario. The groundcover *Trachelospermum* could not maintain the acceptable NDVI threshold at 40% ET_o and *Lantana* and *Oenothera* only held the acceptable

NDVI values under the minimum atmospheric evaporative demand scenario.

Fig. 11 showed how groundcovers responded to 20% ET_o irrigation application




Fig. 8. Response of 10 landscape groundcovers to irrigation scenario equivalent to 80% reference evapotranspiration (ET_o) using the groundcover water response functions. The minimum, mean, and maximum scenarios represent minimum, mean, and maximum cumulative ET_o for that specific date based on the long-term weather data. For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance.




Fig. 9. Response of 10 landscape groundcovers to irrigation scenario equivalent to 60% reference evapotranspiration (ET_0) using the groundcover water response functions. The minimum, mean, and maximum scenarios represent minimum, mean, and maximum cumulative ET_0 for that specific date based on the long-term weather data. For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance.

under minimum, mean and maximum atmospheric evaporative demand scenarios. All groundcovers, except *Trachelospermum*, *Lantana*, and *Oenothera*, performed well under the minimum atmospheric evaporative

demand scenario. At 20% ET_0 irrigation application and mean atmospheric evaporative demand, six groundcovers maintained the acceptable NDVI threshold early in the experimental season, yet their quality fell below the

minimum threshold as the season progressed. A similar trend with a more pronounced reduction in NDVI values was observed under maximum atmospheric evaporative demand (Fig. 11). The NDVI values of groundcovers




Fig. 10. Response of ten landscape groundcovers to irrigation scenario equivalent to 40% reference evapotranspiration (ET_0) using the groundcover water response functions. The minimum, mean, and maximum scenarios represent minimum, mean, and maximum cumulative ET_0 for that specific date based on the long-term weather data. For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance.




Fig. 11. Response of 10 landscape groundcovers to irrigation scenario equivalent to 20% reference evapotranspiration (ET_0) using the groundcover water response functions. The minimum, mean, and maximum scenarios represent minimum, mean, and maximum cumulative ET_0 for that specific date based on the long-term weather data. For each groundcover species, NDVI values that fall in the light orange shaded region represent not meeting the minimal acceptance appearance.

Trachelospermum, *Lantana*, and *Oenothera* were severely decreased under the mean and maximum atmospheric evaporative demands.

Discussion

Performance of the smart irrigation controller. The overestimate of ET_0 by the Weathermatic SL4800 was similar to the 5% to 8% and 3% to 12% overestimation that we observed in two turfgrass irrigation trials in central and southern California, respectively (Haghverdi et al. 2021a, 2021c). A higher range of ET_0 overestimation (9% to 33%) was reported in a study done in Gainesville and Wimauma, FL (Rutland and Dukes 2014), which could be attributed to their different climatic conditions. Our results suggest that the Weathermatic SL4800 controller is reliable for autonomous irrigation scheduling in summer months in semi-arid environments. However, we recommend more studies to assess its accuracy in different climate regions based on long-term data. Also, the average ET_0 for the study periods was within 9% of the 30-year average. Likewise, individual monthly average cumulative ET_0 was within 18% of the long-term average (Table 1). Smart irrigation controllers showed better performance in estimating the irrigation needs; however, programming based on historical data also showed some potential but fluctuated sharply between months of the growing season.

NDVI as an indicator of groundcover growth and quality. The relationship between NDVI and VR was established for 10 landscape groundcovers in this study. We suggest NDVI as a proxy to quantify the growth and health of groundcovers in a fast and consistent

manner, given its high correlation ($0.35 \leq R^2 \leq 0.82$) with VR for almost all the species in this study. The minimum NDVI threshold (equal to VR of 6) ranged from 0.41 to 0.53 among species. We recommend that NDVI thresholds are established for each species separately since the NDVI values are impacted by each species' unique leaf and flowering characteristics (Shen et al. 2009, 2010). We also observed that pruning substantially reduced the NDVI values, especially in the case of woody-type groundcovers. Summer flowering may also impact the NDVI readings of some species, such as *Eremphila*, which bears yellow flowers. Yellow flowers reduce the NDVI readings by increasing the red band canopy reflectance without apparent variation in near-infrared reflectance (Shen et al. 2009). These not water-stress-related fluctuations in the NDVI values should be considered for large-scale remote sensing studies when ground-truth data might not be readily available.

Water conservation potential of the groundcover species. In this study, groundcover *Rhagodia* and *Eriogonum* showed the highest potential for performing well under limited water application. *Rhagodia* and *Eriogonum* formed a dense canopy with almost complete coverage, resisting evaporative loss and conserving soil moisture (Huang 2008). This helped these species stay green and healthy and maintain the acceptable VR at an irrigation rate of 24% to 25% ET_0 . This is roughly one-third of the minimum required irrigation application to sustain hybrid bermudagrass quality in the summer in inland Southern California (Haghverdi et al. 2021c). A slightly lower irrigation level (20% ET_0)

was recommended for *Rhagodia* and *Eriogonum* in coastal southern California (Sisneroz et al. n.a.). The GCWRF estimations suggest that under extremely high atmospheric evaporative demand, the best performing groundcovers (*Rhagodia*, *Eremphila*, *Ruschia*, and *Rosmarinus*) may require substantially higher irrigation applications (60% ET_0) to maintain the acceptable quality throughout the summer. Under the mean atmospheric evaporative demand scenario, however, the groundcovers (*Rhagodia*, *Baccharis*, *Ruschia*, and *Rosmarinus*) are expected to maintain their quality at the 40% ET_0 irrigation application rate (Fig. 10).

A minimum of 49% ET_0 irrigation was found to be sufficient for *Ruschia*, *Rosmarinus*, and *Eremphila* to keep the visually acceptable groundcover quality. A lower irrigation level of 20% ET_0 was reported to maintain the acceptable quality by (Sisneroz et al. n.d.) for *Ruschia* species in Davis and Irvine, CA. This is attributed to differences in field and weather conditions between the two studies, including a) full sun in our study vs. 50% shade in (Sisneroz et al. n.d.) and b) little to no rain in our study vs. considerable rainfall (177 mm) reported by (Sisneroz et al. n.d.) from April to October in Irvine. The least performing species was *Trachelospermum*, with a longer establishment time than other species and visible water stress symptoms even at a 75% ET_0 application rate. Overall, our results showed that groundcovers have irrigation water-saving potentials; however, not all groundcovers are drought-tolerant and perform at lower irrigation rates than the irrigation requirement of turfgrass species in the region. The use of NDVI to assess the quality of groundcovers might be a new normal, but it

comes with challenges, and further research that identifies the time and frequency of data collection for homeowners and stakeholders to get reliable and meaningful results are needed.

Conclusion

A 2-year (2020–21) field study evaluated the effect of deficit irrigation on the NDVI and visual quality rating of 10 landscape groundcovers in inland Southern California. Following are the main conclusions drawn:

- I The Weathermatic SL4800 smart irrigation controller showed a fair potential to schedule autonomous irrigation in summer in semiarid regions with slight overirrigation (on average 4.7% to 7.7%) compared with CIMIS-ET_o.
- II Development of plant-specific plant factors for irrigation scheduling is needed because groundcovers respond differently to different irrigation scenarios, and not all groundcovers can be drought-tolerant and withstand severe deficit irrigation. Three landscape groundcovers, including *Rhodia spinescens*, *Baccharis* × ‘Starn’ Thompson, *Eriogonum fasciculatum* ‘Warriner Lytle’ withstood water stress and maintained their growth and visual quality even at a 24% ET_o irrigation application. Groundcovers, *Ruschia lineolata* nana, *Rosmarinus officinalis* ‘Roman Beauty’, and *Eremphila glabra* have the potential to perform well with ≥49% ET_o irrigation. Landscape groundcovers *Lantana montevidensis*, *Oenothera stubbei*, and *Lonicera japonica* required 75% ET_o or more irrigation to maintain their growth and acceptable visual appearance. Results showed that *Trachelium jasminoides* require a more extended establishment period before deficit irrigation is imposed.
- III NDVI showed a potential to monitor the growth and quality of landscape groundcovers in a fast and consistent manner; however, the growth stages and maintenance activities can affect the readings. Therefore, NDVI should be evaluated, and minimum thresholds should be established for each groundcover. In this study, a minimum NDVI threshold was identified for multiple landscape groundcovers which can be used as an alternative to VR.

References Cited

- Bell GE, Martin DL, Koh K, Han HR. 2009. Comparison of turfgrass visual quality ratings with ratings determined using a handheld optical sensor. *HortTechnology*. 19:309–316. <https://doi.org/10.21273/hortsci.19.2.309>.
- Bremer DJ, Lee H, Su K, Keeley SJ. 2011. Relationships between normalized difference vegetation index and visual quality in cool-season turfgrass: I. Variation among species and cultivars. *Crop Sci*. 51:2212–2218. <https://doi.org/10.2135/cropsci.2010.12.0728>.
- Costello L, Jones K. 2014. Water use classification of landscape species: WUCOLS IV 2014. California Department of Water Resources, Davis, CA. https://ucanr.edu/sites/WUCOLS/WUCOLS_IV_User_Manual/. [accessed 16 Jan 2023].
- Davison E. 1999. Ground covers for Arizona landscapes. Cooperative Extension Publication AZ1110. College of Agriculture and Life Sciences, University of Arizona, Tucson, AZ.
- Dukes MD. 2020. Two decades of smart irrigation controllers in U.S. landscape irrigation. *ASABE*. 63:1593–1601. <https://doi.org/10.13031/trans.13930>.
- Easterday K, C. Kislik C, Dawson TE, Hogan S, Kelly M. 2019. Remotely sensed water limitation in vegetation: Insights from an experiment with unmanned aerial vehicles (UAVs). *Remote Sensing*. 11:1853. <https://doi.org/10.3390/rs11161853>.
- Fitz-Rodríguez E, Choi CY. 2002. Monitoring turfgrass quality using multispectral radiometry. *Trans. ASAE*. 45:865. <https://doi.org/10.13031/2013.8839>.
- García-Navarro MC, Evans RY, Montserrat RS. 2004. Estimation of relative water use among ornamental landscape species. *Sci. Horticulturae*. 99:163–174. [https://doi.org/10.1016/S0304-4238\(03\)00092-X](https://doi.org/10.1016/S0304-4238(03)00092-X).
- Garg A, Sapkota A, Haghverdi A. 2022. SAMZ-Desert: A satellite-based agricultural management zoning tool for the desert agriculture region of Southern California. *Computers Electronics Agric*. 194:106803. <https://doi.org/10.1016/j.compag.2022.106803>.
- Haghverdi A, Reiter M, Sapkota A, Singh A. 2021a. Hybrid bermudagrass and tall fescue turfgrass irrigation in central California: I. Assessment of visual quality, soil moisture and performance of an ET-based smart controller. *Agronomy*. 11:1666. <https://doi.org/10.3390/agronomy11091666>.
- Haghverdi A, Reiter M, Singh A, Sapkota A. 2021b. Hybrid bermudagrass and tall fescue turfgrass irrigation in central California: II. Assessment of NDVI, CWSI, and canopy temperature dynamics. *Agronomy*. 11:1733. <https://doi.org/10.3390/agronomy11091733>.
- Haghverdi A, Singh A, Sapkota A, Reiter M, Ghodsi S. 2021c. Developing irrigation water conservation strategies for hybrid bermudagrass using an evapotranspiration-based smart irrigation controller in inland southern California. *Agric. Water Manage*. 245:106586. <https://doi.org/10.1016/j.agwat.2020.106586>.
- Hargreaves GH, Samani ZA. 1985. Reference crop evapotranspiration from temperature. *Appl. Engineer. Agric*. 1:96–99. <https://doi.org/10.13031/2013.26773>.
- Horst GL, Engelke MC, Meyers W. 1984. Assessment of visual evaluation techniques. *Agron. J*. 76:619–622. <https://doi.org/10.2134/agronj1984.00021962007600040027x>.
- Huang B. 2008. Turfgrass water requirements and factors affecting water usage. *Water quality and quantity issues for turfgrass in urban landscapes*. Council Agric Sci Technol Spec Publ. 27:193–205.
- Huang S, Tang L, Hupy JP, Wang Y, Shao G. 2021. A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *J Forestry Res*. 32:1–6. <https://doi.org/10.1007/s11676-020-01155-1>.
- Leinauer B, VanLeeuwen DM, Serena M, Schiavon M, Sevostianova E. 2014. Digital image analysis and spectral reflectance to determine turfgrass Quality. *Agron J*. 106:1787–1794. <https://doi.org/10.2134/agronj14.0088>.
- Luscier JD, Thompson WL, Wilson JM, Gorham BE, Dragut LD. 2006. Using digital photographs and object-based image analysis to estimate percent ground cover in vegetation plots. *Front Ecol Environ*. 4:408–413. [https://doi.org/10.1890/1540-9295\(2006\)4\[408:UDPAOI\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)4[408:UDPAOI]2.0.CO;2).
- Nazemi Rafi Z, Kazemi F, Tehranifar A. 2019. Effects of various irrigation regimes on water use efficiency and visual quality of some ornamental herbaceous plants in the field. *Agric Water Manage*. 212:78–87. <https://doi.org/10.1016/j.agwat.2018.08.012>.
- Pittenger DR, Shaw DA, Hodel DR, Holt DB. 2001. Responses of landscape groundcovers to minimum irrigation. *J Environ. Hortic*. 19:78–84. <https://doi.org/10.24266/0738-2898-19.2.78>.
- Rutland DC, Dukes MD. 2014. Accuracy of reference evapotranspiration estimation by two irrigation controllers in a humid climate. *J. Irrigation Drainage Engineer*. 140(6):04014011. [https://doi.org/10.1061/\(ASCE\)IR.1943-4774.0000720](https://doi.org/10.1061/(ASCE)IR.1943-4774.0000720).
- Serena M, Velasco-Cruz C, Friell J, Schiavon, M Sevostianova E, Beck L, Sallenave R, Leinauer B. 2020. Irrigation scheduling technologies reduce water use and maintain turfgrass quality. *Agron J*. 112:3456–3469. <https://doi.org/10.1002/agi2.20246>.
- Shen M, Chen J, Zhu X, Tang Y. 2009. Yellow flowers can decrease NDVI and EVI values: Evidence from a field experiment in an alpine meadow. *Can J Remote Sensing* 35:99–106. <https://doi.org/10.5589/m09-003>.
- Shen M, Chen J, Zhu X, Tang Y, Chen X. 2010. Do flowers affect biomass estimate accuracy from NDVI and EVI? *Int J Remote Sensing* 31:2139–2149. <https://doi.org/10.1080/01431160903578812>.
- Shober AL, Davis S, Dukes MD, Denny GC, Brown SP, Vyapari S. 2009. Performance of Florida landscape plants when irrigated by ET-based controllers and time-based methods. *J Environ Hortic*. 27:251–256. <https://doi.org/10.24266/0738-2898-27.4.251>.
- Sisneros JA, Reid K, Oki L, Haver D, Fujino D. n.d. 2018–2020 UC landscape plant irrigation trials. University of California Cooperative Extension. University of California Agricultural and Natural Resources. <https://ucanr.edu/sites/UCLPIT/files/353920.pdf> [accessed 22 Mar 2022].
- US Bureau of Reclamation. 2008. Summary of smart controller water savings studies. US Department of the Interior Bureau of Reclamation. Final Technical Memorandum No. 86-68210-SCAO-01. <https://ucanr.edu/sites/UrbanHort/files/80222.pdf> [accessed 21 Dec 2022].
- Wang T, Chandra A, Jung J, Chang A. 2022. UAV remote sensing based estimation of green cover during turfgrass establishment. *Computers Electronics Agric*. 194:106721. <https://doi.org/10.1016/j.compag.2022.106721>.