

Evaluation of Shrubs for Side Slope Greening and Protection in Urban Landscape

Gabriele Amoroso^{1,3}, Piero Frangi¹, Riccardo Piatti¹, Alessio Fini², Francesco Ferrini², and Marco Faoro¹

ADDITIONAL INDEX WORDS. mulch, low-maintenance, ornamental value, plant management, revegetation, weed control

SUMMARY. Shrubs are often used for side slope greening and protection in the urban landscape. Only a few species are commonly used in the European city environment, bestowing upon cities a monotonous appearance. The aim of this 3-year trial, set up at Fondazione Minoprio (Vertemate con Minoprio; Como, Italy), was to evaluate the performance of 25 shrub species grown on a slope during three seasons (2007–09). Moreover, to evaluate the influence of weeds on plant growth and the weeding time, two mulches (biodegradable textile and polypropylene fabric) were used in comparison with bare soil (control). Shrubs were planted in late Spring 2007. To simulate urban conditions, no pruning or disease control were applied. Irrigation was carried out only in the driest periods of the first summer. Plant height and percent cover were measured every 2 months, whereas plant phenology and state of health were recorded weekly. Chlorophyll fluorescence and chlorophyll content were evaluated on two different drought-tolerant species during the third growing season. At the end of the trial shoot biomass was measured and root characteristics (root density and specific root length) were determined. Weeds were removed twice in the first and third years and three times in the second year. Time for weed removal was recorded for each experimental plot. Results show that the highest growth (height and biomass) was detected in mulched plots, probably because of both limited weed competition for water and nutrients and lower water loss by evaporation. Plants grown in bare soil showed higher root density and finer roots compared with mulched plants; this may be explained by the necessity of the plant root system to explore the soil to reach for water. Differences in growth, groundcovering, and root characteristics were observed among species. As a result of higher mulching cost and the poorer root characteristics of mulched plants, bare soil and fast growing shrubs should be used to limit weed competition and assure a satisfactory slope greening and consolidation.

During the last few years only a few species have been regularly used in the European urban landscape [i.e., herbaceous species or evergreen shrubs such as cotoneaster (*Cotoneaster* spp.) and honeysuckle (*Lonicera* spp.)], but many other species and cultivars could be suitable for this purpose (Assone et al., 2000). Current trends in urban landscaping require the use of low-maintenance species in response to the reduced water availability for

irrigation of urban greenspaces and to the low skill of the maintenance workforce (Franco et al., 2006). Moreover, it is necessary to limit weed competition with transplanted plants to reduce the need for weeding and chemical treatments. Mulch could be successfully used for this purpose (Chalker-Scott, 2007; Ferrini et al., 2008; Stinson et al., 1990). Mulching benefits plant growth and improvement of soil characteristics (Chalker-Scott,

2007), but little information is available on mulch effect on root characteristics of shrub species.

The importance of vegetation on erosion control and slope consolidation is widely documented. In general, vegetation stabilizes slopes (Cazzuffi and Tironi, 2003; Macdonald and Witek, 1994; Stokes et al., 2009). The protective role of vegetation can be explained by many different hydromechanical mechanisms ranging from mechanical reinforcement and restraint by the roots to modification of slope hydrology as a result of rain drop interception by the foliage and soil moisture extraction through transpiration (Gray and Leiser, 1982; Gray and Satir, 1996; Morgan and Rickson, 1994). In urban areas, grass, forbs, or shrubs can be used for slope covering. Grass or other herbaceous vegetation provide protection against rain and wind erosion of surface soils. Conversely, shrubs are more effective in stabilizing soil due to their deep roots (Gray and Satir, 1996). The objective of this work was to measure shoot growth and root characteristics of 25 shrubs species over 3 years under a low-maintenance regime and three mulching systems.

Materials and methods

This experiment was carried out at Fondazione Minoprio (Vertemate con Minoprio, Como, Italy) (lat. 45°44'N, long. 9°04'E, elevation 342 m) over 3 years (2007–09). Daily temperature and rainfall were recorded and monthly averages reported (Fig. 1). The study area was a uniform south-facing slope at a 45° average inclination angle. Before planting, two herbicide treatments were made on 11 Nov. 2006 and 14 Mar. 2007 using broadcast glyphosate (Roundup® 450 Plus; Monsanto, Creve Coeur, MO) and pendimethalin (Stomp® 330-E; BASF, Ludwigshafen,

This work was conducted under a research project “ARBOVERDE—Valutazione di specie arbustive ornamentali per il rinverdimento di scarpate” financed by Regione Lombardia - Agricultural Department, according to the Plan of Research and Development 2006. We gratefully acknowledge Professor Daniel K. Struve (Department of Horticulture and Crop Science, The Ohio State University) for his critical review of the manuscript.

¹Fondazione Minoprio, Centro MiRT, Viale Raimondi, 54, 22070 Vertemate con Minoprio (CO), Italy

²Department of Plant, Soil and Environmental Science, University of Florence, Viale delle Idee, 30, 50019 Sesto Fiorentino (FI), Italy

³Corresponding author. E-mail: g.amoroso@fondazioneminoprio.it.

Units

To convert U.S. to SI, multiply by	U.S. unit	SI unit	To convert SI to U.S., multiply by
29.5735	fl oz	mL	0.0338
0.3048	ft	m	3.2808
0.0929	ft ²	m ²	10.7639
3.7854	gal	L	0.2642
2.54	inch(es)	cm	0.3937
25.4	inch(es)	mm	0.0394
16.3871	inch ³	cm ³	0.0610
1.1209	lb/acre	kg·ha ⁻¹	0.8922
28.3495	oz	g	0.0353
62.5000	oz/lb	g·kg ⁻¹	0.0160
(°F - 32) ÷ 1.8	°F	°C	(1.8 × °C) + 32

Germany) at 38 and 16 mL/100 L, respectively. Ten soil samples were randomly collected from the site and analyzed in Jan. 2007 (Table 1) (Osmond et al., 1997). On the basis of the soil analysis, calcium carbonate (1500 kg·ha⁻¹) and 5N-4.5P-25K (750 kg·ha⁻¹) mineral fertilizer were incorporated 7 cm into the surface in March. In early May 2007, 25 shrub species (Table 2) were obtained from local nurseries in 3-L containers. Shrubs were planted at a density of 4 plants/m². Within a species, all plants came from the same Italian nursery, but not all species came from the same nursery. Two mulches: 1) black polypropylene fabric (3210NE Agritela; Arrigoni, Uggiate Trevano, Italy) and 2) biodegradable textile made of natural fibers [coconut (*Cocos nucifera*), cotton (*Gossypium* spp.), and jute (*Corchorus olitorius*)] [80 g·m⁻² (Isomatural®; Veris srl, Codogno, Italy)], and a non-mulch control were used.

The experiment was a split-plot design with three replicates, where soil management was the main plot and species was the subplot. The size of each subplot was 8 m² for a total experimental area of 1800 m². Two hundred and eighty-eight plants for each species were used in the experiment for a total of 7200 plants. No irrigation or pruning was carried out during the trial, except in the first summer (2007) when plants were weekly irrigated during the driest weeks using a drip irrigation system (seven irrigation events delivering 10 mm·h⁻¹, the irrigation was carried out over 3 h). According to the standard weed management (Rice, 1992), herbicides were applied as a granular broadcast application twice during 2007 on the non-mulched plots: isoxaben + trifluralin [Gallery® T-DG (Dow Agroscience, Indianapolis, IN) at 20.5 kg·ha⁻¹] and 3.6 kg·ha⁻¹ oxadiazon (Ronstar®; Bayer CropScience, Monheim am Rhein, Germany).

Canopy height and percent cover were measured bimonthly during the experiment. Plant height was measured on four randomized plants per subplots while percent cover was assessed visually using a four class scale: 1 = <25%, 2 = 25% to 50%, 3 = 50% to 75%, and 4 = >75%. Plant disease symptoms and phenology were recorded weekly. Weeds were removed by hand twice in the first and third growing seasons and three times during the second growing




Fig. 1. Monthly average temperature and monthly rainfall over 3 years (2007–09); (1.8 × °C) + 32 = °F, 1 mm = 0.0394 inch.

Table 1. Soil physical and chemical properties at the beginning of the trial (Jan. 2007). The parameters reported in the table were obtained collecting soil samples in 10 randomly zones in the slope area. Soil samples were collected at 20-cm (7.9 inches) depth.

Gravel > 2 mm (g·kg ⁻¹) ²	Sand < 2 mm (g·kg ⁻¹)	Silt < 0.05 mm (g·kg ⁻¹)	Clay < 0.002 mm (g·kg ⁻¹)	pH	Carbon/nitrogen ratio
202.4	565.1	412.3	22.6	6.45	11.16

²1 mm = 0.0394 inch, 1 g·kg⁻¹ = 0.0160 oz/lb.

Table 2. List of the 25 shrubs planted on a slope to evaluate their establishment, root characteristics, and canopy groundcover ability under low-maintenance conditions in different types of soil management (bare soil and mulching with biodegradable textile or polypropylene fabric).

Common name	Scientific name
Blue-spirea	<i>Caryopteris × clandonensis</i> 'Heavenly Blue'
Border forsythia	<i>Forsythia × intermedia</i> 'Lynwood'
Boxleaf honeysuckle	<i>Lonicera nitida</i>
Bumald spirea	<i>Spiraea japonica</i> 'Anthony Waterer'
Bush cinquefoil	<i>Potentilla fruticosa</i> 'Goldfinger'
Buttercup winterhazel	<i>Corylopsis pauciflora</i>
Common ninebark	<i>Physocarpus opulifolius</i> 'Diablo'
Common seabuckthorn	<i>Hippophae rhamnoides</i>
Doublefile viburnum	<i>Viburnum plicatum</i> 'Shasta'
English ivy	<i>Hedera helix</i> 'Gold heart'
Fragrant viburnum	<i>Viburnum farreri</i>
Fuzzy deutzia	<i>Deutzia scabra</i> 'Pride of Rochester'
Glossy abelia	<i>Abelia × grandiflora</i>
Goldcup st. johnswort	<i>Hypericum patulum</i> 'Hidcote'
Hybrid deutzia	<i>Deutzia hybrida</i> 'Strawberry Fields'
Japanese kerria	<i>Kerria japonica</i> 'Picta'
Privet honeysuckle	<i>Lonicera pileata</i>
Purpleosier willow	<i>Salix purpurea</i> 'Nana'
Redosier dogwood	<i>Cornus sericea</i> 'Kelsey'
Scorpion senna	<i>Coronilla emerus</i> (synonym: <i>Hippocrepis emerus</i>)
Shrub althea	<i>Hibiscus syriacus</i>
Slender deutzia	<i>Deutzia gracilis</i>
Spanish broom	<i>Spartium junceum</i>
Spanish gorse	<i>Genista lydia</i>
Virginal mock orange	<i>Philadelphus × virginialis</i> 'Minnesota Snowflake'

season. Time needed for weed removal was recorded for each experimental subplot. At the end of the 2nd year, because of the high mortality in non-mulched subplots, doublefile viburnum, english ivy, and fragrant viburnum were excluded from the experiment. In the third growing season leaf chlorophyll content, the maximal quantum yield of photosystem II [variable fluorescence (F_v)/maximum fluorescence (F_m)] and fluorescence transient were measured on goldencup st. johnswort and shrub althea [considered drought-tolerant (Burroughs, 2010) and drought-sensitive (Yeager et al., 2006) species, respectively]. Leaf chlorophyll content was measured on 25 June, 3 July, and 6 Aug. with a chlorophyll meter (SPAD-502; Minolta, Osaka, Japan) on five randomly chosen plants per block (for a total of 30 plants per species). Three youngest fully expanded leaves for each plant were used for this assessment. A clear relationship between chlorophyll meter readings and total leaf chlorophyll concentration was reported by Percival et al. (2008). F_v/F_m and fluorescence transient were measured three times during the driest period of the third growing season (on 3 July, 6 Aug., and 26 Aug. using five randomly selected plants per subplot and one fully expanded leaf per plant). Leaves were adapted to the dark for 30 min before F_v/F_m and fluorescence transient measurements using shading clips. Determinations of F_v/F_m and fluorescence transient were carried out using a portable fluorimeter (Handy Pea Plant Efficiency Analyser; Hansatech Instruments, Norfolk, United Kingdom) and recorded for up to 1 s, with data acquisition every 10 μ s for the first 2 ms, and every 1 ms thereafter, with 12-bit resolution (Strasser et al., 2000). The fluorescence transients were introduced by red light (peak at 650 nm) at 600 μ mol·m⁻²·s⁻¹ photons provided by an array of six light-emitting diodes. Upon the application of a saturating flash of actinic light, fluorescence raises from the ground state value (F_0) to its maximum value, F_m . This allows the determination of the maximal quantum yield of photosystem II (F_v/F_m) (Pinior et al., 2005). Fluorescence values at time intervals corresponding to the step O-J-I-P were recorded and used as original data in the OJIP-test (Strasser and Tsimilli-Michael, 2001).

These include the F_m (step P), the fluorescence intensity at 50 μ s [F_0 (step O)], 100 μ s (F_{100}), 300 μ s (F_{300}), 2 ms (step J), and 30 ms (Step I).

In Dec. 2009, four plants per subplot were cut at the root flare and fresh biomass determined by weighing the plants in the field with a digital balance (440–53N; Kern & Sohn, Balingen-Frommern, Germany) (36 plants per species). Four soil cores were collected randomly from each subplot in Dec. 2009 using a plant root sampler (Profile sampler 05.08; Eijkelkamp, Giesbeek, The Netherlands).

The volume of soil cores was 1250 cm³ (25 × 10 × 5 cm). Samples were collected in between two plants of the same species (≈70 cm from the root flares). After sampling, collected cores were separated in two sections: 0–10 cm and 10–25 cm deep. Samples were placed in plastic bags and held at 7 °C until January. In Jan. 2010, samples were sifted twice first over a 0.5 cm grid and then over a 0.2 cm. Root length was estimated by the intercept method (Tennant, 1975). Root samples were oven-dried (104 °C), and dry weights were recorded. Root density

Table 3. Hand weeding time required in each growing season (weed removal was carried out twice during the first and third growing season and three times during the second growing season) for 25 shrubs grown on a slope under different types of soil management (bare soil and mulched with biodegradable textile or polypropylene fabric). Shrubs were transplanted at a density of 4 plants/m² (0.37 plant/ft²) in May 2007 from 3-L (0.8 gal) containers.

	Hand weeding time (s·m ⁻²) ^z							
	2007		2008		2009		Total	
Species (SP)								
Blue-spirea	66	c-e ^y	116	i-k	50	cd	232	e-g
Border forsythia	92	c-e	145	g-k	24	d	261	d-g
Boxleaf honeysuckle	85	c-e	260	b-d	61	b-d	406	bc
Bumald spirea	87	c-e	153	f-k	9	d	249	d-g
Bush cinquefoil	82	c-e	175	e-j	46	cd	303	c-f
Buttercup winterhazel	74	c-e	188	d-i	42	cd	304	c-f
Common ninebark	60	de	135	g-k	22	d	217	d-f
Common seabuckthorn	118	bc	250	c-e	128	ab	496	b
Doublefile viburnum	77	c-e	214	d-g	—	—	—	—
English ivy	157	ab	348	a	—	—	—	—
Fragrant viburnum	76	ce	206	d-h	—	—	—	—
Fuzzy deutzia	102	c-e	112	i-k	16	d	230	e-g
Glossy abelia	74	c-e	120	i-k	15	d	209	e-g
Goldencup st. johnswort	105	c-e	102	i-k	12	d	219	e-g
Hybrid deutzia	100	c-e	100	jk	11	d	211	e-g
Japanese kerria	109	cd	170	e-j	72	ad	351	c-e
Privet honeysuckle	79	c-e	331	ab	120	a-c	530	ab
Purpleosier willow	54	e	72	k	10	d	136	g
Redosier dogwood	64	de	140	g-k	44	cd	248	d-g
Scorpion senna	68	c-e	107	i-k	52	b-d	227	e-g
Shrub althea	101	c-e	172	e-j	64	ad	337	c-f
Slender deutzia	94	c-e	234	d-f	67	a-d	395	b-d
Spanish broom	60	de	126	h-k	0	d	186	f-g
Spanish gorse	192	a	311	ac	138	a	641	a
Virginal mock orange	72	c-e	135	g-k	15	d	222	e-g
Weed management (WM)								
Polypropylene fabric	47	b	109	b	14	b	170	b
Biodegradable textile	44	b	121	b	24	b	189	b
Bare soil	174	a	301	a	106	a	581	a
Probability								
SP	***		***		***		***	
WM	***		***		***		***	
SP × WM	NS		NS		NS		NS	

^z1 s·m⁻² = 0.0929 s/ft².

^yMeans for each main factor and within each column followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

NS, ***Nonsignificant or significant at $P \leq 0.001$, respectively.

was calculated as root length over soil volume ratio [root length/soil volume (meters per liter)], and specific root length was calculated as the ratio between root length and root dry biomass [root length/root dry biomass (meters per gram)].

All data were processed with analysis of variance, or multivariate analysis of variance when two effects acted in combination. SPSS® (version 17.0; SPSS, Chicago, IL) was used. Means were separated by Duncan's multiple range test at a $P \leq 0.05$ level of significance.

Results and discussion

Without mulch, more time was required for weeding (Table 3). Despite the evident degradation shown by the biodegradable textile during the third growing season, no differences in hand weeding time were observed between plants mulched with this material and with polypropylene fabric over the experiment duration (Table 3). During the first two growing seasons complete canopy closure did not occur and the sunlight could reach the soil. During this period weed control by the biodegradable textile was very important in reducing weed competition with the shrubs, allowing the growth of a shading shrub canopy. After this period the mulch importance as weed regulator was decreased, but the degradation of the material did not affect the weed removal time.

Weeding spanish gorse subplots required the most hand labor, whereas purpleosier willow required the least (Table 3). Blue-spirea, border forsythia, bumald spirea, fuzzy deutzia, glossy abelia, goldencup st. johnswort, hybrid deutzia, redosier dogwood, scorpion senna, spanish broom, virginal mock orange, and purpleosier willow had the lowest hand weeding times (Table 3). Those species with the most rapid canopy closure required the least weeding time. At the end of the first growing season, blue-spirea, japanese kerria, scorpion senna, and spanish broom performed the highest percent cover regardless of the soil management. At the end of the second growing season, only english ivy, privet honeysuckle, spanish gorse, and fragrant viburnum showed the lowest values of canopy closure (Table 4). In relation to weed management, the polypropylene fabric allowed the highest groundcover ability

Table 4. Groundcover ability of 25 shrubs grown on a slope under different types of soil management (bare soil and mulched with biodegradable textile or polypropylene fabric). Shrubs were transplanted at a density of 4 plants/m² (0.37 plant/ft²) in May 2007 from 3-L (0.8 gal) containers.

Species (SP)	Canopy groundcover (1–4 scale) ^z					
	Oct. 2007		Oct. 2008		Oct. 2009	
Blue-spirea	3.89	a ^y	4.00	a	4.00	a
Border forsythia	2.89	ef	3.33	b–f	3.89	ab
Boxleaf honeysuckle	2.00	hi	2.78	f	3.44	a–c
Bumald spirea	2.33	gh	3.44	a–e	4.00	a
Bush cinquefoil	3.11	c–f	3.44	a–e	3.78	a–c
Buttercup winterhazel	2.11	hi	3.22	c–f	3.89	ab
Common ninebark	2.22	gh	3.89	ab	4.00	a
Common seabuckthorn	1.89	hi	3.00	ef	3.33	c
Doublefile viburnum	1.25	jk	2.88	ef	—	—
English ivy	1.00	k	1.11	i	—	—
Fragrant viburnum	1.33	jk	1.67	gh	—	—
Fuzzy deutzia	3.33	b–e	3.67	a–d	3.83	a–c
Glossy abelia	3.00	d–f	4.00	a	4.00	a
Goldencup st. johnswort	3.33	b–e	4.00	a	4.00	a
Hybrid deutzia	3.00	d–f	3.22	c–f	3.78	a–c
Japanese kerria	3.44	a–d	3.67	a–d	3.78	a–c
Privet honeysuckle	1.00	k	1.44	hi	1.33	e
Purpleosier willow	3.22	b–e	3.78	a–c	3.89	ab
Redosier dogwood	1.67	ij	3.11	d–f	3.56	a–c
Scorpion senna	3.67	ab	3.78	ac	3.67	a–c
Shrub althea	2.22	gh	3.44	a–e	3.78	a–c
Slender deutzia	2.00	hi	2.89	ef	3.38	bc
Spanish broom	3.56	a–c	4.00	a	4.00	a
Spanish gorse	1.11	k	2.00	g	2.17	d
Virginal mock orange	2.67	fg	3.44	a–e	3.89	ab
Weed management (WM)						
Polypropylene fabric	2.77	a	3.31	a	3.77	a
Biodegradable textile	2.36	b	3.49	a	3.77	a
Bare soil	2.19	c	2.68	b	3.35	b
Probability						
SP		***		***		***
WM		***		***		***
SP × WM		NS		NS		NS

^zGroundcover percentage was assessed visually: 1 = <25%, 2 = 25% to 50%, 3 = 50% to 75%, 4 = >75%.

^yMeans for each main factor and within each column followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

NS,***Nonsignificant or significant at $P \leq 0.001$, respectively.

starting from the first growing season, whereas from the 2nd year of trial both the tested mulches significantly increased the canopy coverage in comparison with the bare soil (Table 4).

An interaction between species and soil management was detected for plant shoot biomass (data not shown). When statistical differences were detected, plants mulched with polypropylene fabric showed higher shoot biomass compared with plants grown in bare soil with the only exception of scorpion senna (Table 5). Boxleaf and privet honeysuckle, redosier dogwood, and scorpion senna mulched with

biodegradable textile showed a greater shoot biomass compared with plants grown in bare soil (Table 5). As reported by Harris et al. (2004), mulching can increase soil water content by increasing percolation and retention, reducing evaporation and weed competition; moreover mulches can protect soils from extreme temperatures, keeping it cooler in hot conditions and warmer in cold conditions (Chalker-Scott, 2007; Einert et al., 1975; Ferrini et al., 2009; Long et al., 2001). The beneficial effect of mulches can explain, in species sensitive to water stress, the greater plant biomass compared with

non-mulched plants; the degradation of biodegradable textile may have partially reduced the mulching effect in buttercup winterhazel, common seabuckthorn, japanese kerria, shrub althea, and slender deutzia.

Soil covering with mulching materials did not affect shrub althea and goldencup st. johnswort chlorophyll content in June 2009; however, in the second assessment (carried out in early July) goldencup st. johnswort plants mulched with polypropylene textile showed a higher chlorophyll content than those growing on bare soil (Table 6). As reported in literature, F_v/F_m ratio is a reliable indicator of the occurrence of environmental stresses, including water stress, on photosystem II of several woody and herbaceous species (Angelopoulos et al., 1996; Lazár, 2006; Maxwell and Johnson, 2000; Percival and Fraser, 2001; Percival et al., 2006; Yamada et al., 1996). F_v/F_m ratio was not influenced by mulching in goldencup st. johnswort, whereas it was positively affected by polypropylene mulching in shrub althea (Table 6). This finding was also confirmed by chlorophyll a fluorescence transients, which provided evidence that shrub althea mulched with polypropylene fabric was less stressed during Aug. 2009 (Fig. 2). Shrub althea plants grown in bare soil or mulched with biodegradable textile highlighted changes in the acceptor side of photosystem I (IP-phase) (Tóth et al., 2007). In goldencup st. johnswort, chlorophyll a fluorescence transients were little affected by mulching; non-mulched plants showed moderate changes in the redox state of plastoquinone (JI-phase) and in IP-phase compared with plants mulched with polypropylene fabric (data not shown). A relationship between lower stress and higher shoot biomass of shrubs althea mulched with polypropylene can be assumed.

When vegetation is considered in models of slope stability, usually the only root parameters considered are tensile strength and root area ratio (Preti and Giadrossich, 2009; Tosi, 2007), but also the density and thickness of roots play a major role. Thin and fine roots are largely responsible for reinforcing soil on slopes; they act like soil nails on slopes, reinforcing soil (Ganatsas and Tsakalidimi, 2003; Stokes et al., 2009). In our experiment, the root system of plants grown

Table 5. Fresh shoot biomass of 25 shrubs grown on a slope under different types of soil management (bare soil and mulched with biodegradable textile or polypropylene fabric) at the end of the third growing season.

Species	Fresh shoot biomass (g/plant) ^z			Probability
	Polypropylene fabric	Biodegradable textile	Bare soil	
Blue-spirea	558 a ^y	634 a	546 a	NS
Border forsythia	1323 a	1310 a	887 a	NS
Boxleaf honeysuckle	1359 a	1214 a	643 b	*
Bumald spirea	766 a	910 a	624 a	NS
Bush cinquefoil	355 a	327 a	261 a	NS
Buttercup winterhazel	789 a	463 b	375 b	***
Common ninebark	920 a	1004 a	1108 a	NS
Common seabuckthorn	2035 a	1159 b	483 b	***
Fuzzy deutzia	924 a	1252 a	659 a	NS
Glossy abelia	959 a	1011 a	909 a	NS
Goldencup st. johnswort	1410 a	1461 a	1255 a	NS
Hybrid deutzia	830 a	1076 a	726 a	NS
Japanese kerria	1005 a	483 b	605 b	*
Privet honeysuckle	445 a	402 a	126 b	**
Purpleosier willow	1070 a	960 a	863 a	NS
Redosier dogwood	358 a	318 a	152 b	***
Scorpion senna	2034 ab	2800 a	1190 b	**
Shrub althea	1130 a	851 ab	640 b	*
Slender deutzia	480 a	356 b	335 b	**
Spanish broom	2883 a	3508 a	3014 a	NS
Spanish gorse	437 a	368 a	331 a	NS
Virginal mock orange	751 a	784 a	750 a	NS

^z1 g = 0.0353 oz.

^yMeans within the same row followed by different letters are significantly different from each other using the Duncan's mean separation test.

NS, *, **, *** indicate nonsignificant, significant at $P \leq 0.05$, 0.01, or 0.001, respectively.

Table 6. Chlorophyll content and chlorophyll fluorescence (F_v/F_m) in shrub althea and goldencup st. johnswort grown on a slope with different soil cover managements (bare soil and mulched with biodegradable textile or polypropylene fabric). Data were collected during the third growing season after transplanting.

Soil cover management	Chlorophyll content ^z			Chlorophyll fluorescence (F_v/F_m) ^y		
	25 June 2009	3 July 2009	6 Aug. 2009	3 July 2009	6 Aug. 2009	26 Aug. 2009
Shrub althea						
Polypropylene fabric	41.3	42.3	44.6	0.823	0.819 a ^x	0.818 a
Biodegradable textile	41.7	40.5	42.3	0.796	0.770 b	0.763 b
Bare soil	40.1	41.4	42.3	0.788	0.759 b	0.748 b
Significance	NS	NS	NS	NS	**	***
Goldencup st. johnswort						
Polypropylene fabric	55.1	57.5 a	49.3	0.716	0.748	0.757
Biodegradable textile	54.8	55.0 ab	50.2	0.717	0.756	0.726
Bare soil	52.2	52.6 b	46.2	0.702	0.723	0.711
Significance	NS	*	NS	NS	NS	NS

^zChlorophyll content measured with a chlorophyll meter.

^yChlorophyll fluorescence measured with a portable fluorimeter; F_0 = fluorescence intensity at 50 μ s, F_m = maximal fluorescence intensity, F_v = maximal variable fluorescence, $F_v = F_m - F_0$.

^xFor each species means within the same column followed by different letters are significantly different from each other using the Duncan's mean separation test.

NS, *, **, *** indicate nonsignificant, significant at $P \leq 0.05$, 0.01, or 0.001, respectively.

in bare soil had higher specific root length than mulched plants at both soil depth zone investigated; no differences between mulching materials

were observed (Table 7). Among species, the highest specific root length roots, in the first 10-cm soil depth, were observed in privet honeysuckle




Fig. 2. Chlorophyll *a* fluorescence induction curves in shrub althea grown on a slope with different soil cover managements (polypropylene textile, biodegradable textile, and bare soil). Data collected on 26 Aug. 2009. Each single curve represents the average of 10 independent chlorophyll fluorescence measurements; O step = fluorescence intensity at 50 μ s, J step = fluorescence intensity at 2 ms, I step = fluorescence intensity at 30 ms, P step = maximal fluorescence intensity.

Table 7. Root density (root length/soil volume) and specific root length (root length/dry root biomass), recorded at two soil depths [0–10 cm (3.9 inches) and 10–25 cm (9.8 inches) soil depth] in 25 shrubs grown on a slope under different types of soil management (bare soil and mulched with biodegradable textile or polypropylene fabric) at the end of the third growing season.

	Root density ($m \cdot L^{-1}$) ^z		Specific root length ($m \cdot g^{-1}$) ^z at soil depth	
	0–10 cm	10–25 cm	0–10 cm	10–25 cm
Species (SP)				
Blue-spirea	3.0 h ^y	1.4 f	5.7 e	5.5 b
Border forsythia	17.5 bc	14.8 a	8.2 d–e	5.6 b
Boxleaf honeysuckle	12.2 b–f	5.2 b–e	14.2 a–e	7.0 b
Bumald spirea	11.2 c–g	5.3 b–e	16.8 a–e	7.7 b
Bush cinquefoil	7.8 e–h	4.3 c–f	11.7 b–e	11.2 ab
Buttercup winterhazel	9.4 d–h	7.9 bc	10.7 c–e	7.2 b
Common ninebark	12.3 b–f	6.0 b–d	11.3 c–e	6.2 b
Common seabuckthorn	5.8 f–h	3.1 d–f	12.2 b–e	5.5 b
Fuzzy deutzia	28.7 a	12.2 a	17.6 a–d	10.9 ab
Glossy abelia	10.5 d–g	4.1 c–f	13.7 b–e	6.7 b
Goldencup st. johnswort	13.0 b–e	4.3 c–f	16.3 a–e	9.5 ab
Hybrid deutzia	19.0 b	8.2 b	16.5 a–e	7.8 b
Japanese kerria	12.2 b–f	5.4 b–e	8.3 d–e	8.1 b
Privet honeysuckle	7.0 e–h	4.2 c–f	24.8 a	9.2 ab
Purpleosier willow	5.0 gh	3.1 d–f	12.6 b–e	7.1 b
Redosier dogwood	8.8 d–h	5.6 b–e	9.6 d–e	7.1 b
Scorpion senna	4.1 gh	3.6 d–f	8.6 d–e	5.7 b
Shrub althea	13.9 b–e	7.5 bc	13.0 b–e	5.3 b
Slender deutzia	8.4 d–h	3.5 d–f	21.1 a–c	15.1 a
Spanish broom	4.4 gh	3.0 d–f	8.5 d–e	5.0 b
Spanish gorse	4.6 gh	2.1 ef	22.6 ab	9.6 ab
Virginal mock orange	14.9 b–d	8.1 b	22.7 ab	10.3 ab
Weed management (WM)				
Polypropylene fabric	6.5 c	3.9 b	10.3 b	6.8 b
Biodegradable textile	10.7 b	6.2 a	12.0 b	7.4 b
Bare soil	14.6 a	6.8 a	19.6 a	9.4 a
Probability				
SP	***	***	***	*
WM	***	***	***	*
SP \times WM	***	NS	NS	NS

^z1 $m \cdot L^{-1}$ = 12.4193 ft/gal, 1 $m \cdot g^{-1}$ = 93.0102 ft/oz.

^yMeans for each main factor and within each column followed by the same letter are not significantly different ($P \leq 0.05$) according to Duncan's multiple range test.

^{NS,***}Non-significant or significant at $P \leq 0.001$, respectively.

while blue-spirea had the lowest. In the deeper soil zone (between 10 and 25 cm), slender deutzia showed the highest specific root length; bush cinquefoil, fuzzy deutzia, goldencup st. johnswort, privet honeysuckle, spanish gorse, and virginal mock orange did not show significant differences from slender deutzia for this parameter (Table 7). As reported in Table 7, in the shallower soil zone an interaction between species and soil management was detected for root density. When significant differences were found, the highest root density was observed in plant grown in bare soil, whereas the lowest values were reported in plants mulched with polypropylene fabric, with the exception of fuzzy deutzia. In this species, the highest root density was observed in plants mulched with biodegradable textile (Table 8). In the deeper soil zone (between 10 and 25 cm), the greater root density was observed in plants grown using biodegradable mulching and bare soil, whereas the lowest values were observed in plants mulched with polypropylene fabric (Table 7). Border forsythia and fuzzy deutzia showed the highest root density values at this depth, whereas the lowest values were detected in blue-spirea (Table 7). No significant differences in root density from blue-spirea were reported for bush cinquefoil, common seabuckthorn, glossy abelia, goldencup st. johnswort, privet honeysuckle, purpleosier willow, scorpion senna, slender deutzia, spanish broom, and spanish gorse at the 10–25-cm depth (Table 7). The higher root density and the higher specific root length detected in plants grown in bare soil can be probably explained by the necessity of the plant to explore the soil to reach for water. As reported by several authors mulched soil has greater water retention than bare soil (Chalker-Scott, 2007).

The type of soil management did not affect plant phenological stages and plant susceptibility to diseases or pests (data not shown). Septoria (*Septoria cornicola*) caused severe leaf damage on redosier dogwood, leading to early leaf fall. Aphids were detected on young shoots of goldencup st. johnswort, japanese kerria, scorpion senna, shrub althea, spanish broom, and virginal mock orange, but the damage was cosmetic.

Results of this study showed that mulching increased shoot biomass in

Table 8. Root density (root length/soil volume), recorded in the first 10-cm (3.9 inches) soil depth, in 25 shrubs grown on a slope under different types of soil management (bare soil and mulched with biodegradable textile or polypropylene fabric) at the end of the third growing season.

Species	Root density (m·L ⁻¹) up to 10-cm soil depth ^z				Probability
	Polypropylene fabric	Biodegradable textile	Bare soil		
Blue-spirea	2.4 a ^y	1.7 a	4.9 a		NS
Border forsythia	17.4 a	7.3 a	27.9 a		NS
Boxleaf honeysuckle	12.0 a	13.4 a	11.2 a		NS
Bumald spirea	11.1 a	6.8 a	15.6 a		NS
Bush cinquefoil	4.4 a	9.1 a	9.7 a		NS
Buttercup winterhazel	8.3 a	10.1 a	10.0 a		NS
Common ninebark	4.2 a	13.6 a	19.0 a		NS
Common seabuckthorn	3.0 a	8.2 a	6.2 a		NS
Fuzzy deutzia	11.9 b	45.5 a	28.8 ab		*
Glossy abelia	6.3 a	8.8 a	16.4 a		NS
Goldencup st. johnswort	4.6 b	11.1 b	23.5 a		**
Hybrid deutzia	8.9 b	19.7 ab	28.5 a		**
Japanese kerria	4.6 a	18.9 a	13.2 a		NS
Privet honeysuckle	6.9 a	5.7 a	8.3 a		NS
Purpleosier willow	3.7 a	3.7 a	7.7 a		NS
Redosier dogwood	4.9 b	5.9 b	15.4 a		**
Scorpion senna	2.1 b	3.5 ab	6.8 a		*
Shrub althea	12.0 a	11.8 a	17.8 a		NS
Slender deutzia	2.5 a	7.7 a	14.8 a		NS
Spanish broom	4.6 a	3.4 a	5.3 a		NS
Spanish gorse	1.4 b	5.2 ab	7.3 a		*
Virginal mock orange	5.7 b	15.2 ab	23.9 a		*

^z1 m·L⁻¹ = 12.4193 ft/gal.

^yMeans within the same row followed by different letters are significantly different from each other using the Duncan's mean separation test.

NS, *, **, *** indicate nonsignificant, significant at $P \leq 0.05$, 0.01, or 0.001, respectively.

some species and reduced hand weeding time. However considerations are needed about this practice, especially when used on a slope area. As previously reported, the use of mulching leads to a lower root density and lower specific root length compared with bare soil management levels, involving higher management and associated costs (materials, setting an higher transplanting labor). Species or cultivars characterized by fast groundcover ability should be preferred in slope areas to limit the unfavorable effect of the weeds, especially in bare soil. Among the species tested, fuzzy deutzia and virginal mock orange showed a higher number of desirable qualities: fast groundcovering ability, reduced hand weeding time, plant shoot biomass unaffected by soil management and high root density and high specific root length. They should be considered as desirable species for the

slopes along roads and highways in similar conditions to those of the experimental area.

Literature cited

Angelopoulos, K., B. Dichio, and C. Xiloyannis. 1996. Inhibition of photosynthesis in olive trees (*Olea europaea* L.) during water stress and rewatering. *J. Expt. Bot.* 47:1093–1100.

Assone, S., P. Borsotto, E. Fiora, and S. Faccioli Cerea. 2000. Sperimentazione relativa alle principali specie tappezzanti in vista di un possibile impiego nel verde urbano, p. 99–114. *Proc. Natl. Congr. Il verde urbano, storico e contemporaneo: Problematiche legate alla gestione*. Brusson, Italy.

Burroughs, R. 2010. Water-wise landscaping. *Amer. Nurseryman* 210(7):10–20.

Cazzuffi, D. and F. Tironi. 2003. Contribution of roots to slope stability: An overview of typical results for different

plants, p. 101–105. *Proc. Intl. Conf. Fast slope movements: Prediction and prevention for risk mitigation*. Naples, Italy.

Chalker-Scott, L. 2007. Impact of mulches on landscape plants and the environment—A review. *J. Environ. Hort.* 25:239–249.

Einert, A.E., R. Guidry, and H. Huneycutt. 1975. Permanent mulches for landscaping plantings of dwarf crape myrtles. *Amer. Nurseryman* 142(9):62–65.

Ferrini, F., A. Fini, P. Frangi, and G. Amoroso. 2008. Mulching of ornamental trees: Effects on growth and physiology. *Arboriculture Urban For.* 34:157–162.

Ferrini, F., A. Fini, S. Pellegrini, A. Agnelli, M. Platinetti, P. Frangi, and G. Amoroso. 2009. Effect of two organic mulches on soil physical, chemical and biological properties, p. 31–39. In: G. Watson, L. Costello, B. Scharenbroch, and E.F. Gilman (eds.). *The landscape below ground III*. Intl. Soc. Arboriculture, Champaign, IL.

Franco, J.A., J.J. Martinez-Sanchez, J.A. Fernandez, and S. Bañon. 2006. Selection and nursery production of ornamental plants for landscaping and xerogardening in semiarid environment. *J. Hort. Sci. Biotechnol.* 81:3–17.

Ganatsas, P.P. and M.N. Tsakalidimi. 2003. Root system modification of *Pinus brutia* Ten. species under adverse ecological conditions. *Proc. 8th Intl. Conf. Environ. Sci. Technol.* 216–223.

Gray, D.H. and A.T. Leiser. 1982. *Biotechnical slope protection and erosion control*. Van Nostrand Reinhold, New York.

Gray, D.H. and R.B. Satir. 1996. *Biotechnical and soil bioengineering slope stabilization: A practical guide for erosion control*. Wiley-Interscience, New York.

Harris, R.W., J.R. Clark, and N.P. Matheny. 2004. *Arboriculture: Integrated management of landscape trees, shrubs, and vines*. 4th ed. Prentice Hall, Upper Saddle River, NJ.

Lazár, D. 2006. The polyphasic chlorophyll a fluorescence rise measured under high intensity of exciting light. *Funct. Plant Biol.* 33:9–30.

Long, C.E., B.L. Thorne, N.L. Breisch, and L.W. Douglas. 2001. Effect of organic and inorganic landscape mulches on subterranean termite (Isoptera: Rhinotermitidae) foraging activity. *Environ. Entomol.* 30:832–836.

Macdonald, K.B. and B. Witek. 1994. Management options for unstable bluffs in Puget Sound, Washington. Coastal erosion management studies. Vol. 8. Shorelands and Water Resources Program, Washington Department of Ecology, Olympia.

- Maxwell, K. and G.N. Johnson. 2000. Chlorophyll fluorescence—A practical guide. *J. Expt. Bot.* 51:659–668.
- Morgan, R.P.C. and R.J. Rickson. 1994. Slope stabilization and erosion control: A bioengineering approach. E & FN Spon, London.
- Osmond, D.L., C.R. Crozier, and D.H. Hardy. 1997. Careful Soil Sampling—The Key to Reliable Soil Test Information. 18 Apr. 2011. <<http://www.soil.ncsu.edu/publications/Soilfacts/AG-439-30/>>.
- Percival, G.C. and G.A. Fraser. 2001. Measurement of the salinity and freezing tolerance of *Crataegus* genotypes using chlorophyll fluorescence. *J. Arboriculture* 27:233–245.
- Percival, G.C., I.P. Keary, and K. Novis. 2008. The potential of a chlorophyll content SPAD meter to quantify nutrient stress in foliar tissue of sycamore (*Acer pseudo-platanus*), english oak (*Quercus robur*), and european beech (*Fagus sylvatica*). *Arboriculture Urban For.* 34:89–100.
- Percival, G.C., I.P. Keary, and S. Al-Habsi. 2006. An assessment of the drought tolerance of *Fraxinus* genotype for urban landscape planting. *Urban For. Urban Green.* 5:17–27.
- Piniór, A., G. Grunewaldt-Stöcker, H. von Alten, and R.J. Strasser. 2005. Mycorrhizal impact on drought stress tolerance of rose plants probed by chlorophyll a fluorescence, proline content and visual scoring. *Mycorrhiza* 15:596–605.
- Preti, F. and F. Giadrossich. 2009. Root reinforcement and slope bioengineering stabilization by spanish broom (*Spartium junceum* L.). *Hydrol. Earth Syst. Sci.* 13:1713–1726.
- Rice, Jr., R.P. 1992. Nursery and landscape weed control manual. Thomson Publications, Fresno, CA.
- Stinson, J.M., G.H. Brinen, D.B. McConnell, and R.J. Black. 1990. Evaluation of landscape mulches. *Proc. Florida State Hort. Soc.* 103:372–377.
- Stokes, A., C. Atger, A.G. Bengough, T. Fourcaud, and R.C. Sidle. 2009. Desirable plant root traits for protecting natural and engineered slopes against landslides. *Plant Soil* 324:1–30.
- Strasser, R.J., A. Srivastava, and M. Tsimilli-Michael. 2000. The fluorescence transient as a tool to characterize and screen photosynthetic samples, p. 445–483. In: M. Yunus (ed.). *Probing photosynthesis: Mechanisms, regulation and adaptation*. Taylor & Francis, London.
- Strasser, R.J. and M. Tsimilli-Michael. 2001. Stress in plants from daily rhythm to global changes, detected and quantified by JIP test. *Chimie Nouvelle (SRC)* 75: 3321–3326.
- Tennant, D. 1975. A test of a modified line intersect method of estimating root length. *J. Ecol.* 63:995–1001.
- Tóth, S.Z., G. Schansker, and R.J. Strasser. 2007. A non-invasive assay of the plastoquinone pool redox state based on the OJIP-transient. *Photosyn. Res.* 93: 193–203.
- Tosi, M. 2007. Root tensile strength relationships and their slope stability implications of three shrub species in the northern Apennines (Italy). *Geomorphology* 87:268–283.
- Yamada, M., D. Hidaka, and H. Fukamachi. 1996. Heat tolerance in leaves of tropical fruit crops as measured by chlorophyll fluorescence. *Scientia Hort.* 67:39–48.
- Yeager, T., T. Bilderback, D. Fare, C.H. Gilliam, J. Lea-Cox, A. Niemiera, J. Ruter, K.M. Tilt, S.L. Warren, T. Whitwell, and R. Wright. 2006. Best management practices: Guide for producing nursery crops. 2nd ed. Southern Nursery Assn., Atlanta.