

Downloaded from <https://prime-pdfr-watemark.prime-prod.pubfactory.com/> at 2025-11-28 via Open Access. This is an open access article distributed under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). <https://creativecommons.org/licenses/by-nc-nd/4.0/>

NUMBER 5

Postmaster. Please return undeliverable copies and change of address "Form 3579" to the American Society for Horticultural Science, Mount Vernon, Virginia 22121, USA. Return postage guaranteed.

447

removed (defoliated branches); and b) about 8 leaves left on the branch (foliated branches). Fruits were then sealed in gas tight polyethylene bags provided with an inlet and outlet for air with the aid of Apiezon Q (a sealing compound). Some detached fruits were similarly placed in bags as controls. Half the number in each treatment was placed in a dark growth chamber and the remainder in another chamber scheduled for 12 hr of light each day. Both CO_2 and ethylene were monitored daily by injecting a 1 ml aliquot of the outflowing air into a Beckman GC 4 gas chromatograph equipped with Porapak Q column. CO_2 was determined with a thermal conductivity detector and C_2H_4 with a flame ionization system.

Ripening of detached fruits. Avocado fruits were harvested with a 6 cm peduncle (including both pedicel and the true peduncle), attached. The peduncles were trimmed under water, and a #20 serum bottle cap with a hole in the top was fitted over the peduncle and placed in a test tube containing water. Another group of fruits with peduncles cut off close to the fruit were divided into 2 lots: in 1 the peduncle scars were sealed with vaseline and in the other they were left untreated. Fruits were placed in 1.8 liter jars and ventilated with humidified air at 70 ml/min. The effluent air was sampled automatically every 4 hours and passed through a Beckman 215A infrared analyzer for CO_2 determination, and thence to an Aerograph HY-FI Model 600-D gas chromatograph for C_2H_4 determination. Operation of the gas chromatograph was made automatic through the use of an electrically activated sample loop and a Varian Model 480 digital integrator to record the output. The integrator was set to reject peaks of less than 100 counts, which was equivalent to 13 ppb or a production rate of $0.19 \mu\text{l/kg-hr}$. This rejection level was always more than 4 times an average noise peak.

The rate of CO_2 and C_2H_4 evolution from different parts of 'Hass' fruits with and without stems were determined as shown in Fig. 1. The caps of the bottles were fitted with inlet and outlet tubes so that air would be passed continuously over a definite area of the fruit surface. Carbon dioxide and C_2H_4 production were determined from these areas and from separated peduncles by the system described above.

Results

Ripening in the field. Fruit on girdled branches always abscised before any softening could be detected. Once abscission had occurred, prompt ripening was initiated, just as though mechanically detached, which occurred after 5 days in the field. Days from girdling to abscission was greater by 5 days when the girdle was proximal to a flush of leaves (Table 1) than when girdled distal to the nearest leaves. Removal of the leaves when a proximal girdle was made, caused a delay of 2 additional days in the average time to abscission. Clearly, nearby leaves do not delay abscission and ripening. In fact, our results show a slight acceleration of time to abscission when leaves are left attached. It appears that the length of the stem distal to the girdle is the primary factor which regulates the time to abscission.




Fig. 1. Sketch of set-up used for determining the rate of evolution of CO_2 and C_2H_4 from various parts of 'Hass' avocado fruits. The 120 ml polypropylene bottles whose bottoms have been cut away were fixed to the fruits with Terostat IX (Teroson-Werke GmbH, Hiedelberg, Germany).

Ripening on severed branches. Just as in the field studies, ripening was not initiated on fruit attached to branches in the growth chamber, but only occurred after abscission. Here CO_2 and C_2H_4 determinations made on the fruit (Fig. 2) show that the rise in both CO_2 and C_2H_4 occurred just after the fruit abscised. Thus the failure of movement of some inhibitor to the fruit or of a hormone from the fruit to stem must be the signal which initiates ripening. In the branch experiments, fruit on foliated branches always abscised and ripened earlier than fruit on branches from which the leaves were removed. This was especially evident when branches were kept in the dark. The light-dark cycle did not appear to affect the amount of CO_2 and C_2H_4 production but only accelerated the time to abscission.

Ripening of harvested fruits. When compared to detached fruit without peduncles, the climacteric for fruits with peduncles was

Table 1. Effect of girdling and defoliation of branches and detaching fruit on abscission and ripening of 'Hass' avocado fruit in May 1972. Average temperatures in the field were: 25.5°C mean maximum and 16°C mean minimum. Fruits in the laboratory were kept at 20°C .

Treatment	Av. days to abscission	Av. days to soften
Proximal girdling defoliated	23	28
Proximal girdling foliated	21	26
Distal girdling	16	21
Detached fruit in field		5
Detached fruit in lab		7




Fig. 2. Respiration pattern (a) and (c), ethylene production (b) and (d) of 'Fuerte' fruits in detached form Δ , on foliated branch \circ , and on defoliated branch \bullet ; at $20 \pm 2^\circ\text{C}$. Arrows indicate the time at which the fruits abscised.




Fig. 3. Respiration pattern of 'Hass' avocado fruits with no peduncle (\bullet), peduncle scar covered with vaseline (Δ) and with 6 cm long peduncle (\circ) 20°C . Each point represents average of 8 fruits and arrow indicates the time the peduncle separated.




Fig. 4. Respiration rate (a) and (c) and ethylene evolution (b) and (d) from different parts of 'Hass' avocado fruits at 20°C. Symbols represent: ● = stem end, Δ = midsection and ○ = stylar end.

delayed by at least one day; but there was no difference in onset of the climacteric between fruits whose stem scar was sealed with vaseline and where the peduncle was cut off as close as possible (Fig. 3). Peduncles separated from the fruit at an early stage of the climacteric rise.

When the rates of evolution of CO_2 and C_2H_4 from the stem end, stylar end and midsection of the fruits were determined per unit area, (Fig. 4), ethylene was always detected first at the stem end of the fruit; this preceded its detection from other parts of the fruits by as much as 8 to 24 hrs. As the climacteric progressed, the rates of evolution of both CO_2 and C_2H_4 from the stylar end and midsection exceeded rates from the stem end. The presence of 6 cm long peduncles delayed onset of the climacteric by 2 days, and also caused reduced C_2H_4 production from all parts of the fruit. No C_2H_4 production could be detected from 2 g of separated peduncles, which indicated that the production rate was less than $0.19 \mu\text{l/kg-hr}$.

Discussion

The results show that abscission and subsequent ripening of attached avocado fruits was not delayed by the presence or proximity of the leaves (Table 1 and Fig. 2), and therefore, do not support the hypothesis that leaves are a source of ripening inhibitor (5, 15). On the contrary, ripening was hastened when a few leaves were left on the severed branches (Fig. 2c and d) causing these fruits to ripen as if they were detached. Although water loss was reduced to a minimum, the negative effect of leaves may partially be due to greater water loss from these branches and, consequently, from the fruits. Littmann (9) and recently, Adato and Gazit (1) have shown that increased water loss hastened ripening of avocado fruits. The important issue is, however, that leaves do not contribute to extending the time to abscission and ripening over that provided by the peduncle and branch alone.

Experiments of Burg and Burg (5) focused attention on the leaves as the source of ripening inhibitors. It is, however, equally possible that the peduncle and stem may act as a sink for a ripening hormone produced in the fruit, as was suggested earlier by Biale (2). At the present, we cannot say whether substances controlling ripening are being moved into or from the fruit, or in fact whether movement of substances in both directions is necessary. We have been unsuccessful in making any extract of peduncles which, when vacuum infiltrated into preclimacteric fruit, exert any effect on ripening (14). This leaves open the possibility that the peduncle may act as a sink for hormones

produced in the fruit. On the other hand, we have shown (14) that low concentration of IAA ($1 \mu\text{M}$) infiltrated into preclimacteric avocado fruit does delay ripening and reduce the amount of C_2H_4 produced during the climacteric, much as does attachment of the peduncle. This observation suggests that the peduncle is supplying an auxin to the fruit and ripening is initiated only when the source of auxin is depleted and the level in the fruit is reduced to some critical value. Treatment with 1 mM IAA caused earlier and higher ethylene production compared to control fruit, which induced ripening earlier. Ethylene production in response to auxin treatment has been reported for a number of tissues (10, 12).

Ethylene production and softening of the fruit was always detected earlier in the stem end than in other parts of the fruit. This was not due to production of C_2H_4 by the peduncle which we found to produce no C_2H_4 detectable in our flow system. Earlier ripening in the stem end may be due to the extensive vascular system transferring a ripening regulator more effectively to or from the tissue of this part of the fruit.

Light has been reported by Peacock (11) to accelerate ripening in banana fruit and far-red light to accelerate senescence in *Marchantia polymorpha* (6). While our results (Fig. 2) show some acceleration of ripening on alternating 12 hr light-dark cycles, we were (as was Peacock) unable to keep the fruit tissue under light at exactly the same temperature as fruit held in the dark. We believe that the slightly more rapid ripening in light was mainly a temperature effect. We have not been able to demonstrate any effect of red or far-red light on ripening.

We conclude that attachment of avocado fruit to a branch or even the peduncle, delays abscission and reduces ethylene production during the climacteric. Ripening occurs only after the fruit has abscised from the peduncle. Nearby leaves on the branch either have no effect or actually accelerate the time to abscission and subsequent ripening. We believe that an auxin is at least one factor involved in the inhibition of ripening of attached avocado fruit.

Literature Cited

- Adato, I., and S. Gazit. 1974. Water-deficit stress, ethylene production, and ripening of avocado fruits. *Plant Physiol.* 53:45-46.
- Biale, J. B. 1960. Respiration of fruits. *Handbuch der Pflanzenphysiologie* 12:536-592.
- Burg, S. P. 1962. Postharvest ripening of avocados. *Nature* 194:398-399.
- . 1964. Studies on formation and function of ethylene gas in plant tissue. In: *Régulateurs Naturels de la Croissance Végétale* Ed. J. P. Nitsch pp. 719-725. C.N.R.S. Paris.
- , and E. A. Burg. 1964. Evidence of a naturally occurring inhibitor of ripening. *Plant Physiol.* 39: (suppl) x.
- De Greif, J. A. and H. Fredericq. 1972. Enhancement of senescence by far-red light. *Planta* 104:272-274.
- Gazit, S., and A. Blumenfeld. 1970. Response of mature avocado fruits to ethylene treatment before and after harvest. *J. Amer. Soc. Hort. Sci.* 95:229-231.
- Hansen, E. 1966. Postharvest physiology of fruits. *Ann. Rev. Plant Physiol.* 17:459-480.
- Littmann, M. D. 1972. Effect of water loss on the ripening of climacteric fruits. *Queensl. J. Agr. and Animal Sci.* 29:103-113.
- Morgan, P. W., and W. C. Hall. 1964. Accelerated release of ethylene by cotton following application of indolyl-3-acetic acid. *Nature* 201:99.
- Peacock, B. C. 1972. Effect of light on initiation of fruit ripening. *Nature New Biol.* 235:62-63.
- Pratt, H. K., and J. D. Goeschl. 1969. Physiological roles of ethylene in plants. *Ann. Rev. Plant Physiol.* 20:541-584.
- Smock, R. M. 1972. Influence of detachment from tree on the respiration of apples. *J. Amer. Soc. Hort. Sci.* 94:509-511.
- Tingwa, P. O. and R. E. Young. 1975. The effect of indole-3-acetic acid and other growth regulators on the ripening of avocado fruits. *Plant Physiol.* 55:937-940.
- Young, R. E. 1968. Regulation of respiration associated with slicing of plant tissue. Proc. Int. Symp. on Plant Biochem. Reg. in viral and other diseases or injury. *Phytopath. Soc. Japan*. Tokyo.