

14. Metzger, B. 1972. Cyclic lighting of carnations. *Colo. Flower. Gro. Assoc. Bul.* 265:1-4.
15. Novovesky, I.M.P. 1967. Effects of photoperiod and CO₂ enrichment on carnations. *Colo. Flower. Gro. Assoc. Bul.* 209:1-2.
16. Phillips, D. J. 1968. Growth and development of a carnation shoot. *Colo. Flower. Gro. Assoc. Bul.* 219:1-4.
17. Porkorny, F. A. and J. R. Kamp. 1960. Photoperiodic control of growth and flowering of carnation. *Ill. State Flor. Bul.* 202:6-8.
18. Tucker, D. J. 1975. Far-red light as a suppressor of side shoot growth in the tomato. *Plant Sci. Lett.* 5:127-130.
19. Vince, D., J. Tucker, and R. Spencer. 1964. Some effects of wavelengths of the supplementary light on the photoperiodic behavior of the long-day plants, carnation and lettuce. *Physiol. Plant* 17:119-125.

J. Amer. Soc. Hort. Sci. 104(3):319-322. 1979.

Boron Tolerance of Twenty-five Ornamental Shrub Species¹

L. E. Francois and R. A. Clark²

U. S. Salinity Laboratory, U. S. Department of Agriculture, Riverside, CA 92501

Additional index words. *Carissa grandiflora*, *Raphiolepis indica*, *Hibiscus rosa-sinensis*, *Nerium oleander*, *Buxus microphylla*, *Callistemon citrinus*, *Leucophyllum frutescens*, *Cordyline indivisa*, *Syzygium paniculatum*, *Podocarpus macrophyllus*, *Platyclusus orientalis*, *Rosmarinus officinalis*, *Abelia X grandiflora*, *Lantana camara*, *Juniperus chinensis*, *Ilex cornuta*, *Pittosporum tobira*, *Euonymus japonica*, *Feijoa sellowiana*, *Ligustrum japonicum*, *Viburnum tinus*, *Elaeagnus pungens*, *Xylosma congestum*, *Photinia X Fraseri*, *Mahonia aquifolium*.

Abstract. Boron (B) tolerance of 25 ornamental shrub species was determined in large, outdoor sand cultures. Tolerant species such as Natal plum (*Carissa grandiflora* (E. H. Mey.) A. D. C.), Indian hawthorn (*Raphiolepis indica* (L.) Lindl.), Chinese hibiscus (*Hibiscus rosa-sinensis* L.), oleander (*Nerium oleander* L.), Japanese boxwood (*Buxus microphylla* Siebold and Zucc.), bottlebrush (*Callistemon citrinus* (Curtis) Stapf), ceniza (*Leucophyllum frutescens* (Berland.) I. M. Johnst.), and blue dracaena (*Cordyline indivisa* (G. Forst) Steud.) were affected little, if at all, by 7.5 mg B/liter in the irrigation water. Sensitive species like yellow sage (*Lantana camara* L.), juniper (*Juniperus chinensis* L.), Chinese holly (*Ilex cornuta* Lindl. and Paxt.), Wax-leaf privet (*Ligustrum japonicum* Thunb.), laurustinus (*Viburnum tinus* L.), thorny elaeagnus (*Elaeagnus pungens* Thunb.), xylosma (*Xylosma congestum* (Lour.) Merrill), photinia (*Photinia X Fraseri* Dress.), and Oregon grape (*Mahonia aquifolium* (Pursh) Nutt.) were severely damaged or killed by 7.5 mg B/liter and moderately damaged by 2.5 mg B/liter in the irrigation water. B tolerance and B accumulation in the leaves were not correlated and no correlation was found between B tolerance and salinity tolerance.

Boron (B) toxicity of plants in the United States is primarily confined to a few irrigated areas in the West and secondarily to over fertilization with B fertilizer. Although excess B problems do not involve a large area, plant injury can be severe. The most common source of B in soils is from irrigation water pumped from wells with high B contents and, to a lesser extent, from very shallow water tables and soils naturally high in B. Most irrigation waters drawn from surface streams are low in B (5).

The B tolerance of a wide variety of plants has been tested over the years including major crop plants (4, 6, 8, 12, 13) and native species (3, 11). However, information on B tolerance of ornamentals, is severely lacking and especially needed where urban development is taking place on land that was previously irrigated with water containing B. This study was conducted to determine the B tolerance of 25 ornamental species often used in landscaping where excess B is a problem.

Methods and Materials

Twenty-five shrub species (Table 1), growing in 4-liter containers, were purchased from a commercial nursery, and transplanted on March 7, 1973, into 15 large outdoor sand tanks. Before transplanting, the soil was washed from the roots. Five tanks were used for each B treatment and each species was replicated twice within a treatment by planting 10 species in each tank. The tanks were 4.3 m long × 0.76 m wide × 0.76 m

deep, filled with a coarse river sand. Each sand tank had a 2400-liter irrigation reservoir in addition to 400 liters of irrigation water retained in the sand. Culture solutions were surface applied automatically twice daily to completely flush the sand tanks and were returned to the reservoir. Boron treatments, 0.5, 2.5, and 7.5 mg B/liter, were added as H₃BO₃ to tap water on April 23, 1973. These levels included 0.1 mg B/liter as impurities in the nutrient salts and 0.06 mg B/liter in tap water. Boron levels were monitored monthly to maintain desired levels. The study was terminated on January 20, 1976 (33 months later).

The base nutrient solution contained 2, 1.5, 1.0, 0.5 mm of Ca(NO₃)₂, KCl, MgSO₄ · 7 H₂O, and NH₄H₂PO₄, respectively, and 0.5 mg/liter Fe as iron citrate, 0.25 mg/liter Mn as MnCl₂, 0.025 mg/liter Zn as ZnSO₄, 0.01 mg/liter Cu as CuSO₄, and 0.005 mg/liter Mo as molybdic acid. Tap water was added to the irrigation reservoirs about every 3 weeks to replenish water lost by evapotranspiration. Nutrient salts were also added in proportion to the amount of water replenished. Sulfuric acid was added to lower the pH to the range of 6 to 7. Culture solutions were changed completely once each year in midwinter.

All shrubs were pruned during the winter, or more frequently as required. Pruning weights were added to top weights to determine total top growth (uncorrected for initial top weight). Appearance assessment was made throughout the experiment, because aesthetic value is often more important than growth rate for ornamentals.

Leaf samples for B analysis were collected annually in October and when injury symptoms developed. Injured and uninjured leaves of comparable age were sampled from the same plant whenever possible. Boron was determined colorimetrically by the Carmine method (9).

¹Received for publication November 6, 1978.

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper must therefore be hereby marked *advertisement* solely to indicate this fact.

²Research Agronomist and Chemist, respectively.

Table 1. Leaf symptoms of B injury and growth reduction of 25 shrub species at 2.5 and 7.5 mg B/liter compared with the control treatment of 0.5 mg B/liter. (LB = low concentration boron; HB = high concentration).

Species ^z	General observations	Growth reduction (%)
<i>Tolerant</i>		
Natal plum (<i>Carissa grandiflora</i> (E. H. Mey.) A. DC. cv. Tuttlei)	LB: No injury HB: No injury	0 0
Indian hawthorn (<i>Raphiolepis indica</i> (L.) Lindl. cv. Enchantress)	LB: No injury HB: No injury	0 0
Chinese hibiscus (<i>Hibiscus rosa-sinensis</i> L.)	LB: No injury HB: Slight premature leaf drop	0 0
Oleander (<i>Nerium oleander</i> L.)	LB: No injury HB: Narrow (1 to 2 mm) marginal chlorosis-slight tip burn	21 24
Japanese boxwood (<i>Buxus microphylla</i> Siebold and Zucc. var. <i>japonica</i> (Mull. Arg.))	LB: No injury HB: General marginal chlorosis with necrotic older leaves	0 0
Bottlebrush (<i>Callistemon citrinus</i> (Curtis) Stapf)	LB: Slight marginal coloration similar to HB HB: Marginal anthocyanin coloration (5 mm from leaf tip) progressed inward in semicircle pattern toward midrib; marginal and tip necrosis developed as leaves matured	0 0
Ceniza (<i>Leucophyllum frutescens</i> (Berland.) I. M. Johnst. cv. Compactum)	LB: No injury HB: Older leaves dropped prematurely	15 24
Blue dracaena (<i>Cordyline indivisa</i> (G. Forst) Steud.)	LB: Tip burn, 5 to 7 cm (1973) 10 to 13 cm (1975) HB: Tip burn, 7 to 10 cm (1973) 18 to 22 cm (1975)	0 0 0 0
<i>Semi-tolerant</i>		
Brush cherry (<i>Syzygium paniculatum</i> Gaertn.)	LB: Slight anthocyanin spotting oldest leaves HB: Moderate anthocyanin spotting; oldest leaves dropped prematurely; general appearance chlorotic	0 11
Southern yew (<i>Podocarpus macrophyllus</i> (Thunb.) D. Don var. <i>Maki</i> Endl)	LB: Slight tip burn with narrow chlorotic band between burn and remainder of leaf HB: Moderate to severe tip burn (1 cm) with narrow chlorotic area like LB – leaves on lower ¼ plant exhibited burn	0 8
Oriental arborvitae (<i>Platycladus orientalis</i> (L.) Franco)	LB: Slight chlorosis to necrosis on tips older leaves HB: Severe necrosis older leaves – only outside perimeter of plant was still green	27 30
Rosemary (<i>Rosmarinus officinalis</i> L.)	LB: Tip necrosis older leaves HB: Tip necrosis all leaves	20 51
Glossy abelia (<i>Abelia × grandiflora</i> (Andre) Rehd.)	LB: Bronzing and tip burn older leaves HB: Bronzing all leaves, slight leaf drop	56 70
<i>Sensitive</i>		
Yellow sage (<i>Lantana camara</i> L.)	LB: Tip and marginal leaf burn intermediate and older leaves, some hastened leaf drop HB: Moderate to severe leaf burn all leaves, severe leaf drop	14 82
Juniper (<i>Juniperus chinensis</i> L. cv. <i>Armstrongii</i>)	LB: Moderate tip burn older leaves HB: Severe tip burn all leaves except outside perimeter of new leaves – center leaves of plant dead	20 47
Chinese holly (<i>Ilex cornuta</i> Lindl. and Paxt. cv. <i>Burfordii</i>)	LB: Some marginal burn and interveinal chlorosis HB: Tip and marginal burn all leaves – premature leaf drop	17 88
Japanese pittosporum (<i>Pittosporum tobira</i> (Thunb.) Ait.)	LB: Marginal and tip burn distal half older leaves – premature leaf drop HB: Premature leaf drop all leaves except very youngest – young leaves chlorotic with moderate to severe marginal and tip burn – small rosettes young leaves at branch tips	50 50

(Continued)

Table 1. (Continued)

Species ^Z	General observations	Growth reduction (%)
Spindle tree (<i>Euonymus japonica</i> Thunb. cv. <i>Grandifolia</i>)	LB: Slight tip burn, slight leaf drop HB: Severe chlorosis and tip burn all leaves, severe leaf drop	4 100
Pineapple guava (<i>Feijoa sellowiana</i> O. Berg.)	LB: Slight tip burn 1st year – moderate leaf drop, moderate tip and marginal burn 1974 and 1975 HB: Severe leaf drop – all leaves showed severe tip and marginal burn – youngest leaves also chlorotic	13 35
Wax-leaf privet (<i>Ligustrum japonicum</i> Thunb.)	LB: No apparent injury symptoms except reduced growth HB: Terminal 1/2 to 2/3 of branches dead – necrotic spotting older leaves, nearly completely defoliated	17 100
Laurustinus (<i>Viburnum tinus</i> L. cv. <i>Robustum</i>)	LB: Marginal and tip burn intermediate and older leaves, moderate leaf drop HB: Severe tip and marginal burn, all leaves, except very youngest	0 100
Thorny elaeagnus (<i>Elaeagnus pungens</i> Thunb. cv. <i>Fruitlandii</i>)	LB: Older leaves interveinal and marginal chlorosis on distal half of leaf HB: Severe chlorosis with marginal necrosis, severe leaf drop nearly all but youngest leaves, remaining leaves hyponastic	11 70
Xylosma (<i>Xylosma congestum</i> (Lour.) Merrill)	LB: Older leaves anthocyanin mottling and tip burn, more severe by mid-Summer, severe leaf drop older leaves HB: Many branches dead, anthocyanin mottling and severe tip burn all leaves, nearly complete leaf drop	23 100
Photinia (<i>Photinia</i> × <i>Fraseri</i> Dress)	LB: Marginal and tip burn older leaves HB: Severe leaf burn, severe leaf drop – stem tips dead, death mid-1974	0 100
Oregon grape (<i>Mahonia aquifolium</i> (Pursh) Nutt.)	LB: Tip necrosis young leaves, severe leaf drop older leaves HB: Severe leaf drop except very youngest – severe burn older and intermediate leaves, tip burn young leaves, barely survived 1st year (1973)	50 100

^ZSpecies listed in descending order of tolerance.

Results and Discussion

The B tolerance of the 25 shrub species is presented in Table 1 in decreasing order of tolerance and are classified as tolerant, semitolerant, or sensitive. There is no sharp line of demarcation between categories. These classifications are based on both growth reduction and overall shrub appearance. Some species, like oleander and ceniza, are classified as tolerant since their




Fig. 1. Leaf injury on Bottlebrush.

esthetic value is considered more important than reduction in growth. In general, the semitolerant and sensitive shrubs suffered in appearance, as well as growth reduction.

Evaluation of species response to B is given in Table 1 in terms of injury description and percent growth reduction. Early injurious effects of B were usually observed on the leaves as marginal chlorosis or necrosis, tip burn, cupping of leaves, necrotic spots between veins, reduced leaf size, premature leaf drop or development of anthocyanins. Similar injury symptoms have been reported for several plants (5, 10). These types of injury are reported as caused by excess B carried to the leaves in the transpirational stream, where it moves from the veins into the interveinal tissue and accumulates at the leaf tip and margins (1,6). Bottlebrush developed leaf injury which may be unique to that species. The initial visual injury symptom was the development of an anthocyanin semicircle on the leaf margin about 5 mm from the tip. As injury became more severe the semicircle increased in circumference with a chlorotic area between the anthocyanin on the margin and a very narrow band of anthocyanin between the chlorotic area and the healthy leaf tissue (Fig. 1). The leaves never became completely injured.

No correlation was found between B tolerance and B

Table 2. Leaf B concentrations of 25 shrub species at three nutrient B levels.

Species	Leaf B concentration (mg/kg dry weight)			B damaged leaves ^z
	0.5	2.5	7.5	
Natal plum	62	261	632	—
Indian hawthorn	57	94	164	440
Chinese hibiscus	53	67	140	—
Oleander	83	240	288	667
Japanese boxwood	106	416	646	1250
Bottlebrush	85	277	706	917
Blue dracaena	68	85	180	—
Ceniza	65	81	157	—
Brush cherry	108	389	550	1030
Southern yew	74	202	294	690
Oriental arborvitae	80	167	391	413
Rosemary	55	152	327	626
Glossy abelia	153	288	1074 ^y	1470
Yellow sage	92	286	399	716
Juniper	43	126	292	325
Chinese holly	109	426	674 ^x	994
Japanese pittosporum	63	234	669	742
Spindle tree	59	182	206	416
Wax-leaf privet	35	134	318	446
Pineapple guava	74	300	782	1020
Laurustinus	106	364	729	1212
Thorny elaeagnus	113	329	348	585
Xylosma	101	259	551	785
Photinia	90	144	254	1102
Oregon grape	128	347	618	994

^zLowest concentration B found in damaged leaves.

^xMarginal chlorosis.

^yDull bronzed and tip burn.

accumulation in leaves (Table 2). The highly tolerant Natal plum accumulated 261 and 632 B/kg dry weight at the 2.5 and 7.5 mg/liter B treatments, respectively, with no leaf injury nor growth reduction. In contrast, the spindle tree, wax-leaf privet and thorny elaeagnus showed considerable leaf burn and growth reduction at 416, 446, and 585 mg B/kg dry weight, respectively.

Leaves of the blue dracaena exhibited tip burn at all B levels. The leaf tips, when analyzed separately from the healthy leaf tissue, showed extremely high B accumulation. The control plants, with only 0.5 mg B/liter in the irrigation water, contained 8615 mg B/kg dry weight in the tips, whereas the remaining tissue contained only 68 mg B/kg dry weight. The 2.5 and 7.5 mg/liter B treatments showed similar results with 9960 and 11297 mg/B per kg in the tips, respectively. Eaton (5) reported similar findings when he analyzed the burned tips of eureka lemon (*Citrus limon* (L.) Burm. F.) leaves separately from the rest of the leaf.

The injury which develops is closely related to the type of leaf venation. The blue dracaena, rosemary, and southern yew, all with parallel venation, showed tip burn and no marginal injury. Shrubs with pinnate or palmate venation, however, exhibited both tip and marginal burn. Similar symptoms have been reported by Oertli and Kohl (10) for other species.

Since B accumulates predominantly in the leaf tissue, without later translocation (6), fall or winter pruning removes much of the accumulated B. Consequently, the new spring growth will contain low B levels.

Climatic conditions can play an important role in B tolerance, since the movement of B to the leaves is governed by transpiration. Under the hot (35°C), dry (30% relative humidity) summer conditions in Riverside, California, B will accumulate considerably faster than under a cool, wet climate. Injury may occur in the hot, dry climate but not in the cool, wet climate, although soil B concentrations are similar.

No correlation was found between B tolerance and the reported salt tolerance of these shrubs (2, 7). Yellow sage and juniper, both sensitive to B, are moderately tolerant to salinity; whereas Chinese hibiscus, which is tolerant to B, is very sensitive to salinity. Natal plum and Oleander are tolerant to both stress conditions, whereas Chinese holly and pineapple guava are sensitive to both conditions. While excess B may be associated with saline irrigation water, the B tolerance of individual species to such conditions are not necessarily related to salt tolerance.

Literature Cited

- Asen, S., and O. W. Davidson. 1950. The boron distribution in greenhouse rose plants. *Proc. Amer. Soc. Hort. Sci.* 56:433-438.
- Bernstein, L., L. E. Francois, and R. A. Clark. 1972. Salt tolerance of ornamental shrubs and ground covers. *J. Amer. Soc. Hort. Sci.* 97:550-556.
- Chatterton, N. J., C. M. McKell, J. R. Goodin, and F. T. Bingham. 1969. *Atriplex polycarpa*. II. Germination and growth in water cultures containing high levels of boron. *Agron. J.* 61:451-453.
- Cooper, W. C., and A. Peynado. 1959. Chloride and boron tolerance of young-line citrus trees on various rootstocks. *J. Rio Grande Valley Hort. Soc.* 13:89-96.
- Eaton, F. M. 1935. Boron in soil and irrigation waters and its effect on plants. *USDA Tech. Bul.* 448.
- _____. 1944. Deficiency, toxicity, and accumulation of boron in plants. *J. Agr. Res.* 69:237-277.
- Francois, L. E., and R. A. Clark. 1978. Salt tolerance of ornamental shrubs, trees, and iceplant. *J. Amer. Soc. Hort. Sci.* 103:280-283.
- Haas, A. R. C. 1929. Toxic effect of boron on fruit trees. *Bot. Gaz.* 88:113-131.
- Hatcher, J. T., and L. V. Wilcox. 1950. Colorimetric determination of boron using carmine. *Anal. Chem.* 22:567-569.
- Oertli, J. J., and H. C. Kohl. 1961. Some considerations about the tolerance of various plant species to excessive supplies of boron. *Soil Sci.* 92:243-247.
- Pratt, P. F., E. C. Nord, and F. L. Bair. 1971. Early growth tolerances of grasses, shrubs, and trees to boron in tunnel spoil. *USDA For. Serv. Res. Note* PSW-232.
- Purvis, E. R., and W. J. Hanna. 1938. Boron studies: I. The susceptibility of various plants to boron toxicity as influenced by soil type. *Proc. Soil Sci. Soc. Amer.* 3:205-209.
- Wilcox, L. V. 1960. Boron injury to plants. *USDA Info. Bul.* 211.