

Keeping Qualities of 'Stayman' and 'Delicious' Apples Treated with Calcium Chloride, Scald Inhibitors, and other Chemicals¹

R. E. Hardenburg and R. E. Anderson²

Horticultural Crops Quality Laboratory, Horticultural Science Institute, SEA-AR, U.S. Department of Agriculture, Beltsville, MD 20705

Additional index words. *Malus domestica*, storage, decay physiological diseases

Abstract. 'Stayman' apples (*Malus domestica* Borkh.) dipped in 4% CaCl₂ were firmer after 5-6 months storage in air at 0°C and developed less senescent breakdown than untreated fruit. Use of the thickener Keltrol with CaCl₂ or vacuum infiltrating (VI) CaCl₂ resulted in the firmest fruit after storage and the highest flesh Ca levels. 'Stayman' stored 5 or 6 months in CA at 0°C were 1.5 kg firmer than air-stored fruit. Dipping apples in CaCl₂ prior to CA storage provided little additional benefit. Liquid concentrate formulations of diphenylamine (DPA) and ethoxyquin used alone or in combination with 4% CaCl₂ gave excellent scald control on 'Stayman' stored 5 or 6 months at 0° + 6 days at 20°. However, the same chemicals gave poor scald control for 'Starkrimson Delicious' picked early to midseason. The antioxidants BHT and BHA evaluated as postharvest dips at 2,000 ppm were less effective than DPA or ethoxyquin in controlling scald on 'Stayman'. 'Delicious' apples were not significantly firmer after 6 months storage in air at 0°C when dipped in 3% CaCl₂ alone or with Keltrol, or when CaCl₂ was VI. When CaCl₂ was contaminated with *Penicillium expansum* spores, VI of the solution greatly increased decay of both 'Delicious' and 'Stayman'.

Bitter pit of apples has been reduced or controlled in numerous cultivars with CaCl₂ dips before storage or transport (3, 6, 7, 18, 20, 22, 24, 25, 27). CaCl₂ is also used commercially as dip or drench treatments to reduce storage breakdowns of 'Jonathan', 'Spartan', and 'Cox's Orange Pippin' apples (1, 4, 12, 16, 21), the incidence of breakdown being related to the final Ca concentration in the fruit (6, 17). Uptake of Ca solution by fruit was primarily through lenticels and increased with higher concentration of solution; absorbed Ca moved inward through tissues but remained highest near the surface (2, 8, 11, 12, 14). Use of various surfactants, thickeners such as Keltrol, and lecithin with CaCl₂ often has increased effectiveness of the treatment, delayed softening (3, 5-8, 10, 13, 14, 19), or reduced Ca injury (7, 18). For some cultivars both vacuum and pressure infiltration of CaCl₂ solution may improve control of breakdown and bitter pit over simple dipping (7, 15, 22, 23, 26).

The objective of our studies was to evaluate the post-storage qualities and Ca content of 'Delicious' and 'Stayman' apples when treated with CaCl₂ alone and in combination with scald inhibitors and other chemicals, and when applied by different methods after harvest.

Materials and Methods

'Delicious' apples from 4 orchards in Virginia, Maryland, and Pennsylvania were evaluated during the 1978-79 season, using 75 to 90 apples per treatment per orchard. 'Stayman' apples from 4 orchards also were evaluated during the 1978-79 season, using 90

to 115 apples per treatment per orchard. For another 'Stayman' test, fruit was obtained from 8 orchards, 4 in 1978-79 and 4 in 1979-80. All fruit were composited after picking. Initial fruit firmness averaged 7.1 to 7.3 kg for the 'Delicious' and 6.8 to 8.1 kg for the 'Stayman' fruit.

Laboratory grade USP CaCl₂ (76%), made up as 2, 3, or 4% (w/v) solutions, was used in most tests. In 1 test, laboratory grade CaCl₂ was compared with a commercial grade CaCl₂ (78% Dow flake) on 'Stayman' apple. The food thickener Keltrol (xanthan gum) used at 0.2 or 0.3% (w/v) to aid retention of CaCl₂ on apples was prepared in a Waring blender with water. In some tests fruit was stored without rinsing following treatment; in others it was rinsed after 2 weeks in storage in an attempt to reduce chemical injury. Some fruit was vacuum infiltrated (VI) with CaCl₂ for 1 or 3 min under 250 mm Hg vacuum and then held in the solution 1 min after vacuum release. Another comparison involved VI with either clean CaCl₂ solution or CaCl₂ solution contaminated with *Penicillium expansum* spores (ca. 3×10^4 spores/ml), as might occur after several hours of use.

Two commercial scald inhibitors (Chemley Products Co.) were compared: liquid concentrate "160" DPA used at 2000 ppm and liquid concentrate "260" ethoxyquin (ETOQ) used at 2700 ppm. Two other antioxidants, butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) were evaluated for scald control at 2,000 ppm. Some treatments included the wetting agent Tween-20 (0.1% v/v). Apples were dipped for 30 sec, then drained briefly and packed in fiberboard cartons.

Storage was at 0°C with 90% relative humidity for 4 to 6 months and then fruit were ripened 6 days at 20°. The CA chamber for the 'Stayman' apples was maintained at 3% O₂ and 2% CO₂. Firmness was measured on opposite pared sides of 16 or 20 apples per treatment per orchard with a Magness-Taylor penetrometer. Apples also were examined for decay, scald, other defects and possible injury after storage and after ripening. All fruit were eventually cut for internal examination. For Ca analysis, 10 fruit per treatment from each farm were washed in distilled water and a stem to calyx sector cut to the cord was removed. Peel and

¹Received for publication April 18, 1981.

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked advertisement solely to indicate this fact.

²The authors acknowledge the valued assistance of George A. Brown, Biological Laboratory Technician.

³Mention of a trademark, proprietary product or vendor does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture and does not imply its approval to the exclusion of other products or vendors that may also be suitable.

Table 1. Effects of scald inhibitors, CaCl_2 and other postharvest treatments on condition of 'Delicious' apples after 6 months at 0°C plus 6 days at 20° .²

Treatment	Firmness (kg)	Total scald (%)	Decay (%)	Calyx injury (%)	Flesh Ca (ppm)
Control, H_2O dip	6.1 a ^y	66 a	4.1 a	0.5 a	242 a
2,000 ppm DPA ^x	6.0 a	41 b	2.2 a	0.5 a	260 ab
2,700 ppm ETOQ ^x	6.0 a	45 bc	1.9 a	1.2 a	259 ab
3% CaCl_2 (lab grade)	6.0 a	59 ab	1.8 a	2.2 a	339 bc
3% CaCl_2 + 2,000 ppm DPA	5.8 a	36 c	1.5 a	2.2 a	286 abc
3% CaCl_2 + 2,700 ppm ETOQ	5.9 a	43 bc	1.7 a	1.8 a	292 abc
3% CaCl_2 + 0.2% Keltrol	5.7 a	47 bc	4.0 a	5.1 a	459 d
3% CaCl_2 + 0.2% Keltrol, rinse after 2 wks.	5.9 a	51 abc	2.3 a	3.3 a	328 abc
3% CaCl_2 , VI 1 min, clean soln ^w	6.1 a	45 bc	11.3 b	10.7 b	456 d
3% CaCl_2 , VI 1 min, contaminated soln	6.1 a	55 abc	38.9 c	11.1 b	374 c
3% CaCl_2 , VI 1 min, rinse immediately	6.2 a	52 abc	11.8 b	1.1 a	288 abc
3% CaCl_2 , 45°C , 30 sec, warm fruit	6.0 a	43 bc	2.5 a	1.5 a	341 bc
3% CaCl_2 , 45°C , 1 min, cold fruit	5.9 a	59 ab	1.4 a	2.6 a	294 abc

²Values are means of fruit from 4 orchards, 1978-79 season, 75-90 apples per treatment per orchard. All postharvest chemical treatments are for 30 sec unless stated otherwise. Initial fruit firmness averaged 7.2 kg (15.9 lb.).

^yMean separation within columns by Duncan's multiple-range test, 5% level.

^xChemley Products Co., liquid conc "160" No Scald DPA and liquid conc. "250" No Scald Ethoxyquin.

^wVI = vacuum infiltrated 1 min 250 mm Hg vacuum. Contaminated soln contained *Penicillium* spores.

core tissue were discarded. These composited flesh samples then were freeze dried, ground and 0.5 g samples were ashed, dissolved in 5 ml 2 N HCl, filtered and diluted into 100-ml flasks, and analyzed for Ca with a Jarrell-Ash atomic absorption spectrophotometer. All Ca values are reported on a dry-wt basis.

Results and Discussion

'Delicious' apples. In earlier experiments with 'Delicious', a 4% CaCl_2 treatment, with or without 0.3% Keltrol, reduced senescent breakdown and bitter pit (6). Neither disorder was encountered in the tests reported here. A 30-sec dip in 3% CaCl_2 alone or with 0.2% Keltrol increased flesh Ca levels but not flesh firmness after 6 months of storage (Table 1). Increasing the dip temperature to 45°C did not result in further increases in flesh Ca. The greatest flesh Ca concentrations were obtained when Keltrol or VI were used with CaCl_2 , so long as the apples were not subsequently rinsed. However, the VI fruit developed more decay and calyx burn, and were no firmer or otherwise in better condition than fruit dipped in CaCl_2 . Rinsing fruit in water immediately after VI prevented calyx injury but reduced Ca uptake during storage.

'Delicious' apples that were VI with CaCl_2 contaminated with *Penicillium* spores developed almost 40% decay after 6 months of

storage plus 6 days at 20°C (Table 1). This problem will have to be overcome if VI becomes a commercial practice.

The 'Delicious' fruit, which were picked early to midseason, developed extensive storage scald even though they were treated with liquid concentrate formulations of DPA or ethoxyquin. Three of the 4 lots were 'Starkrimson', which has been reported to be more susceptible to scale than other strains if the apples are harvested early (R. M. Smock, Cornell Fruit Handling and Storage Newsletter, July 1979). 'Starkrimson' also usually develops less water core than other 'Delicious' strains at harvest (9), so early picking should be avoided. Dip treatments of 3% CaCl_2 with either DPA or ethoxyquin also failed to control scald but caused no significant increases in skin injury. In an earlier study (6), 4% CaCl_2 + 0.3% Keltrol + ethoxyquin caused severe skin injury to both 'Delicious' and 'Golden Delicious'.

'Stayman' apple. In 5 seasons of testing this cultivar usually has responded favorably to CaCl_2 treatment (5, 6). A postharvest dip in 4% CaCl_2 retarded softening of 'Stayman' stored 4 months at 0°C (Table 2). Softening was also retarded after only 2 months of storage when laboratory grade CaCl_2 was used. After 4 months of storage the apples treated with laboratory grade CaCl_2 were more firm and had less breakdown than apples treated with com-

Table 2. Effects of commercial and laboratory grade CaCl_2 , with and without the adjuvant Keltrol, on the condition of 'Stayman' apples stored 2 and 4 months at 0°C .²

Treatment ^y	Firmness (kg)		Flesh Ca (ppm)		Breakdown (%)
	2 mo.	4 mo.	2 mo.	4 mo.	4 mo.
Control, H_2O dip	5.6 a ^x	4.6 a	200 a	230 a	20.0 a
4% CaCl_2 , comm. grade	5.8 ab	4.9 b	285 ab	275 a	8.3 b
4% CaCl_2 , lab grade	6.1 bc	5.1 c	304 ab	274 a	4.9 c
4% CaCl_2 , comm. grade + 0.3% Keltrol	6.0 bc	5.4 d	482 c	612 b	0.8 d
4% CaCl_2 , lab grade + 0.3% Keltrol	6.3 c	5.5 d	384 bc	563 b	0.8 d

²Values are means for fruit from 4 orchards. Initial fruit firmness 6.8 kg (15.0 lbs).

^yPostharvest 30-sec dips. Commercial grade CaCl_2 was 78% Dow flake; lab grade CaCl_2 was 76% Fisher USP. Breakdown recorded after 4 months 0°C + 6 days 20° .

^xMean separation within columns by Duncan's multiple-range test, 5% level.

Table 3. Effects of scald inhibitors and CaCl₂ postharvest dips on condition of 'Stayman' apples after storage in air and CA at 0°C plus 6 days at 20°C^z

Type storage and treatment ^y	Firmness (kg) ^x		Scald (%)	Flesh Ca (ppm)	Calyx injury (%)
	1st exam	2nd exam			
Air storage:					
Control, H ₂ O dip	5.3 ab ^w	4.8 a	72 a	304 a	3.4 abc
2,000 ppm DPA	5.2 a	4.8 a	4 e	290 a	2.0 ab
2,700 ppm ETOQ	5.1 a	4.8 a	3 e	295 a	2.2 ab
4% CaCl ₂ + 0.1% Tween	5.4 bc	5.0 b	53 b	357 b	4.7 abc
4% CaCl ₂ + DPA	5.3 ab	4.9 ab	3 e	365 b	6.0 c
4% CaCl ₂ + ETOQ	5.3 ab	4.9 ab	3 e	330 ab	5.1 bc
4% CaCl ₂ + 0.2% Keltrol	5.5 bc	5.3 c	51 b	505 c	8.9 d
4% CaCl ₂ + 0.2% Keltrol + DPA	5.6 c	5.2 c	2 e	490 c	11.9 e
4% CaCl ₂ + 0.2% Keltrol + ETOQ	5.5 bc	5.2 c	3 e	513 c	10.0 de
2,000 ppm BHT + Tween	5.3 ab	4.8 a	21 cd	----	1.7 a
2,000 ppm BHA + Tween	5.3 ab	4.8 a	32 c	----	2.7 ab ^v
CA storage:					
Control, H ₂ O dip	6.8 d	6.0 d	17 de	316 a	1.5 a
4% CaCl ₂ + 0.1% Tween	6.9 d	6.1 d	12 de	368 b	2.9 abc

^zAvg. values for 4 orchards in 1978 and 4 orchards in 1979, with 6 and 5 months of storage, respectively. Mean initial fruit firmness 8.1 kg (17.8 lb.).

^yPostharvest 30-sec dips with laboratory grade CaCl₂ and/or DPA and ethoxyquin liquid concentrates from Chemley Products Co.

^xFirst exam after storage, second exam after holding at 20°C.

^wMean separation within columns by Duncan's multiple range test, 5% level.

^vBHA (butylated hydroxyanisole) caused chemical burn on 37% of the fruit in 1979.

mercial grade CaCl₂. Keltrol increased flesh Ca and flesh firmness after 4 months of storage, as reported for 'Spartan' apples by Mason et al. (14). Senescent breakdown was reduced from 20% in control fruit stored 4 months + 6 days at 20°, to less than 1% in fruit that was dipped in 4% CaCl₂ + Keltrol. Breakdown also was reduced by a CaCl₂ dip without Keltrol.

In another series of 8 tests, 'Stayman' dipped in CaCl₂ + Tween and CaCl₂ + Keltrol were somewhat firmer than control

fruit after storage plus 6 days at 20°C (Table 3). The increased flesh firmness appeared to be directly related to the Ca in the flesh.

CaCl₂ appeared to be compatible with DPA or ethoxyquin for use on 'Stayman' apples. There was no skin injury from the combined treatment other than slight calyx darkening on a few apples. Calyx injury, although not a serious market defect, was more extensive when Keltrol was used.

Table 4. Effects of postharvest CaCl₂ treatments on the condition of 'Stayman' apples stored at 0°C plus 6 days at 20°C^z.

Season and treatment ^y	Firmness (kg)	Decay (%)	Flesh Ca (ppm)	Calyx injury (%)
1978-79 (6 months of storage):				
Control, H ₂ O dip	4.8 a ^x	3.0 a	335 a	2.1 a
4% CaCl ₂ (lab grade)	5.0 b	0.7 a	414 ab	7.6 ab
4% CaCl ₂ + 0.2% Keltrol	5.2 c	1.0 a	543 c	8.1 ab
4% CaCl ₂ + 0.2% Keltrol, rinse after 2 wks	5.1 bc	0.7 a	441 b	3.3 ab
4% CaCl ₂ VI 1 min, clean soln	5.2 c	1.9 a	444 b	11.7 ab
4% CaCl ₂ VI 1 min, contaminated ^w	5.2 c	20.4 b	----	12.6 b
1979-80 (5 months of storage):				
Control, H ₂ O dip	4.8 a	0.7 a	274 a	4.8 ab
4% CaCl ₂ (lab grade)	5.0 ab	0.7 a	301 a	4.6 ab
4% CaCl ₂ + 0.2% Keltrol	5.3 c	2.7 a	467 b	9.9 c
4% CaCl ₂ VI 1 min, clean	5.4 c	0.9 a	341 a	9.9 c
4% CaCl ₂ VI 3 min, clean	5.3 c	0.7 a	317 a	8.7 bc
2% CaCl ₂ VI 1 min, clean	5.1 bc	0.9 a	291 a	5.8 abc
2% CaCl ₂ VI 3 min, clean	5.2 bc	1.6 a	271 a	4.1 a

^zAverages for 4 orchards in each season.

^yPostharvest 30-sec dips unless stated otherwise. VI = vacuum infiltrated at 250 mm Hg vacuum. Tween-20 used with all CaCl₂ treatments in 1979-80 test.

^xMean separation within columns and within seasons by Duncan's multiple-range test, 5% level.

^wSolution contaminated with *Penicillium* spores (ca. 3×10^4 spores/ml).

Liquid concentrate formulations of DPA and ethoxyquin used alone or in combination with 4% CaCl_2 gave excellent scald control on 'Stayman' (Table 3). A CaCl_2 dip alone reduce scald significantly in these tests but did not provide adequate control. Both BHT and BHA, possible substitutes for DPA and ethoxyquin, reduced scald significantly, but DPA and ethoxyquin were much more effective. Also, BHA caused severe skin burn (37% damaged) in 1980.

'Stayman' apples stored in CA averaged 1.5 kg firmer than comparable air-stored fruit (Table 3). Dipping fruit in CaCl_2 before CA increased flesh Ca but did not significantly affect flesh firmness. Significant effects of CaCl_2 on 'Stayman' in CA, including a reduction in senescent breakdown, were noted previously (6).

Additional tests with CaCl_2 for 'Stayman' are summarized in Table 4. Rinsing fruit 2 weeks after dipping in CaCl_2 + Keltrol did not significantly reduce the calyx burn found after 6 months of storage. Flesh Ca after storage was ca 100 ppm lower in rinsed than in non-rinsed apples but the rinsed fruit was not significantly softer. One-minute VI with CaCl_2 resulted in firmness values similar to those produced by CaCl_2 + Keltrol. VI with CaCl_2 soln contaminated with *Penicillium* spores resulted in excessive decay (20%).

In the 1979-80 season, a 4% CaCl_2 + Tween dip did not increase flesh firmness. However, flesh firmness was significantly increased by 4% CaCl_2 + Keltrol and by VI of 2 or 4% CaCl_2 for 1 or 3 min. An advantage of VI over dip treatment is that apples can be rinsed immediately after treatment, reducing the possibility of fruit injury and corrosion of equipment (23).

In conclusion, there was no advantage of dips or VI with 3% CaCl_2 for 'Delicious' when bitter pit or storage breakdown were not problems. Scald of some strains of 'Delicious' was not controlled with either DPA or ethoxyquin when fruit was picked early to midseason some years. A dip of 4% CaCl_2 on 'Staymans' usually gave firmer fruit and greatly reduced senescent breakdown after storage. Uptake of Ca and firmness were highest when CaCl_2 was VI or applied with Keltrol. Liquid concentrate formulations of DPA and ethoxyquin used alone or combined with CaCl_2 provided excellent scald control for 'Stayman' apples.

Literature Cited

1. Bangerth, F., D. R. Dilley, and D. H. Dewey. 1972. Effect of postharvest calcium treatments on internal breakdown and respiration of apple fruits. *J. Amer. Soc. Hort. Sci.* 97:679-682.
2. Betts, H. A. and W. J. Bramlage. 1977. Uptake of calcium by apples from postharvest dips in calcium chloride solutions. *J. Amer. Soc. Hort. Sci.* 102:785-788.
3. DeVilliers, J. F. and A. N. Hanekom. 1977. Factors by which the postharvest uptake of calcium by Golden Delicious apples is influenced. *Decid. Fruit Grow.* 27(3):85-91.
4. Dewey, D. H. and D. R. Dilley. 1974. Increasing storage and market life of Jonathan apples. *Mich. State Univ. Ext. Bul. E-627 (Rev.)*.
5. Hardenburg, R. E. and R. E. Anderson. 1977. Softening and breakdown of 'Stayman' apples reduced in air and CA storage by postharvest dip in calcium chloride solutions. *HortScience* 12:387. (Abstr.)
6. Hardenburg, R. E. and R. E. Anderson. 1979. Effects of postharvest calcium and other chemical treatments on softening, scald, bitter pit and breakdown of stored apples. *Proc. 15th Intern. Congr. Refrig. Venice, Sept. 22-29, Vol. 3, paper C2-29*.
7. Johnson, D. S. 1979. New techniques in the post-harvest treatment of apple fruits with calcium salt. *Commun. Soil Sci. & Plant Anal.* 10:373-382.
8. Lidster, P. D. and S. W. Porritt. 1978. Some factors affecting calcium uptake by apples dipped after harvest in calcium chloride solution. *Can. J. Plant Sci.* 58:35-40.
9. Lord, W. J., R. A. Damon, Jr., and J. F. Anderson. 1980. A comparison of tree size, productivity, and fruit quality of 'Delicious' strains. *J. Amer. Soc. Hort. Sci.* 105:883-887.
10. Mason, J. L. 1976. Calcium concentration and firmness of stored 'McIntosh' apples increased by calcium chloride solution plus thickener. *HortScience* 11:504-505.
11. Mason, J. L. and B. G. Drought. 1975. Penetration of calcium into 'Spartan' apple fruits from a post-harvest calcium chloride dip. *J. Amer. Soc. Hort. Sci.* 100:413-415.
12. Mason, J. L., B. G. Drought, and J. M. McDougald. 1974. Effect of calcium chloride dip on senescent breakdown, firmness and calcium concentration in 'Spartan' apple. *HortScience* 9:596.
13. Mason, J. L., J. J. Jasmin, and R. L. Granger. 1975. Softening of 'McIntosh' apples reduced by a post-harvest dip in calcium chloride solution plus thickener. *HortScience* 10:524-525.
14. Mason, J. L., J. M. McDougald, and B. G. Drought. 1974. Calcium concentration in apple fruit resulting from calcium chloride dips modified by surfactants and thickeners. *HortScience* 9:122-123.
15. Menendez, R. A. and B. W. Poovaiah. 1980. Factors influencing calcium infiltration into apple fruits. *HortScience* 15:391. (Abstr.)
16. Padfield, C. A. S. 1979. Use of calcium to reduce storage breakdown in Spartan apples. *N. Zeal. J. Expt. Agr.* 7:379-381.
17. Porritt, S. W. and P. D. Lidster. 1978. Effect of time of washing on calcium uptake, breakdown and condition of Spartan apples dipped in calcium chloride solution after harvest. *Can. J. Plant Sci.* 58:41-44.
18. Reid, M. S. and C. A. S. Padfield. 1975. Control of bitter pit in apples with lecithin and calcium. *N. Zeal. J. Agr. Res.* 18:383-385.
19. Riley, R. G. and P. E. Kolattukudy. 1976. Effect of treatment with calcium ion-containing formulations on the firmness of Golden Delicious apples. *HortScience* 11:249-251.
20. Schumacher, R., F. Fankhauser, and W. Stadler. 1976. Experiments with calcium chloride, antitranspirants and boric acid to reduce the occurrence of bitter pit. *Schweiz. Zeitschr. für Obst-und Weinbau* 112:300-304.
21. Scott, K. J. and R. B. H. Wills. 1975. Postharvest application of calcium as a control for storage breakdown of apples. *HortScience* 10:75-76.
22. Scott, K. J. and R. B. H. Wills. 1977. Vacuum infiltration of calcium chloride: a method for reducing bitter pit and senescence of apples during storage at ambient temperatures. *HortScience* 12:71-72.
23. Scott, K. J. and R. B. H. Wills. 1979. Effects of vacuum and pressure infiltration of calcium chloride and storage temperature on the incidence of bitter pit and low temperature breakdown of apples. *Austral. J. Agr. Res.* 30:917-928.
24. Sharples, R. O. and D. S. Johnson. 1976. Post-harvest chemical treatments for the control of storage disorders of apples. *Ann. Appl. Biol.* 83:157-167.
25. Shear, C. B. (ed.) 1979. International symposium on calcium nutrition of economic plants. *Commun. Soil Sci. & Plant Anal.* 10(1&2).
26. Taylor, A., B. W. Poovaiah, and J. A. Hopfinger. 1978. Calcium infiltration treatments aid in Golden Delicious bitter pit control. *Goodfruit Grow.* 29(8):2, 24.
27. Webster, D. H. and F. R. Forsyth. 1979. Partial control of bitter pit in Northern Spy apples with a post-harvest dip in calcium chloride solution. *Can. J. Plant Sci.* 59:717-723.