

7. Purvis, A. C., K. Kawada, and W. Grierson. 1979. Relationship between midseason resistance to chilling injury and reducing sugar level in grapefruit peel. *HortScience* 14:227-229.
8. Purvis, A. C. and W. Grierson. 1982. Accumulation of reducing sugar and resistance of grapefruit peel to chilling injury as related to winter temperatures. *J. Amer. Soc. Hort. Sci.* 107:139-142.
9. Steponkus, P. L. 1971. Cold acclimation of *Hedera helix*. Evidence for a two phase process. *Plant Physiol.* 47:175-180.
10. Ting, S. V. and R. L. Rouseff. 1979. Proline content in Florida frozen concentrated orange juice and canned grapefruit juice. *Proc. Fla. State Hort. Soc.* 92:143-145.
11. Weiser, C. J. 1970. Cold resistance and injury in woody plants. *Science* 169:1269-1278.
12. Yelenosky, G. 1975. Cold hardening in citrus stems. *Plant Physiol.* 56:540-543.
13. Yelenosky, G. 1977. The potential of citrus to survive freezes. *Proc. Intern. Soc. Citriculture* 1:199-203.
14. Yelenosky, G. 1979. Accumulation of free proline in citrus leaves during cold hardening of young trees in controlled temperature regimes. *Plant Physiol.* 64:425-427.
15. Yelenosky, G. and C. L. Guy. 1977. Carbohydrate accumulation in leaves and stems of 'Valencia' orange at progressively colder temperatures. *Bot. Gaz.* 138:13-17.
16. Young, R. and W. D. Bell. 1974. Photosynthesis in detached leaves of cold-hardened citrus seedlings. *J. Amer. Soc. Hort. Sci.* 99:400-403.

*J. Amer. Soc. Hort. Sci.* 107(2):226-231. 1982.

## Growth and Respiration of Dormant Flower Buds of *Pyrus communis* and *Pyrus calleryana*<sup>1</sup>

Mary E. Cole,<sup>2</sup> Theophanes Solomos,<sup>3</sup> and Miklos Faust<sup>4</sup>

Fruit Laboratory, Agricultural Research, U. S. Department of Agriculture, Beltsville, MD 20705

*Additional index words.* *Prunus armeniaca*, *Prunus salicina*, *Prunus avium*, *Prunus domestica*, *Prunus serrulata*, *Malus hupehensis*, *Malus domestica*

**Abstract.** Respiration of flower-buds of *Pyrus communis* L., a late blooming species, and *P. calleryana*, an early blooming species, was investigated throughout the winter. Respiration of *P. calleryana* Decne at 5°C was twice as high as that of *P. communis*, whereas the respiration rates were similar at 25°. A large portion (60-70%) of the respiration at 5° was cyanide resistant in *P. calleryana* and much less in *P. communis*. The combination of inhibitors, cyanide (KCN) and salicylhydroxamic acid (SHAM), still only partially inhibited respiration. The residual respiration was much higher for *P. calleryana* than for *P. communis*. The nature of the residual respiration is not known.

It appears that respiration of *P. calleryana* is less sensitive to cold than that of *P. communis*. *P. calleryana* also appears to complete the development of its buds during the winter and blooms early. In contrast, *P. communis*, the species which has a respiratory pathway less operational at relatively low temperatures, delays the completion of its buds until the spring when the temperature warms up, and consequently flowers later. This hypothesis was evaluated by measuring the mid-winter respiration of 5 early and 5 late blooming species of fruit trees, and the bloom date was evaluated relative to the type of respiration as described above.

Resting buds of fruit trees in general and pears in particular enter into dormancy with partially developed buds. The buds apparently complete their development by the time growth is resumed. The time of bloom varies from year to year and from species to species. There can be differences of up to one month for a species in a given location (2). The major factor regulating the time of bloom is temperature. A certain amount of cold is required to break the rest. After the cold requirement has been satisfied, a certain amount of heat is needed to enable the tree to bloom. It is conceivable that the time of bloom depends on the rate of development of buds during dormancy which in turn is influenced by temperature.

The time of bloom can be genetically transmitted (9) and intermediate bloom types created between early and late blooming pears. Generally the low chilling types (*P. calleryana*) also bloom early. The late blooming types (*P. communis*) bloom earlier when grafted on roots of early blooming types (*P. calleryana*) (24). Strausz (20) stated that this translocatable reduction of chilling requirement was due to a shift in growth promoters.

Pear buds are metabolically active during the winter. Thom (21) measured respiration of dormant buds of *P. communis* 'Hardy' and found that O<sub>2</sub> consumption increased six-fold when growth resumed in the spring. Zimmerman et al. (26) found that incorporation of uracil and valine into alcohol insoluble components of buds was high during the winter indicating that pear buds were not "resting" but metabolizing actively without visible in-

<sup>1</sup>Received for publication May 9, 1981.

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

<sup>2</sup>Present Address: Dept. of Agr. Economics, University of Maryland, College Park, MD, 20742.

<sup>3</sup>Professor, Dept. of Horticulture, University of Maryland, College Park, MD 20742.

<sup>4</sup>Plant Physiologist, Fruit Laboratory, SEA, U. S. Department of Agriculture, Beltsville, MD 20705.

<sup>5</sup>Mention of a trademark name or a proprietary product does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture and does not imply its approval to the exclusion of other products that may be suitable.

dication of growth. Chaudhry et al. (5) found that at the time of bud break concentrations of soluble-N, ester-P, and RNA-P increased markedly.

Strausz (20) found that *P. communis* had considerably higher levels of abscisic acid (ABA) than did *P. calleryana* during the winter. However, ABA did not disappear from the whole buds when rest was broken, nor did it disappear gradually during chilling. An unidentified growth promoter did not appear in significantly large amounts until growth had resumed.

Brown and Kotob (4) reported that the fresh and dry weights of *P. communis* flower buds slightly increased during October and November. Development of bud weight increases have not been reported for *P. calleryana* although the buds of *P. calleryana* are larger than those of *P. communis* at the end of the growing period.

None of the information available to date explains the major physiological differences in time of bloom between early and late blooming types. The purpose of this work was to examine the respiratory mechanism of *P. communis* and *P. calleryana* buds to discover the sources of energy for dormant bud development and find explanations for differences in bloom.

### Materials and Methods

**Plant material.** Flower buds from two species of pear, *Pyrus communis* 'US 309' and *P. calleryana* 'Bradford' were used in the studies reported here. For comparative studies additional deciduous fruit tree species were included. These species were: *Malus hupehensis* (Pamp.) Rehd; *M. sylvestris* Mill; 'Golden Delicious'; *Prunus salicina* Lindl.; *P. domestica* L.; *P. avium* L.; *P. serrulata* Lindl.; 'Kwanzan'; *P. armeniaca* L.; 'Wilson Delicious'; and *P. armeniaca* L. 'US 566'.

**Measurement of respiration.** The scales were removed from the flower buds leaving the partially developed flower clusters (later referred to as inflorescences) intact. Enough inflorescences were prepared to obtain approximately 100 mg of tissue per flask. These were weighed and placed in Gilson respiration flasks. The concentrations of the reagents used were as follows: 3mM phosphate buffer, pH 5.3, 3mM SHAM and 0.2mM KCN. The method of Robbie (17) was used to absorb CO<sub>2</sub> and maintain constant cyanide concentration in the fluid. The rate of oxygen uptake is expressed as  $\mu\text{l O}_2$  at NTP/g<sup>-1</sup>h<sup>-1</sup> (23).

The respiration rate of the pear flower bud inflorescences was measured at 25°, 18°, 10°, and 5°C periodically throughout dormancy. Since respiration at 18° and 10° was intermediate between the 25° and 5° data it is not reported here but is available elsewhere (6). The survey of respiration rates of the other species was performed in early February.

**Ambient orchard temperatures.** Chill units were calculated by summing the hours when the temperatures were between 7.2° and 0°C for each day. Heat sums, in degree-days above 4.5°, were calculated by summing the differences between 4.5° and the average daily temperatures after January 31.

**Measurement of bud growth.** Whole buds and inflorescences were weighed to determine bud growth each time buds were prepared for respiration.

### Results

**Respiration rates of *Pyrus calleryana* inflorescences.** At 25°C the rate of O<sub>2</sub> uptake of *P. calleryana* inflorescences increased steadily from late September until early March and then began to increase rapidly (Fig. 1). SHAM had no effect on the rate of respiration at 25° (Fig. 1B). The effect of cyanide on the rate of oxygen




Fig. 1. Respiration of *Pyrus calleryana* inflorescences measured at 25°C.

uptake varied with the calendar date. Thus, the inhibitor decreased respiration by about 50% from late September through the middle of October. This inhibition was decreased to about 20% in November, to gradually increase to 30–35% in late February and early March. It should be noted that the absolute values of the cyanide-resistant respiration followed the same pattern as the control samples up to early March. Thereafter, cyanide-resistant respiration declined sharply while that of the control increased rapidly. These changes in the absolute magnitudes of the two respiratory components resulted in respiration being strongly inhibited by cyanide in the latter part of March.

In most plant tissues the combined application of KCN and SHAM does not inhibit completely the rate of oxygen uptake (7). The nature of this "residual" rate of respiration is unknown. Residual respiration increased slightly during the fall and winter, to decrease to almost zero in the latter part of March (Fig. 1A, B). At this stage, the inhibition of oxygen uptake by the combined application of SHAM and KCN was similar to that produced by cyanide alone.

At 5°C, respiration of the controls also increased steadily through the year (Fig. 2A). Further, the increment was similar to




Fig. 2. Respiration of *Pyrus calleryana* inflorescences measured at 5°C.

that of 25°: about 600% (Fig. 2A). However, cyanide resistant respiration was higher at 5° than at 25°. In early October, cyanide reduced respiration by 30% while in January the inhibition was only 12%. In addition, not only did the percentage inhibition decrease during the fall and winter, but the absolute values of the cyanide-resistant component of respiration increased during the same period. In late March, cyanide inhibited respiration by 44%. Residual respiration increased during the period of rest, to become zero by late March (Fig. 2A, B).

**Respiration of *P. communis* inflorescences.** The respiration of *P. communis* inflorescences was also measured at 25° and 5°C from late fall until spring. As with *P. calleryana*, respiration of *P. communis* also increased during the above period. At 25° the respiration of the controls increased by about four-fold (Fig. 3A), cyanide inhibited respiration by 37% in late September and the inhibitory effect tended to increase with time (Fig. 3B), in contrast to the decrease with flower bud axes of *P. calleryana* (Fig. 1B). The residual respiration was also lower with *P. communis* than *P. calleryana*. The combined application of 3 mM SHAM and 0.2 mM KCN decreased respiration by 46% more than with cyanide alone in September (Fig. 3B). This difference decreased to only 10% from December through February. In early March it increased to 26%, to fall again to zero by the end of March (Fig. 3B).

The pattern of respiration at 5°C was similar to that at 25° (Figs. 4A, 3A). The inhibition by cyanide was lower at 5° than 25° (Fig. 4B). Thus, cyanide inhibited respiration by an average of 45% and 59% at 5° and 25°, respectively. Residual respiration was low

at 5°. In September and October it averaged 30% below that caused by cyanide alone. The difference increased to 58% in November and then slowly decreased to 3% by late March.

**Bud growth.** The fresh weight of *P. calleryana* whole buds increased gradually from the fall until mid-February and then began to increase sharply. With *P. communis*, whole flower buds doubled in weight during November after which bud weight remained constant from early December until late February when active growth began. The growth of the inflorescences followed a pattern almost identical to that of *P. calleryana*. Throughout the year the weight of the inflorescences of *P. calleryana* flower buds was a much smaller percentage of the total bud weight than for *P. communis* (Fig. 5).

**Respiration of inflorescences of several species of fruit trees.** As measured at 5°C the mid-winter respiration rates of inflorescences of the earlier blooming fruit trees were higher than those of the later blooming types, with the exception of *P. avium* (Table 1).

The average percentage of respiration resistant to cyanide was higher for the earlier blooming types than for the later blooming ones.

With the exception of *Prunus serrulata* the respiration, which was resistant to the combined treatment of SHAM and KCN, was higher for the earlier blooming types.




Fig. 3. Respiration of *Pyrus communis* inflorescences measured at 25°C.

Table 1. Mid-winter respiration rates of the flower buds of several species of deciduous fruit trees measured at 5°C.

Species	Date of full bloom	Inflorescences (g)	Respiration ( $\mu\text{l}/\text{O}_2 \text{ g}^{-1} \text{ fresh wt}$ )				%of control		
			Control	3 mM SHAM	0.2 mM KCN	KCN + SHAM	3 mM SHAM	0.2 mM KCN	KCN+ SHAM
<i>Prunus armeniaca</i> 'Wilson Delicious'	Apr. 5	169	177	149	104	92	84	59	52
<i>Prunus armeniaca</i> 'US 566'	Apr. 8	137	193	177	154	122	92	80	63
<i>Prunus salicina</i>	Apr. 8	403	202	181	182	170	90	90	85
<i>Pyrus calleryana</i>	Apr. 10	48	184	182	139	81	99	76	46
<i>Prunus avium</i>	Apr. 11	82	82	79	71	45	97	87	55
<i>Pyrus communis</i>	Apr. 17	88	111	114	51	14	103	46	13
<i>Prunus domestica</i>	Apr. 18	211	112	103	59	33	92	53	30
<i>Malus hupenhensis</i>	Apr. 18	133	165	163	85	56	99	52	34
<i>Malus domestica</i> 'Golden Delicious'	Apr. 24	48	81	85	63	31	104	77	38
<i>Prunus serrulata</i>	Apr. 26	105	87	98	75	55	113	86	63

Calculations using linear regression showed that there was no correlation between the number of inflorescences per gram and the rate of respiration.

**Ambient orchard temperatures.** Rest is assumed to be completed in pears after exposure to about 1,100 hours of chilling. This would have occurred about the second week of January in the Beltsville area (Fig. 6).

The heat sums required for bloom are shown in Figure 6. *P. calleryana* bloomed on April 10 after 180 degree-days of heat. *P. communis*, which flowered on April 17, required 233 degree-days.

## Discussion

Since outdoor temperatures were low during dormancy, ranging from  $-10^{\circ}$  to  $10^{\circ}\text{C}$ , the respiration at  $5^{\circ}$  may represent more closely what occurred in the buds in nature whereas the data obtained at  $25^{\circ}$  indicate the capacity of buds to respire. Both rates, the assumed natural respiration and the capacity of respiration were high for both species. Rates were in the range of 50 to 200 and 400 to 2000  $\mu\text{l O}_2 \text{ g}^{-1} \text{ fw hr}^{-1}$  for  $5^{\circ}$  and  $25^{\circ}$  respectively.

At  $5^{\circ}\text{C}$  the midwinter respiration of *P. calleryana* was double that of *P. communis*. In contrast, the difference in respiration be-





Fig. 4. Respiration of *Pyrus communis* inflorescences measured at  $5^{\circ}\text{C}$ .





Fig. 5. Fresh weights of whole flower buds and inflorescences of *Pyrus calleryana* (upper) and *P. communis* (lower).




Fig. 6. Chill units (hours) and heat units ( $^{\circ}\text{C} \times \text{days}$ ) for the 1979–1980 season.

tween the two species at  $25^{\circ}$  was much less indicating that *P. calleryana* has a mechanism which provides an advantage to this species at low temperature only. The largest increase in respiration of *P. calleryana* at  $5^{\circ}$  occurred during the month of January, whereas the increase of *P. communis* came a month later. At  $25^{\circ}$  both species have shown further elevation of respiration during the month of March. The increase in respiration during January and February may indicate the end of the rest period for each species, respectively. The increase in respiration during March indicated the beginning of spring metabolic activity even though actual flowering occurred later in mid-April. These results are consistent with those of Thom (21) who found that *P. communis* buds kept in cold storage at  $2.5^{\circ}$  increased their respiration continuously for five months. However, the pear data differ from the respiration of buds in warm climates which was shown by Lavee (14) to decrease when buds go into dormancy and increase when dormancy ends.

It is noteworthy that the dormant buds of pears respire at low temperatures ( $5^{\circ}\text{C}$ ) and the inhibitor experiments indicate that at this temperature the capacity for both alternate and residual oxidases is much larger than at  $25^{\circ}$ . Up to 80% of the respiration of *P. calleryana* buds was resistant to cyanide inhibition. The cyanide resistant respiration of *P. communis* buds was much lower. This is consistent with the result of other researchers who demonstrated that chilling can stimulate respiration in many species. Goodwine et al. (11) and Shulze et al. (18) showed that a temperature of  $1^{\circ}$  stimulates cyanide resistant respiration in potato tubers. Leopold and Musgrave (15) showed that chilling of soybean cotyledons engages the cyanide resistant alternate pathway, and Yoshida and Tagawa (25) found that most of the respiration of *Cornus callus* tissue was cyanide resistant below  $15^{\circ}$ .

Cyanide-resistant respiration is a widespread phenomenon in higher plants (7, 12, 19). It has been suggested by Bonner (3) that the cyanide-resistant pathway may be a general constituent of all plants with resistance varying from as low as a few percent (7, 12) to as much as 100 percent in *Arum maculatum* (13). On the basis of the present data, it is not clear whether cyanide-resistant respiration is engaged in the absence of cyanide. It should be noted, however, that both the absolute value of cyanide-resistant respiration and the percent inhibition decrease in late March, when the growth resumes, and hence the demand for energy increases.

The combination of both inhibitors, SHAM and KCN, still only partially inhibited respiration. The residual respiration was much higher for *P. calleryana* than for *P. communis*. The nature of the residual respiration is not known. Tucker (22) proposed that

there may be a third pathway of respiration. Eskin et al. (8), Parish and Leopold (16), and Goldstein et al. (10) all pointed out the possibility that lipoxygenase activity may be responsible for residual  $\text{O}_2$  uptake. Since SHAM appears to inhibit lipoxygenase activity (10) this possibility is unlikely in our experiments.

During these experiments, the picture which emerged was that the species (*P. calleryana*) having a higher rate of respiration, including the cyanide resistant alternate pathway and residual respiration, had the advantage of completing the development of its buds at low temperature and bloomed early. In contrast, the species which did not have a relatively active respiratory pathway operational at low temperatures, or had it to a lesser degree, needed to wait until completion of bud development took place during the spring when the temperature warmed up.

To check this theory we determined midwinter respiration of several early and late blooming species at  $5^{\circ}\text{C}$ . In general the survey of 10 species has shown that the above supposition was correct with one exception. *Prunus serrulata* bloomed late but its respiration was more like that of the early-blooming species.

The results presented here strongly suggest that buds, which respire through a pathway that is not inhibited, or is inhibited to a lesser degree, by cold, complete their development and bloom earlier than do buds which do not have this ability. Much of the development may occur in early spring. This study also revealed that before bloom the buds switch to the cyanide sensitive pathway which is also temperature sensitive. Thus, decreasing the temperature at this time appears to be an effective way to delay bloom. In fact this is what happened in apples when sprinkler irrigation was used for delaying bloom (1). Controlling temperature by sprinkling is effective but impractical. Further work is necessary to find chemical controls for the respiratory shift which could delay bloom effectively.

#### Literature Cited

1. Anderson, J. L., G. L. Ashcroft, E. A. Richardson, J. F. Alfaro, R. E. Griffin, G. R. Hanson, and J. Keller. 1975. Effects of evaporative cooling on temperature and development of apple buds. *J. Amer. Soc. Hort. Sci.* 100:229–231.
2. Antsey, T. H. 1966. Prediction of full bloom date for apple, pear, cherry, peach, and apricot from air temperature data. *Proc. Amer. Soc. Hort. Sci.* 88:57–66.
3. Bonner, W. D. 1965. Mitochondria and electron transport. p. 89–123. In: J. Bonner and J. E. Varner (eds.). *Plant biochemistry*. Academic Press, New York.
4. Brown, D. S. and F. A. Kotob. 1957. Growth of flower buds of apricot, peach, and pear during the rest period. *Proc. Amer. Soc. Hort. Sci.* 69:158–164.
5. Chaudhry, W. M., T. C. Broyer, and L. C. T. Young. 1970. Chemical changes associated with the breaking of the rest period in vegetative buds of *Pyrus communis*. *Physiol. Plant.* 23:1157–1169.
6. Cole, M. E. 1980. Growth and respiration of dormant flower buds of *Pyrus communis* and *Pyrus calleryana* Decne. MS Thesis. Univ. of Maryland, College Park.
7. Day, A. D., G. P. Arron, and G. G. Laties. 1980. Nature and control of respiratory pathways in plants. The interaction of cyanide-resistant with cyanide sensitive pathway. p. 197. In: D. D. Davies (ed.) *The biochemistry of plants; a comprehensive treatise*. Vol. 2. Academic Press.
8. Eskin, N. A. M., S. Grossman, and A. Pinsky. 1977. Biochemistry of lipoxygenase in relation to food quality. *Crit. Rev. Food Sci. Nut.* 9:1–40.
9. Faust, M., R. Zimmerman, and T. van der Zwet. 1976. Genetic transmission of bloom date in pears. *HortScience* 11:59–60.

10. Goldstein, A. H., J. O. Anderson, and R. G. McDaniel. 1980. Cyanide-insensitive and cyanide-sensitive O<sub>2</sub> uptake in wheat. *Plant Physiol.* 66:488–493.
11. Goodwine, W., M. Mikai, S. Kaitems, and C. Frenkel. 1979. Development of respiration and cyanide-resistant respiration in potato tubers as influenced by low temperatures and hypobaric conditions. *Plant Physiol.* 63:158 (Abstr.).
12. Henry, M. F. and E. J. Nyns. 1975. Cyanide-insensitive respiration. An alternative mitochondrial pathway. *Sub-Cell. Biochem.* 4:1–65.
13. James, W. O. and H. Beevers. 1950 The respiration of *Arum spadix*. A rapid respiration, resistant to cyanide. *New Phytol.* 49:353–374.
14. Lavee, S. 1972. Dormancy and bud break in warm climates: considerations of growth regulator involvement. *Acta. Hort.* 34:225–234.
15. Leopold, A. C. and M. E. Musgrave. 1979. Respiratory changes with chilling injury of soybeans. *Plant Physiol.* 64:702–705.
16. Parrish, D. J. and A. C. Leopold. 1978. Confounding of the alternate respiration by lipoxygenase activity. *Plant Physiol.* 62:470–472.
17. Robbie, W. A. 1946. The quantitative control of cyanide in manometric experiments. *J. Cell. & Comp. Physiol.* 27:181–209.
18. Shulze, C., G. Wulster, H. Janes, and C. Frenkel. 1979. Interaction of low temperature and oxygen regimes on the stimulation of respiration, and cyanide-resistant respiration, in potato tubers. *Plant Physiol.* 63:103 (Abstr.)
19. Solomos, T. 1977. Cyanide-resistant respiration in higher plants. *Annu. Rev. Plant Physiol.* 28:279–297.
20. Strausz, S. D. 1970. A study of the physiology of dormancy in the genus *Pyrus*. PhD Thesis. Oregon State Univ., Corvallis.
21. Thom, L. C. 1951. A study of the respiration of Hardy pear buds in relation to the rest period. PhD Thesis. Univ. of California, Berkeley.
22. Tucker, M. L. 1978. The significance of cyanide-insensitive respiration in the ripening and concomitant rise in respiration of banana fruit slices. MS Thesis. Univ. of Maryland, College Park.
23. Umbreit, W. W., R. H. Burris, and J. F. Stauffer. 1972. *Manometric and Biochemical Techniques*. Burgess Pub. Co., Minneapolis.
24. Westwood, M. N. and N. E. Chestnut. 1964. Rest period chilling requirement of Bartlett pear as related to *Pyrus calleryana* and *P. communis* rootstocks. *Proc. Amer. Soc. Hort. Sci.* 84:82–87.
25. Yoshida, S. and F. Tagawa. 1979. Alteration of the respiratory function in chill-sensitive callus due to low temperature stress. I. Involvement of the alternate pathway. *Plant & Cell. Physiol.* 20:1243–1250.
26. Zimmerman, R. H., M. Faust, and A. W. Shreve. 1970. Glucose metabolism of various tissue of pear buds. *Plant Physiol.* 46:839–841.

*J. Amer. Soc. Hort. Sci.* 107(2):231–234. 1982.

## Selection for Resistance in *Phaseolus vulgaris* L. to White Mold Disease Caused by *Sclerotinia sclerotiorum* (Lib.) de Bary

M. H. Dickson, J. E. Hunter, M. A. Boettger, and J. A. Cigna

*Departments of Seed and Vegetable Sciences and Plant Pathology, New York State Agricultural Experiment Station, Geneva, NY 14456*

*Additional index words.* juvenile stem test, blossom test, genetic resistance

**Abstract.** Three populations of beans were screened for resistance to white mold using a mycelium/juvenile stem inoculation (JSI) method on 3-week-old plants and an ascospore/blossom test on plants in bloom was used to test resistance of survivors of the JSI test. There were few survivors in the JSI test: 0.8% in the 4-way cross of susceptible x susceptible, 2% in the intermediate x intermediate crosses and 3.8% in 10 *P. coccineus* lines with intermediate resistance. JSI tests of the progeny produced more survivors, 17% from the first 2 populations. There was good agreement between the JSI test on juvenile plants and the ascospore/blossom tests on blossoming plants respectively. The JSI test appears to be an efficient method with which to identify individual plants with moderate resistance.

White mold disease of beans has been neglected by bean breeders until recently. In 1973 Adams et al. (2) identified some *Phaseolus coccineus* lines with resistance. Abawi et al. (1) reported resistance in *P. vulgaris*, derived crosses with *P. coccineus*. Most *P. vulgaris* accessions were relatively susceptible. In field trials *P. vulgaris* cultivars Anderson et al. (3), Coyne et

al. (5) and Schwartz et al. (10) reported some resistance which could be attributed to avoidance due to plant growth habit in some cases and resistance in others.

Abawi et al. (1) used blooming plants that were sprayed with a suspension of ascospores. Hunter et al. (8) modified Abawi's ascospore-blossom inoculation (ABI) method (1). They also developed a second procedure, called limited term inoculation (8) which is referred to as juvenile stem inoculation (JSI) in this paper.

The high reliability of JSI and ABI (8) screening methods made possible the screening of many accessions of *Phaseolus* spp. (7), which led to the identification of a number of plant introductions with varying levels of resistance.

<sup>1</sup>Received for publication June 1, 1981. Journal Paper No. 3364 from the New York State Agricultural Experiment Station, Geneva, NY 14456.

The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.