

Effect of Liming Rate on Nitrification in a Pine Bark Medium

Alexander X. Niemiera¹ and Robert D. Wright²

Department of Horticulture, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061

Additional index words. container-grown, nitrogen, pH

Abstract. A 100% pine bark medium was amended with dolomitic lime at 0, 3, or 6 kg·m⁻³ and periodically fertilized with 210 ml of a nutrient solution containing 100 ppm N as (NH₄)₂SO₄. At the 3- and 6-kg lime treatments, medium solution NH₄-N concentrations decreased rapidly while NO₃-N concentrations increased. At 0 kg lime, the NH₄-N decrease was slower than at the 3- and 6-kg lime treatments and NO₃-N was not found. Similarly treated bark without plants was used to determine a NO₃-N accumulation rate (NAR). NAR was greatest at 6 kg of lime, except at the last 2 sampling dates, when NAR did not differ between 3 and 6 kg of lime. This lack of difference was attributed to a limiting NH₄-N supply at 6 kg of lime. In a 2nd experiment, NAR of bark treated with 6 kg of lime per m³ and fertilized with 300 ppm NH₄-N was 3 times greater than with bark treated with 100 ppm NH₄-N, thus supporting the contention that, over time, the NH₄-N supply of the 100-ppm treatment limited nitrification. These results indicate that the stimulative influence of lime on nitrification is subject to medium pH and NH₄-N status that changes over time.

Liming container media is a routine practice in the nursery industry. Liming increases medium pH, supplies Ca (and Mg in the case of dolomitic lime), and influences nutrient availability. An increased medium pH stimulates nitrification, the biological oxidation of NH₄⁺ to NO₃⁻. Thus, plant response to lime rate may be in part a response to the ionic form of N in the medium solution. Niemiera and Wright (7) treated pine bark with 6 kg of lime per m³ and fertilized with (NH₄)₂SO₄ and found a depletion of NH₄-N and an accumulation of NO₃-N in the medium solution. Chrusic and Wright (1) treated pine bark with dolomitic lime in the range of 0 to 8 kg of lime per m³ and reported that the 8-kg lime treatment had the lowest solution NH₄-N : NO₃-N ratio. This result was attributed to increased nitrifier populations and an increased adsorption of NH₄-N to bark particles. However, bark was fertilized with NH₄NO₃ in their experimentation, and the application of NO₃-N to the bark makes the influence of lime on nitrification difficult to discern. Niemiera and Wright (7) used a nitrification inhibitor in their experiment, and found the medium solution NH₄-N concentrations remained relatively high, indicating that NH₄-N depletion without NI was unequivocally a result of nitrification.

The ionic form of N has a significant influence on plant nutrient composition and growth (3, 4, 6). Chrusic and Wright (1) found that the growth of 3 woody genera was greatest at the 0 or 2 kg lime per m³ bark treatments and attributed this response in part to a greater NH₄-N : NO₃-N ratio. Other researchers working with woody plants (5, 9) have shown a similar response to such a ratio. Growers of container-grown crops commonly fertilize with NH₄-N and amend their media with lime. However, lime rates vary greatly within the industry, depending on the species grown and grower preference. The purpose of this research was to determine how lime rates affect nitrification in a pine bark medium.

Materials and Methods

Experiment 1. Pine bark used in this experiment was identical to bark described in a previous paper (7). Bark was moistened and amended with 0.58 kg of urea per m³ to stimulate the establishment of nitrifier populations, and 1 kg of Micromax per m³ (Sierra, Milpitas, Calif.). Bark then was treated with 0, 3, or 6 kg of dolomitic lime per m³. This lime contained 51% CaCO₃, 40% MgCO₃, and had a CaCO₃ equivalent of 100. On 24 Jan. 1984, rooted cuttings of *Ilex crenata* Thunb. 'Convexa' were transplanted into 1-liter plastic containers and grown in a glasshouse with day/night temperatures of 24°/18°C. Irrigation techniques and method of medium solution extraction were the same as in experiment 2 of a previous paper (7). Containers were arranged in a randomized complete block design with 3 containers per treatment in each of 5 blocks. Medium solutions were analyzed for pH, NH₄-N, and NO₃-N using ion-selective electrodes.

A set of containers filled with amended pine bark but not containing plants was subjected to the same irrigation schedule and used for NO₃-N accumulation rate (NAR) determinations. Sufficient numbers of containers were prepared to allow for one container per treatment in each of 5 blocks to be periodically sampled.

Periodic determinations of NAR were made following the 2nd nutrient solution irrigation. Following irrigation, containers were allowed to drain for 1 hr, bark was stirred (to ensure uniformity of sampling), and a subsample of bark (about 50 cc) was removed. Containers with remaining bark were enclosed in a plastic bag and stored in an incubator at 25°C for 96 hr. The subsample was put into a PVC tube (2.5 × 15.2 cm) with one end covered with cheesecloth to retain bark during the leaching process. Distilled water (210 ml) was dripped onto bark-filled tubes at a rate of 70 ml·hr⁻¹ and leachates collected (hour 0 N content). This process was repeated 96 hr later with another bark subsample from the same containers. After the leaching process, bark from each tube was dried and weighed. Leachates were analyzed for NH₄-N and NO₃-N. The volume of leachate collected was multiplied by NH₄-N and NO₃-N concentrations to determine the total amounts of leachable NH₄-N and NO₃-N in the bark sample. The total then was divided by the weight of bark to arrive at an amount of N per grams of bark. The

Received for publication 29 Oct. 1985. Research supported in part by the Virginia Nurserymen's Association. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Graduate Research Assistant.

²Professor.

hour 0 amount of $\text{NO}_3\text{-N}$ then was subtracted from the amount at hour 96 and the difference divided by 96 to get a NAR expressed as micrograms of $\text{NO}_3\text{-N}$ per gram of bark per hour. Leachable $\text{NH}_4\text{-N}$ was measured to give an indication of the relative amount of nitrifiable substrate on the bark at each NAR determination.

Experiment 2. NAR was determined every 6 days using bark treated with 6 kg of lime per m^3 and methodology as described previously. At days 42 and 48, some of the containers that had been fertilized prior to day 42 with 100 ppm $\text{NH}_4\text{-N}$ were treated with 300 ppm $\text{NH}_4\text{-N}$. This experiment was conducted to determine if the $\text{NH}_4\text{-N}$ supply of the 100 ppm N treatment limited nitrification over time.

Results and Discussion

Medium solution $\text{NH}_4\text{-N}$ concentrations decreased over time in all treatments; however, the decrease occurred earlier, more rapidly, and to a greater extent in the 6-kg lime treatment compared to the 3-kg lime treatment (Fig. 1). Ammoniacal-N concentrations at 0 kg of lime declined gradually over time. Periods of increasing $\text{NO}_3\text{-N}$ concentration in the medium solution coincided with the decline in $\text{NH}_4\text{-N}$. These periods occurred between days 29 and 70 and days 56 and 84 for the 6- and 3-kg lime treatments, respectively. Nitrate-N was not detected at any time in the 0-kg lime treatment.

Measurable NAR first occurred at days 41 and 84 for the 6- and 3-kg lime treatments, respectively, while $\text{NO}_3\text{-N}$ accumulation did not occur at 0 kg of lime (Fig. 2). NAR was greatest at 6 kg of lime until day 84, after which there was no significant difference between NAR at 3 and 6 kg of lime. The stimulative effect of the 6-kg lime treatment on nitrification is in agreement with the fact that nitrifier growth and metabolism are pH de-




Fig. 2. Influence of lime rate on NAR. Bars indicate SE.

pendent (2). Yet, similar $\text{NO}_3\text{-N}$ concentrations were measured in the medium solution at the 3- and 6-kg treatments between days 70 and 145 (Fig. 1), and no differences in NAR occurred at the last 2 sampling dates in Expt. 2 between these treatments (Fig. 2). The reason for the lack of differences between treatments may be explained by a difference in substrate availability. Medium solution $\text{NH}_4\text{-N}$ concentrations were generally lower during the period of considerable $\text{NO}_3\text{-N}$ accumulation at 6 kg compared to the 3 kg lime treatment.

The amount of leachable $\text{NH}_4\text{-N}$ per gram of bark at hour 0 beyond day 41 and the amount of this N remaining after 96 hr was inversely related to the liming rate (Table 1). Furthermore, only 10% of this hour 0 $\text{NH}_4\text{-N}$ content remained after 96 hr beyond day 56. For example, the percentage of the hour 0 leachable N remaining after 96 hr at day 107 was 90%, 38%, and 12% for the 0-, 3-, and 6-kg lime treatments, respectively. Beyond day 70, the hour 0 leachable $\text{NH}_4\text{-N}$ contents of bark at 6 kg of lime were significantly less than other treatments (Table 1). Thus, relatively low medium solution $\text{NH}_4\text{-N}$ concentrations, relatively low hour 0 $\text{NH}_4\text{-N}$ contents in the NAR determinations, and low amounts of this leachable $\text{NH}_4\text{-N}$ remaining after 96 hr indicate that nitrifier activity at 6 kg of lime apparently was limited by a low $\text{NH}_4\text{-N}$ supply. A nitrifier population exposed to a favorable pH and a limiting $\text{NH}_4\text{-N}$ supply (as with the 6-kg lime treatment) could, over time, produce an amount of $\text{NO}_3\text{-N}$ similar to that produced by a population exposed to a less favorable pH and a greater $\text{NH}_4\text{-N}$ supply (as with the 3-kg lime treatment).

Experiment 2 was conducted to determine if the $\text{NH}_4\text{-N}$ supply of the 100-ppm $\text{NH}_4\text{-N}$ treatment limited nitrification over time. NAR of the 100-ppm N treatment increased until maximum values occurred at days 19 and 25, after which rates de-




Fig. 1. Influence of lime rate on medium solution $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ concentrations. SE = 2.0 if bars are not indicated.

Table 1. Influence of lime treatment on the amount of leachable $\text{NH}_4\text{-N}$ measured at hour 0 and hour 96 during periodic NAR determinations.

Lime treatment (kg lime/m ³ bark)	Days						HSD ²
	41	56	70	84	98	107	
	<i>Hour 0 leachable NH₄-N (μg NH₄-N/g bark)</i>						
0	290	356	362	363	386	324	89
3	275	279	324	292	303	235	64
6	265	278	206	163	142	123	68
HSD	55 ^y	54	66	62	72	74	
	<i>Leachable NH₄-N remaining after 96 hr (μg NH₄-N/g bark)</i>						
0	295	376	362	364	341	291	93
3	280	278	305	275	195	87	57
6	260	264	22	17	13	14	49
HSD	55	90	47	54	51	46	

²Minimum significant difference within rows, HSD 0.05.

^yMinimum significant difference within columns, HSD 0.05.

Table 2. Influence of N treatment on NAR and on the amounts of $\text{NH}_4\text{-N}$ measured at hour 0 and hour 96 during periodic NAR determinations (Expt. 2).

Measurement	N treatment (ppm)	Days									
		1	7	13	19	25	31	37	43	49	HSD ^z
NAR	100	0.21	0.44	1.5	2.1	2.1	1.1	0.81	0.78	0.47	0.51
($\text{NO}_3\text{-N } \mu\text{g}\cdot\text{g}^{-1} \text{ bark}\cdot\text{hr}^{-1}$)	300								2.3 ^{*y}	1.7 [*]	0.33
Hour 0 leachable $\text{NH}_4\text{-N}$	100	229	159	174	141	92	58	42	31	28	21
($\text{NH}_4\text{-N } \mu\text{g}\cdot\text{g}^{-1} \text{ bark}$)	300								114 [*]	104 [*]	20
Leachable $\text{NH}_4\text{-N}$ remaining after 96 hr	100	208	122	150	99	17	7	8	4	3	17
($\text{NH}_4\text{-N } \mu\text{g}\cdot\text{g}^{-1} \text{ bark}$)	300								7	6	4

^zMinimum significant difference within rows, HSD = 0.05.

^y*Means significantly different within columns, *t* test = 0.05.

creased (Table 2). The hour 0 amount of leachable $\text{NH}_4\text{-N}$ decreased from 229 μg of $\text{NH}_4\text{-N}$ per gram of bark at day 1 to 28 μg of $\text{NH}_4\text{-N}$ per gram at day 49 (Table 2). In association with this decrease, the amount of hour 0 $\text{NH}_4\text{-N}$ remaining after 96 hr decreased from 208 μg of $\text{NH}_4\text{-N}$ per gram of bark at day 1 to 17 μg of $\text{NH}_4\text{-N}$ per gram of bark at day 25. Bark previously treated with 100 ppm of $\text{NH}_4\text{-N}$ was treated with 300 ppm of $\text{NH}_4\text{-N}$ at days 43 and 49. NAR increased at least 3-fold in response to the increased N supply. Bark of the 300-ppm N treatment contained at least 3 times the leachable $\text{NH}_4\text{-N}$ at hour 0 as bark of the 100-ppm N treatment. The increased nitrifier activity was evidenced further by the fact that only 6% of the hour 0 leachable $\text{NH}_4\text{-N}$ remained after 96 hr. Thus, results of Expt. 2 support the hypothesis that, over time, nitrifiers were limited by the N supply of the 100-ppm N treatment applied at 6-day intervals.

Medium solution pH decreased over time for all treatments (Fig. 3). Greater decreases in pH occurred during the period of rapid $\text{NO}_3\text{-N}$ accumulation at 3 and 6 kg of lime compared to similar periods at 0 kg of lime. A similar result was found in previous work (7). Nitrification is pH-dependent, with maximum nitrification occurring at pH 7 to 8 (2). Since medium pH decreased over time, the influence of medium pH on nitrification was contingent on the prevailing pH.

This work has characterized the medium solution N and pH status of a pine bark medium treated with different rates of lime. Addition of lime stimulated nitrification; however, this influence varied over time, since medium pH and $\text{NH}_4\text{-N}$ supply decreased over time. Thus nitrifier activity is subject to the

dynamic nature of the medium solution and the influence of a particular lime rate on nitrification is dependent on the set of conditions that are prevalent at any one point in time.

The medium solution $\text{NH}_4\text{-N} : \text{NO}_3\text{-N}$ ratio can have an influence on dry weight accumulation (3, 4, 6), and this influence is apparently species-specific. Nursery plants are sold on the basis of size, hence a ratio that maximizes growth would increase profits. Such a ratio could be effected by manipulating the amount of lime added to the medium. Chrusic and Wright (1) showed growth of azalea, holly, and juniper plants to be greater at 0 or 2 kg of lime per m^3 of bark than at higher rates of lime and attributed this growth response in part to the relatively high $\text{NH}_4\text{-N} : \text{NO}_3\text{-N}$ ratio. A growth response to lime treatment may be in part a response to the ionic form of N as affected by nitrification and not a response to the increased pH or Ca availability. In agreement with this statement, Starr and Wright (8) showed no growth response to medium solution Ca concentrations >10 ppm.

Literature Cited

1. Chrusic, G.A. and R.D. Wright. 1983. Influence of liming on holly, azalea, and juniper growth in pine bark. *J. Amer. Soc. Hort. Sci.* 108:791-795.
2. Focht, D.D. and W. Verstraete. 1977. Biochemical ecology of nitrification and denitrification. *Adv. Microbiol. Ecol.* 1:135-214.
3. Hageman, R.H. 1980. Effect of form of nitrogen on plant growth, p. 47-62. In: M. Stelly (ed.). *Nitrification inhibitors—potentials and limitations*. Amer. Soc. Agron. Spec. Pub. 38.
4. Haynes, R.J. and K.M. Goh. 1978. Ammonium and nitrate nutrition of plants. *Biol. Rev.* 53:465-510.
5. McFee, W.W. and E.L. Stone. 1968. Ammonium and nitrate as nitrogen sources for *Pinus radiata* and *Picea glauca*. *Soil Sci. Soc. Amer. Proc.* 32:879-884.
6. Mengel, K. and E.A. Kirkby. 1982. *Principles of plant nutrition*, 3rd ed. Intl. Potash Inst., Bern, Switzerland.
7. Niemiera, A.X. and R.D. Wright. 1986. Influence of nitrification on the medium solution and growth of azalea, holly, and juniper in a pine bark medium. *J. Amer. Soc. Hort. Sci.* 111:708-712.
8. Starr, K.D. and R.D. Wright. 1985. Calcium and magnesium requirements of *Ilex crenata* 'Helleri'. *J. Amer. Soc. Hort. Sci.* 109:857-860.
9. Townsend, L.R. 1969. Influence of form of nitrogen and pH on growth and nutrient level in the leaves and the roots of lowbush blueberry. *Can. J. Plant Sci.* 49:333-338.




Fig. 3. Influence of lime rate on medium solution pH. SE = 0.2 if bars are not indicated.