

Purification of a Zinc Binding Protein from Xylem of *Citrus jambhiri*

Kathryn C. Taylor,¹ Danielle R. Ellis,² and Luciano V. Paiva³

Department of Plant Sciences, 303 Forbes Building, University of Arizona, Tucson, AZ 85721

ADDITIONAL INDEX WORDS. kunitz soybean proteinase inhibitors, metal-binding proteins, papaya latex trypsin inhibitor, sporamin

ABSTRACT. Zinc in xylem and phloem of the citrus rootstock, rough lemon [*Citrus jambhiri* (L.)] was associated with a Zn-binding protein, designated citrus vascular Zn-binding protein (CVZBP). The apparent molecular mass of the CVZBP was 19.5 kDa after nondenaturing size exclusion chromatography and 21.8 kDa after sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Ion exchange chromatography demonstrated that CVZBP was anionic, requiring 0.43 N NaCl for elution from quaternary aminoethyl Sepharose. Antiserum to the protein cross-reacted more with total protein extracts from leaf midveins than with total protein from the rest of the leaf lamina, further suggesting a vascular location of the Zn-binding protein. Quantitative analysis indicated that ≈ 2 to 3 mol of Zn were associated with 1 mol of native protein. Binding studies with the partially purified CVZBP demonstrated a capacity to bind several divalent cations: Cd, Ni, Pb, and Zn. Reaction with Ellman's reagent suggested that the protein has significant sulfhydryl group content that may be involved in metal binding. N-terminal sequencing demonstrates identity with papaya latex trypsin inhibitor, sporamin, or other Kunitz soybean proteinase inhibitors.

Tree health and longevity appear to be affected by tree Zn status (Marshner, 1995). In addition, Zn status has been associated with citrus blight, a citrus (*Citrus* L. sp.) disorder of unknown cause. This decline disorder is characterized by Zn accumulation in trunk vascular tissues above the bud union (Taylor and Geitzenauer, 1998; Taylor et al., 1988, 1989, 1996; Williams and Albrigo, 1984) and Zn deficiency in the leaves of affected trees. Zinc redistribution is followed by physiological changes that result in decreased water conductivity and tree decline (Albrigo et al., 1986; Brlansky et al., 1984; Vasconcellos and Castle, 1994; Williams and Albrigo, 1984). An abundance of Zn in trunk phloem of decline-affected citrus (Albrigo et al., 1986) is accompanied by accumulation of Zn-binding factors that appear responsible for trunk phloem Zn sequestration (Taylor et al., 1988). One of these Zn-binding factors, the 5 kDa Zn-binding protein (ZBP), was purified and characterized previously (Taylor and Geitzenauer, 1998; Taylor et al., 1996).

While these previous studies were focused on the site of earliest Zn accumulation, the phloem, the greatest Zn accumulation occurs in the xylem of blight affected trees. A Zn-binding factor associated similarly with Zn accumulation in xylem has not been identified previously. Another Zn-binding protein was isolated from citrus vascular tissue (Taylor et al., 1989). The following report is a further characterization of that protein. This novel Zn-binding protein is abundant in xylem and has identity to some members of the Kunitz soybean proteinase inhibitor (KSPI) family.

Materials and Methods

PLANT MATERIAL. As source material for purification of the citrus vascular Zn-binding protein (CVZBP), xylem evacuate

samples were collected from healthy or blight affected trees of *Citrus jambhiri* grown in a quarantine greenhouse facility on the campus of University of Arizona, Tucson, during 1996. Xylem evacuate was collected by applying 4 mL of 10 mM Tris-Cl, pH 7.4 to one end of a 10 cm root piece >1 cm diameter while a vacuum was pulled at the other end. Bark was removed from the root piece down to the cambium before evacuation. Before purification, xylem evacuate was stored at -80°C . Phloem tissue was sampled as a bark patch taken 15 to 30 cm above the soil line. Contaminants on the bark surface were removed by scraping away the brown-green outer layer to a 5×10 cm rectangle of clean whitish phloem tissue. The patch was scored through the scraped bark to the wood and separated from the tree at the cambial layer. The samples were taken from convex surfaces of healthy and blight-affected trees in active growth.

ISOLATION AND PURIFICATION OF CVZBP. To quantitate recovery of CVZBP, three independent purifications were performed. Three separate xylem evacuates (Derrick et al., 1990) were lyophilized and adjusted to 1 mL each with deionized water. Xylem evacuate was fractionated by ion exchange chromatography (IEC) (Taylor et al., 1996) over quaternary aminoethyl Sepharose (QAE) (Pharmacia, Piscataway, N.J.). Total Zn was determined for aliquots of each fraction by determination of $A_{213.9}$ by atomic absorption spectrometry (model 3100, Perkin-Elmer, Shelton, Conn.) and for 254 nm absorbance (A_{254}) by ultraviolet (UV)-spectrophotometry. Fractions with elevations of A_{254} and coincident elevated Zn were pooled and the volume reduced in a centrifugal vacuum concentrator (model RC 10. 10; Jouan, Winchester, Va.). Concentrated Zn-containing eluant from QAE chromatography was separated using a 2.5×14 cm gel filtration (GF) column of Bio-Gel P-30 (Bio-Rad, Hercules, Calif.) in 10 mM Tris-Cl buffer, pH 8.0 (4°C) in 3 mL fractions. Fractions within the peaks of A_{254} and elevated Zn content were pooled, trichloroacetic acid precipitated (Deutscher, 1990), and further purified by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) (Laemmli, 1970). For all isolation steps, total protein was determined at A_{595} with Quantigold (Diversified Biotech, Newton Centre, Mass.) or with the Bradford assay (Bio-Rad), depending upon protein concentration with each stage of purification.

Received for publication 17 Sept. 2001. Accepted for publication 18 Apr. 2002. This research was supported by the Arizona Citrus Research Council.

¹Current address: Department of Horticulture, University of Georgia, Southeastern Fruit and Tree Nut Laboratory, 21 Dunbar Rd., Byron, GA 31008; e-mail kctaylor@uga.edu; fax: 478-956-2929.

²Current address: Children's Nutrition Research Center, 1100 Bates St., Houston TX 77077.

³Current address: Department de Quimica, Universidade Federal de Lavras, UFLA 37200-000, Lavras, MG Brazil.

Table 1. Total protein and Zn, specific Zn content ($\mu\text{g Zn/mg protein}$), and protein concentration during purification of CVZBP from healthy and blight affected tissues of *Citrus jambhiri*.²

Sample ^y	Healthy tissue				Blight-affected tissue			
	Total protein (mg)	Total Zn (μg)	Zn content ($\mu\text{g}\cdot\text{mg}^{-1}$ protein)	Protein concn (mg·g ⁻¹ fresh wt)	Total protein (mg)	Total Zn (μg)	Zn content ($\mu\text{g}\cdot\text{mg}^{-1}$ protein)	Protein concn (mg·g ⁻¹ fresh wt)
Crude extract								
Phloem	30	72.8	0.0024	3.02	30	274.4	0.0091	5.87
Xylem	10	62.1	0.0062	2.03	10	102.0	0.0102	4.49
IEC								
Phloem	6.0	12.3	2.1	0.60	4.3	16.4	3.8	0.84
Xylem	8.5	38.8	4.6	1.85	7.9	34.7	4.4	2.69
GF								
Phloem	0.292	1.5	5.1	0.03	0.271	1.7	6.3	0.05
Xylem	0.425	4.1	9.7	0.13	0.639	6.2	9.6	0.29

²Each value is based on three column purifications.

^yIEC = ion exchange chromatography and GF = gel filtration chromatography.

GF purified CVZBP (50 mg) was electrophoresed for 1.5 h at 150 V, constant voltage, in a pH 8.3 Laemmli buffer system (Laemmli, 1970). Gels were silver stained (Morrissey, 1981). Following SDS-PAGE of samples containing 200 mg protein, two nonstained lanes were sliced across their width into 0.5 cm fractions down the length of the gel. Each gel fraction was eluted in 1.0 mL of 0.5% HCl (Trace metal grade) in high-performance liquid chromatography (HPLC) grade water. Total Zn content eluted from each gel slice was determined by atomic absorption spectrometry. The isoelectric point (pI) for the GF purified CVZBP was determined as described previously (Taylor et al., 1996).

ELLMAN'S DETERMINATION OF SULFHYDRYL GROUPS IN CVZBP. Ellman's reactions (Ellman, 1959) were performed to assess the sulfhydryl content associated with the GF-purified fractions, since they are often involved in protein associations with transition metals (Vallee and Auld, 1990). Fractions eluted from GF columns were assessed for the presence of sulfhydryl containing residues in the CVZBP. This was accomplished by additional postcolumn sulfhydryl group derivatization with 75 μM Ellman's reagent, 5,5'-dithiois (2-nitrobenzoic acid in 50 mM potassium phosphate buffer (pH 7.6). Sulfhydryl determinations were made at 410 nm.

IMMUNOCHEMISTRY. IEC and GF purified CVZBP were separated by preparative SDS-PAGE. Gels were stained in Coomassie Brilliant Blue R in 5 distilled water : 1 acetic acid : 5 methanol (by volume) for 5 min to identify the band of interest. After a 30 to 45 min destain, CVZBP bands were cut out of the gels. The gel pieces were used for anti-serum preparation as described previously (Taylor et al., 1996). Crude phloem extract was used in immunoblot analyses to assess purity of the serum. In addition, leaves were dissected to assess location of the CVZBP with relation to vascular tissue. For all immunoblot analyses, 10 mg of crude protein extract was loaded onto 20% SDS-PAGE gels. Following SDS-PAGE, proteins were transferred to 0.45 μm poly(vinylidene difluoride) membrane in the buffer of Towbin et al. (1979). Electrotransfer was accomplished with the Bio-Rad Mini Trans-Blot apparatus in 45 min. Blots were blocked overnight in 1% bovine serum albumin in buffer [10 mM Tris-Cl, pH 8.0, 150 mM NaCl, 0.05% Tween-20 (TBST)]. Detection was performed by incubating blots in a 1:1000 dilution of anti-CVZBP rabbit serum in TBST followed by incubation in a 1:200 dilution of alkaline

phosphatase labeled goat antirabbit serum (Sigma, St. Louis) in TBST. The immunoblot was developed as in King et al. (1985).

DETERMINATION OF AMINO ACID COMPOSITION AND N-TERMINAL SEQUENCE. For amino acid composition and N-terminal sequencing, a denaturing gel was electroblotted in CAPS buffer (Matsudaira, 1987), lightly stained with Coomassie Brilliant Blue R, and the major band was cut out of the membrane. Amino acid analysis of the CVZBP was completed at the University of Arizona, Laboratory for Protein Sequencing and Analyses, Tucson, using an amino acid analyzer (ABI 420 A/H; Applied Biosystems, Foster City, Calif.). The protein was hydrolyzed in vapor phase using 6 N HCl at 155 °C for 75 min, then derivitized with phenylisothiocyanate to form phenylthiocarbamyl amino acid derivatives which were extracted and transferred to on-line HPLC for analysis of A₂₅₄. Dimethyl sulfoxide (DMSO) hydrolysis was required before amino acid analysis to determine cysteine as cysteic acid (Spencer and Wold, 1969). Results of the 6 N HCl and DMSO hydrolyses were combined for reporting purposes.

Using Edman chemistry, the N-terminal amino acid sequence of the purified CVZBP was determined at the University of Arizona Laboratory for Protein Sequencing and Analysis using a 477A Protein/Peptide sequencer (Applied Biosystems) interfaced with a 120A HPLC (C-18 PTH column, reverse phase chromatography) analyzer to determine PTH amino acids. The N-terminal sequence was analyzed by BLAST searches for identity and similarity with other reported sequences in National Center for Biotechnology Information Genbank.

VOLTAMETRIC DETERMINATION OF RESIDUAL METAL-BINDING CAPACITY OF CVZBP. The voltametric method, differential pulse polarography, was used as a measure of the residual complexation capacity for the divalent cations, Zn, Cd, Ni, and Pb (Street and Petersen, 1982; Taylor et al., 1988). Differential pulse polarography was used to determine the capacity of IEC/GF-purified CVZBP to reduce the level of free metal ion in a 5 mM metal ion solution in 0.1 M KNO₃. To the metal ion solution, 10 mg total protein was added. A polarographic analyzer with a static mercury electrode (model 384B and model 303A, respectively; EG and G Princeton Applied Research, Princeton, N.J.) were used for the assays. Differential pulse polarography was performed over the range of -0.800 to -1.200 V with a drop/step time of 0.6 mV·s⁻¹. Calculations of free metal ion concentration were made on the basis of the total 10 mL assay mixture. A




Fig. 1. Elution profile of protein extracts from QAE-sepharose ion exchange column. Zinc content was coincident with elevated UV_{254nm} of metal-ion chromophores.




Fig. 2. Elution profile of pooled QAE fractions from Bio-Gel P30 column. Zinc containing peaks also absorbed UV at 254 nm. Insert: Calibration curves for the elution of the CVZBP from the Bio-Gel P30 Column. The size exclusion elution profile was calibrated for molecular masses with 3 mg each of soybean trypsin inhibitor (20,100 M_r) and adrenocorticotrophic hormone (ACTH, 4,500 M_r). The void volume was determined by blue dextran (V_o, 2,000,000 M_r) elution.

reduction of the concentration of free metal ion was assumed to be an indication of the level of metal ion complexed by the CVZBP.

Results

ACCOUNT OF ZINC AND PROTEIN THROUGH THE PURIFICATION OF CVZBP. The CVZBP was most abundant in xylem tissue of blight affected citrus. After IEC and GF purification, the CVZBP was present in higher amounts in xylem evacuate than phloem tissue on a total protein basis (1.4- to 2.4-fold more in xylem than phloem) (Table 1). Four to seven times more of the total protein from the xylem evacuate was accounted for by the CVZBP than total protein from phloem (4.3% to 6.4%, xylem; 0.9% to 1.0%, phloem). The ratio of Zn to purified protein indicated that purification of the CVZBP from phloem and xylem was as much as 2125- and 1560-fold, respectively (Table 1). CVZBP was eluted from xylem evacuate of blight affected citrus at 2.2-fold greater level than from xylem evacuate of healthy citrus, while CVZBP elution was similar from phloem tissue extracts of blight and healthy citrus (Table 1). Quantitative analysis of Zn and protein from the three separate purifications of CVZBP from xylem and phloem agreed.

The anionic CVZBP was eluted from QAE Sepharose by 0.43 M NaCl (Fig. 1). Based upon two methods to estimate molecular mass, chromatography and electrophoresis, we placed the apparent molecular mass of this Zn-binding protein at 22 kDa. The Zn peak contained a protein of \approx 19.5 kDa (Fig. 2), as determined by nondenaturing GF chromatography. This relative mobility was less than the apparent molecular mass of 21.8 kDa determined by the denaturing conditions of SDS-PAGE (Fig. 3). Direct preparative isoelectric focusing data indicated that the protein had a pI of 5.5 (data not presented).

DIRECT CORRELATION OF PUTATIVE CVZBP WITH PRESENCE OF ZINC. We were able to directly assess the concentration of Zn in SDS-PAGE gels, after elution from the sequentially dissected gel pieces. Zn was correlated with a 21.8 kDa protein band (Fig. 3), which was the band that cross-reacted with the anti-CVZBP serum. In addition, the Zn-containing preparative isoelectric focusing fractions contained protein that had an apparent molecular mass of 22 kDa (gels not shown).

ELLMAN'S DETERMINATION OF SULFHYDRYL GROUPS IN CVZBP. To assess the sulfhydryl group content of the IEC/GF purified CVZBP, Ellman's reactions (Ellman, 1959) were performed. Assessing only the xylem evacuate from blight affected citrus we determined that the peak of partially purified sample contained \approx 6.7 μ M sulfhydryl groups (Fig. 2). More importantly, the sulfhydryl content was coincident with the Zn and A₂₅₄ elution profiles.

RESIDUAL METAL BINDING CAPACITY OF CVZBP. The capacity of CVZBP to continue to bind Zn under the generally denaturing conditions of SDS-PAGE suggested an extremely stable association of this protein with Zn. Therefore, CVZBP was assessed for its capacity to not only sequester additional levels of Zn, but also Cd, Ni, and Pb above the Zn already bound by the protein, to develop a profile of the capacity of the protein to bind relative quantities of other divalent ions. In this binding study, 1 mg·mL⁻¹ of the IEC/GF purified CVZBP was placed in 5 mM concentrations of each metal individually. The protein bound all metal ions tested at differing molar concentrations (Table 2) in the following sequence Pb > Cd \geq Ni \geq Zn.

IMMUNOCHEMICAL DETERMINATION OF CVZBP ACCUMULATION IN VASCULAR TISSUE. Although CVZBP was extracted from the




Fig. 3. Direct correlation of CVZBP with presence of Zn. Purified CVZBP separated on SDS-PAGE gel adjacent to graph that indicates the level of Zn eluted from 0.5-cm gel slices from the 11 cm gel. Lane 1 = immunoblot of purified CVZBP; lane 2 = silver stained lane from the same gel; and lane 3 = molecular mass markers (Mkr). This gel was overloaded and over stained to show apparent purification of CVZBP.

Table 2. Residual metal-binding capacity of Pb, Ni, Cd, and Zn by GF-purified CVZBP above the level of Zn associated with the protein after its isolation. Five micromolar solutions of each metal was supplied to a $1 \mu\text{g}\cdot\text{mL}^{-1}$ solution of GF-purified CVZBP.

Metal ion	Metal binding (mean \pm SD) ²
Cd	0.79 ± 0.15
Ni	0.65 ± 0.11
Pb	1.15 ± 0.32
Zn	0.55 ± 0.06

²Each mean is based on 10 observations.

phloem and xylem of *C. jambhiri*, it was necessary to determine its specificity to vascular tissues. In an immunochemical determination of CVZBP accumulation, total protein was extracted from whole leaves of *C. jambhiri* and leaves that had been dissected into midveins and remaining laminae to determine if the proteins might be specific to the vascular tissue in the leaf. The CVZBP was most abundant in the more densely vascular midvein extracts and apparently absent in the whole leaf and laminal extracts (Fig. 4). Along with the xylem and phloem isolations of this protein, the midvein localization further suggested a vascular location for CVZBP. When equivalent amounts (10 μg) of IEC/GF purified CVZBP from xylem and phloem were separated by SDS-PAGE followed by immunoblot analysis, they were equally reactive to the anti-CVZBP serum.

AMINO ACID COMPOSITION. Amino acid analysis (Table 3) revealed that CVZBP contained seven cysteine and four histidine residues. These residues are often involved in metal binding of Zn-finger containing proteins (Berg, 1986; Vallee et al., 1991) such as metallothionein. In addition, the composition was rich in aspartate/asparagine and glutamate/glutamine. The acid forms

may participate in metal binding through carboxyl moieties of the R-groups (Vallee and Auld, 1990).

N-TERMINAL AMINO ACID SEQUENCE. The first 27 residues of the amino acid sequence contained no cysteine molecules. However, the sequence was indicative of the N-terminal sequence of sporamin (Hattori et al., 1985), papaya latex trypsin inhibitor (Odani et al., 1996), and other Kunitz-type proteinase inhibitors (Fig. 5), members of the KSPI superfamily of genes. The N-terminal sequence had 52% identity and 67% similarity with the N-terminal sequence of papaya latex trypsin inhibitor, 48% identity and 59% similarity with the N-terminal sequence of sporamin A and B precursors (Hattori et al., 1985, 1989; Koide et al., 1997), and 48% identity with the N-terminal sequence of chymotrypsin inhibitor WCI-3 (Habu et al., 1997; Sakata et al., 1997). The KSPI family includes serine, aspartic and cysteine proteinase inhibitors from a variety of plant families (Gruden et al., 1997; Jokufu and Goldberg, 1989; Leah and Mundy, 1989; Spencer and Hodge, 1991; Trumper et al., 1994), along with other proteins of similar identity but unknown function (Masuda et al., 1995; Theerasilp et al., 1989) such as miraculin.

Discussion

We isolated and characterized a novel CVZBP that appears to be localized primarily in the xylem and is 2.2 times more abundant in xylem of blight affected citrus than xylem of healthy citrus. The CVZBP was associated with the xylem Zn accumulation that characterizes citrus blight. CVZBP has N-terminal identity with papaya latex trypsin inhibitor, sporamin A and B precursors, and chymotrypsin inhibitor, members of the KSPI protein family. The N-terminal amino acid residue of the CVZBP (Fig. 5) was aspartic acid. Lack of an N-terminal methionine residue suggests that CVZBP has a propeptide that is removed after processing. The presence of propeptides has been demonstrated for several members of the KSPI family (Gruden et al., 1997; Koide et al., 1997; Masuda et al., 1995; Spencer and Hodge, 1991).

We estimated the molecular mass of this protein at 22 kDa, despite the molecular mass disparity between nondenaturing and denaturing conditions. This difference with purification conditions is not surprising. Amino acid analysis (Table 3) predicted an intermediate molecular mass of 21.4 kDa for a protein containing 175 amino acid residues. It is likely that the CVZBP was not completely denatured by the typical SDS-PAGE method of Laemmli (1970), since Zn remained associated with the protein throughout SDS-PAGE (Fig. 3), resulting in a size estimate that was larger than the composition suggested.

A key characteristic of members of the KSPI family is the




Fig. 4. Western blot analysis for CVZBP in citrus. (A) Lane 1 = molecular mass marker; lane 2 = positive control, 5 μg of protein extract of trunk phloem tissue; lane 3 to 10 μg of protein extract of leaf midvein; lane 4 = 10 μg of protein extract of whole leaves; and lane 5 = 10 μg of protein extract of leaves without midvein. (B) Lane 1 = 10 μg of protein extract of trunk phloem tissue; and lane 2 = 10 μg of protein xylem evacuate (500 \times concentrate).

Table 3. Amino acid composition of CVZBP.

Amino acid residues	nM (%)	Estimated no. of residues
Asx ^z	12.6	22
Valine	10.8	19
Glycine	9.4	16
Arginine	9.0	16
Serine	8.0	14
Threonine	6.3	14
Glx ^z	6.3	11
Leucine	8.1	11
Alanine	5.1	9
Cysteine	4.1	7
Proline	3.4	6
Isoleucine	3.4	6
Phenylalanine	3.3	6
Lysine	3.3	6
Histidine	2.3	4
Tyrosine	1.7	3
Tryptophan	1.7	3
Methionine	1.2	2
Total	100	175

^zAsx and Glx = asparagine + aspartate and glutamine + glutamate, respectively.

presence of two disulfide domains, one in the N-terminal region and one at the C-terminus. The CVZBP amino acid composition suggested that the sequence contained seven cysteines and four histidines, which can be involved in metal binding. This result was supported by the Ellman's reaction determination (Fig. 2) that demonstrated the presence of sulhydryl groups consistent with the presence of cysteine residues. Several KSPIs contain a similar number of cysteine residues, with four arranged into a domain near the carboxy end of the molecules. Two of these four residues have been demonstrated to form a disulfide bond in miraculin (Igeta et al., 1991). Although some other KSPI members contain additional cysteines, the presence of an additional disulfide bond is demonstrated only in miraculin, which suggests that in at least one member of this family the additional cysteines are crosslinked and this motif is not available to bind metal. However in the CVZBP, it is not improbable that Zn could be stably bound by any of these cysteine pairs replacing the disulfide bonds, with a metal ion crosslinked between them as in the Cd-binding domain of metallothionein. Two KSPI members, the 21 kDa seed protein from *Theobroma cacao* L. (cacao) (Spencer and Hodge, 1991) and miraculin (Igeta et al., 1991), have amino acid sequence identity with the metal binding domains of metallothionein (deMiranda et al., 1990; Evans et al., 1990) and Ferridoxin II (Berg, 1986) relative to cysteine arrangement in the C-terminal third of the protein. In metallothionein, one domain is correlated with sequestration of three metal ions (Vallee and Auld, 1990), similar to the stoichiometry we have noted in CVZBP purification. In fact our Zn/CVZBP quantitation (gel filtration, Table 1) suggests Zn:CVZBP molar ratios in this range (with 1.77 and 3.33 for phloem and xylem purified CVZBP, respectively). Differential pulse polarography data (Table 2) for GF-purified metal binding fractions

from citrus phloem tissue suggested that this protein bound Zn in excess of that found associated with these fractions in vivo. Like metallothionein, the CVZBP also had the capacity to bind Cd, as well as Pb and Ni.

Although a highly speculative conclusion at this juncture, we may reasonably hypothesize that isolation and characterization of the gene that encodes CVZBP would predict a deduced amino acid sequence with a cysteine content and arrangement typical of other metal binding proteins like metallothionein, yet similar to the sequences of other KSPI family members, such as miraculin and cacao seed protein (with sequence similarity to the Cd-binding domain of metallothionein). Whether the primary function of this protein is that of metal sequestration we cannot predict. That conclusion will require much functional analysis. We can suggest that CVZBP is a novel metal-binding protein with N-terminal sequence consistent with members of the KSPI family that appear to function in plant defense and possibly development. While the primary function of CVZBP is yet uncharacterized, the vascular location of CVZBP does not eliminate the possibility of any of these suggested functions.

Literature Cited

Albrigo, L.G., J.P. Syvertsen, and R.H. Young. 1986. Phloem zinc accumulation in citrus trees affected with blight. *HortScience* 16:158-160.
 Berg, J.M. 1986. Potential metal-binding domains in nucleic acid binding proteins. *Science* 232:485-486.
 Brlansky, R.H., Jr., J.B. Hollick, T.J. Parsons, H.R.G. Clark, and M.P. Gordon. 1984. Relationship of xylem plugging to reduced water uptake and symptom development in citrus trees with blight and blight like declines. *Phytopathology* 74:1325-1328.
 de Miranda, J.R., M.A. Thomas, D.A. Thurman, and A.B. Tomsett. 1990. Metallothionein of cDNAs associated with tracheary element differentiation in cultured zinnia cells. *Plant Physiol.* 103: 815-821.
 Derrick, K.S., R.F. Lee, R.H. Brlansky, L.W. Timmer, B.G. Hewitt, and G.A. Barthe. 1990. Proteins associated with citrus blight. *Plant Dis.* 74:168-170.
 Deutscher, M.P. 1990. Guide to protein purification. Academic Press, Inc., San Diego.
 Ellman, G.L. 1959. Tissue sulhydryl groups. *Arch. Biochem. Biophys.* 82:70-77.
 Evans, I.M., L.N. Gatehouse, J.A. Gatehouse, N.J. Robinson, and R.R.D. Croy. 1990. A gene from pea, *Pisum sativum* L., with homology to metallothionein genes. *FEBS Lett.* 262:29-32.
 Gruden, K., B. Strukelj, M. Ravniker, M. Poljsak-Prijatelj, I. Mavric, J. Brzin, J. Pungercarn, and I. Kregar. 1997. Potato cysteine proteinase

Fig. 5. N-terminal amino acid sequence of purified CVZBP.

Protein	Sequence	% ID/ % SIM	Swiss-Prot database no.
CVZBP	-DPLVDVHGN KVEASRDYYL VSAIRGAG	100	
Papaya LTI	PK·I·ID·K P·LYGV·FV ···W···	52/67	P80691
Sporamin B	--VL·IN·D ER·GEN·M- ···W···	48/59	P14716
Sporamin A	--VL·IN·D ER·GGN·M- ···W···	48/59	P14715
Trp Inh DE3	---·LE·L·NGGT·LPH·WAL·	48/56	P07475
Trp Inh DE5	---L·D·F LNRGGG·I·P·F·K·	44/56	P09941
WBA1bumin-1	--VY·AE·LVNRGK·TI·FSD···	44/56	P15465
K trp inh	-RE·L·D·F LNRGGG·I·P·F·K·	44/56	P25272
Cocoa Seed	--VL·TD·D ELQTNQ·V L·S·S···	37/56	P32765
Wh α-Am Inh	--VH·TD·E LR·DAN·V LP·N·AH·	37/52	P16347
STI3	---VL·NE·PL·NGGT·I L·D·TAF·	33/52	P01070
Miraculin	-N·VL·ID·E ·LRTGTN·I ·PVL·DH·	33/48	P13087

- inhibitor gene family: Molecular cloning, characterization and immunocytochemical localization studies. *Plant Mol. Biol.* 34:317–323.
- Habu, Y., S. Peyachoknagul, Y. Sakata, F. Fukasawa, and T. Ohno. 1997. Evolution of a multigene family that encodes the Kunitz chymotrypsin inhibitor in winged bean: A possible intermediate in the generation of a new gene with a distinct pattern of expression. *Mol. Gen. Genet.* 254:73–80.
- Hattori, T., T. Nakagawa, M. Maeshima, K. Nakamura, and T. Asahi. 1985. Molecular cloning and nucleotide sequence of cDNA for sporamin, the major soluble protein of sweet potato tuberous roots. *Plant Mol. Biol.* 5:313–320.
- Hattori, T., N. Yoshida, and K. Nakamura. 1989. Structural relationship among the members of a multigene family coding for the sweet potato tuberous root storage protein. *Plant Mol. Biol.* 13:563–572.
- Igeta, H., Y. Tamura, K. Nakaya, Y. Nakamura, and Y. Kurihara. 1991. Determination of disulfide array and subunit structure of taste-modifying protein, miraculin. *Biochem. et Biophys. Acta.* 1079:303–307.
- Jofuku D.K. and R.B. Goldberg. 1989. Kunitz trypsin inhibitor genes are differentially expressed during the soybean life cycle and in transformed tobacco plants. *Plant Cell* 1:1079–1093.
- King, S.M., T. Otter, and G.B. Witman. 1985. Characterization of monoclonal antibodies against *Chlamydomonas* flagellar dyneins by high resolution protein blotting. *Proc. Natl. Acad. Sci. USA* 82:4717.
- Koide, Y., H. Hirano, K. Matsuoka, and K. Nakamura. 1997. The N-terminal propeptide of the precursor to sporamin acts as a vacuole-targeting signal even at the C terminus of the mature part in tobacco cells. *Plant Physiol.* 114:863–870.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:68–685.
- Leah, R. and J. Mundy. 1989. The bifunctional α -amylase/subtilisin inhibitor of barley: Nucleotide sequence and patterns of seed-specific expression. *Plant Mol. Biol.* 12:673–682.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. 2nd ed. Academic Press, Inc., San Diego.
- Masuda, Y., S. Nirasawa, K. Nakaya, and Y. Kurihara. 1995. Cloning and sequencing of a cDNA encoding and taste-modifying protein, miraculin. *Gene* 161:175–177.
- Matsudaira, P. 1987. Sequence from picomole quantities of proteins electroblotted onto polyvinylidene difluoride membranes. *J. Biol. Chem.* 262:10035–10038.
- Morrissey, J.H. 1981. Silver stain for proteins in polyacrylamide gels: A modified procedure with enhanced uniform sensitivity. *Anal. Biochem.* 117:307–310.
- Odani, S., Y. Yokokawa, H. Takeda, S. Abe, and S. Okani. 1996. The primary structure and characterization of carbohydrate chains of the extracellular glycoprotein proteinase inhibitor from latex of *Carica papaya*. *European J. Biochem.* 241:77–82.
- Sakata, Y., Y. Chiba, H. Fukushima, N. Matsubara, Y. Habu, S. Naito, and T. Ohno. 1997. The RY sequence is necessary but not sufficient for the transcription activation of a winged bean chymotrypsin inhibitor gene in developing seeds. *Plant Mol. Biol.* 34:191–197.
- Spencer, M.E. and R. Hodge. 1991. Cloning and sequencing of the cDNA encoding the major albumin of *Theobroma cacao*. *Planta* 183:528–535.
- Spencer, R.L. and F. Wold. 1969. A new convenient method for estimation of total cystine-cysteine in proteins. *Anal. Biochem.* 32:185.
- Street, J.J. and W.M. Petersen. 1982. Anodic stripping and differential pulse polarography, p.112–141. In: A.L. Page, R.H. Miller, and D.R. Keeney (eds.). *Methods of soil analysis: Part 2. Chemical and microbiological properties*. 2nd ed. Amer. Soc. of Agron., Soil Sci. Soc. of Amer., Madison, Wis.
- Taylor, K.C., L.G. Albrigo, and C.D. Chase. 1988. Zinc complexation in the phloem of blight-affected citrus. *J. Amer. Soc. Hort. Sci.* 113:407–411.
- Taylor, K.C., L.G. Albrigo, and C.D. Chase. 1989. Characterization of a Zn-binding peptide associated with a decline disorder in citrus, p. 385–394. In: D. Winge and D. Hamer (eds.). *UCLA symposium on metal ion homeostasis: Molecular biology and chemistry*. Alan R. Liss, Inc., New York.
- Taylor, K.C., L.G. Albrigo, and C.D. Chase. 1996. Purification of a Zn-binding phloem protein with sequence identity to chitin-binding proteins. *Plant Physiol.* 110:657–664.
- Taylor, K.C. and H.L. Geitzenauer. 1998. 5-kD zinc-binding protein accumulation in macrophylla-decline-affected citrus. *J. Amer. Soc. Hort. Sci.* 123:357–360.
- Theerasilp, S., H. Hitotsuya, S. Nakajos, K. Nakayas, Y. Nakamuras, and Y. Kurihara. 1989. Complete amino acid sequence and structure characterization of the taste-modifying protein, miraculin. *J. Biol. Chem.* 264:6655–6659.
- Towbin, H., T. Staehlin, and T. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. *Proc. Natl. Acad. Sci. USA* 76:4350–4354.
- Trumper, S., H. Follmann, and I. Haberlein. 1994. A novel dehydroascorbate reductase from spinach chloroplasts homologous to plant trypsin inhibitor. *FEBS Lett.* 352:159–162.
- Vallee, B. and D. Auld. 1990. Zinc coordination, function and structure of zinc enzymes and other proteins. *Biochemistry* 29:5647–5659.
- Vallee, B.L., J.E. Coleman, and D.S. Auld. 1991. Zinc fingers, zinc clusters and zinc twists in DNA binding domains. *Proc. Natl. Acad. Sci. USA* 88:999–1003.
- Vasconcellos, L.A.B.C. and W.S. Castle. 1994. Trunk xylem anatomy of mature healthy and blighted grapefruit trees of several rootstocks. *J. Amer. Soc. Hort. Soc.* 119:185–194.
- Williams, G.L. and L.G. Albrigo, 1984. Some inorganic element changes in trunk phloem of healthy and blight-affected citrus trees. *J. Amer. Soc. Hort. Sci.* 109:437–440.